



AJKPSC

Updated
Edition!
2026

BIOLOGY

Lecturer Notes


According to the Latest Syllabus of Azad Jammu & Kashmir Public Service Commission Exams

Also Helpful For Following Exams:

- *ESE, SST & TGT*
- *Subject Specialist*
- *Assistant Professor*

Premium Features:

- To The Point Topic Wise Detailed Notes
- Authentic, Concise, Error less & Result Oriented Notes + One Liners + Practice MCQs
- Solved Past Papers & Most Repeated Important MCQs
- According to Azad Jammu & Kashmir Public Service Commission Syllabus
- 100% Success Guaranteed



BY

MUBASHIR AHMAD
Lecturer Biology (HED)

MK PUBLICATIONS
LET'S MAKE IT HAPPEN!

For Online Order:
0348-4233593

Contact: 0342-4470091 | 0333-2605045

Email: mkpreparations@gmail.com



@MK PREPARATIONS



Table Of Content

1. Introduction to Biology	15. Kingdom Plantae
2. Cell Biology	16. Forms and Functions in Plants
3. Cell Cycle	17. Kingdom Animalia
4. Biological Molecules & Enzymes	18. Nutrition and Digestion
5. Bioenergetics	19. Gaseous Exchange
6. Chromosomes & DNA	20. Circulation & Transport
7. Variation & Genetics	21. Homeostasis
8. Biotechnology	22. Support & Movement
9. Evolution	23. Nervous and Sensory System
10. Ecology and Ecosystems	24. Endocrine System
11. Viruses	25. Immune System
12. Bacteria & Archaea	26. Reproduction and Development
13. Protists	27. Solved Past Papers
14. Fungi	





Chapter 1

Introduction to Biology

Biology is the scientific study of life. The term originates from the Greek words *bios* (life) and *logos* (study/discourse). It encompasses the investigation of all living organisms—their structure, function, growth, origin, evolution, and distribution—as well as their interactions with each other and the non-living environment. A biologist studies both the **living (biotic)** components of nature and the **non-living (abiotic)** factors that influence them.

Life is defined operationally by a set of distinctive characteristics that collectively distinguish living organisms from non-living matter. These characteristics are not exclusive to any one organism but are a shared basis for life as we know it.

Characteristics of Life

All living organisms exhibit the following key characteristics:

- **Cellular Organization:** All living things are composed of one or more **cells**, which are the basic structural and functional units of life.
- **Metabolism:** Living organisms acquire and utilize energy through a complex set of controlled chemical reactions. This includes **anabolism** (building complex molecules) and **catabolism** (breaking down molecules).
- **Homeostasis:** Organisms maintain a relatively stable internal environment despite changes in the external conditions.
- **Growth and Development:** Organisms increase in size (**growth**) and undergo orderly, genetically programmed changes (**development**) throughout their lifespan.
- **Reproduction:** Living things produce new individuals of their own kind, ensuring the continuity of their species. This can be asexual (involving one parent) or sexual (involving two parents).
- **Heredity and Variation:** Genetic material (**DNA**) is passed from parents to offspring, transmitting traits (**heredity**). Slight changes in this material lead to **variation**, which is the raw material for evolution.
- **Response to Stimuli (Sensitivity/Sensitivity):** Organisms detect and respond to changes (stimuli) in their internal and external environments.
- **Adaptation:** Over generations, populations of organisms evolve traits that enhance their survival and reproduction in a specific environment.

Historical Development of Biology

The study of life has evolved through significant contributions:

- **Ancient Foundations:** Early knowledge from civilizations for medicine, agriculture, and animal domestication.
- **Aristotle (4th Century BCE):** Conducted systematic observation and classification of animals; regarded as the **Father of Zoology**.
- **Theophrastus:** A student of Aristotle, recognized as the **Father of Botany** for his foundational work on plants.
- **Carl Linnaeus (18th Century):** Formally used the term "Biology" and established the **binomial nomenclature** system for naming species.
- **Robert Hooke (1665):** Coined the term "**cell**" after observing cork tissue under a microscope.
- **Antonie van Leeuwenhoek (1670s):** First to observe and describe living microorganisms ("animalcules") using improved microscopes.
- **Matthias Schleiden (1838) & Theodor Schwann (1839):** Formulated the foundational **Cell Theory**, stating that all plants and animals are composed of cells.
- **Rudolf Virchow (1855):** Added the principle "*Omnis cellula e cellula*" (all cells arise from pre-existing cells), completing the classical Cell Theory.



Introduction to Biology: One Liner

- **Biology** is the scientific study of life.
- The term Biology originates from the Greek words *bios* (life) and *logos* (study/discourse).
- **Carl Linnaeus** first formally used the term "Biology" in 1736.
- **Lamarck and Treviranus** made the term "Biology" common in the 1800s.
- Biology encompasses the study of structure, function, growth, origin, evolution, and distribution of living organisms.
- Biology also studies interactions between organisms and with their non-living environment.
- A biologist studies both **biotic** (living) and **abiotic** (non-living) components of nature.
- **Jabir Bin Hayan (721-815 AD)** pioneered experimental investigation in chemistry.
- **Abdul Malik Asmai (740-828 AD)** is considered the first Muslim scientist to conduct detailed anatomical and behavioral studies of animals.
- **Bu Ali Sina (Avicenna) (980-1037 AD)** is honored as the founder of medicine.
- Bu Ali Sina's book "**Al-Qanun-fi-al-Tib**" is a monumental medical encyclopedia used for centuries.
- Life is defined operationally by a set of shared characteristics.
- **Cellular Organization:** All living things are composed of one or more **cells**.
- **Metabolism:** Living organisms acquire and utilize energy through controlled chemical reactions.
- Metabolism includes **anabolism** (building complex molecules) and **catabolism** (breaking down molecules).
- **Homeostasis:** Organisms maintain a relatively stable internal environment.
- **Growth:** Organisms increase in size.
- **Development:** Organisms undergo orderly, genetically programmed changes.
- **Reproduction:** Living things produce new individuals of their own kind.
- Reproduction can be **asexual** (one parent) or **sexual** (two parents).
- **Heredity:** Genetic material (**DNA**) is passed from parents to offspring.
- **Variation:** Slight changes in genetic material lead to differences, the raw material for evolution.
- **Response to Stimuli:** Organisms detect and respond to changes in their internal and external environments.
- **Adaptation:** Over generations, populations evolve traits that enhance survival and reproduction.
- **Aristotle (4th Century BCE)** is regarded as the **Father of Zoology**.
- **Theophrastus**, a student of Aristotle, is recognized as the **Father of Botany**.
- **Robert Hooke (1665)** coined the term "**cell**" after observing cork tissue.
- **Antonie van Leeuwenhoek (1670s)** was the first to observe and describe living microorganisms.
- **Matthias Schleiden (1838) & Theodor Schwann (1839)** formulated the foundational **Cell Theory**.
- **Rudolf Virchow (1855)** added that all cells arise from pre-existing cells ("**Omnis cellula e cellula**").
- **Charles Darwin & Alfred Russel Wallace (1859)** independently proposed the **Theory of Evolution by Natural Selection**.
- **Watson & Crick (1953)** discovered the double helix structure of DNA.
- The **Human Genome Project** was completed in 2003.
- The three major divisions are **Zoology**, **Botany**, and **Microbiology**.
- **Zoology** is the study of animals.
- Sub-branches of Zoology include **Entomology** (insects), **Ichthyology** (fish), **Ornithology** (birds), **Mammalogy**, and **Herpetology**.
- **Botany** is the study of plants.

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. The term **Biology** is derived from Greek words "bios" and "logos" meaning:

- A) Animal and discourse
- B) Life and study
- C) Plant and logic
- D) Nature and science

Answer: Life and study

2. The study of internal structure of organisms is called:

- A) Morphology
- B) Histology
- C) Anatomy
- D) Physiology

Answer: Anatomy

3. Who is known as the **Father of Zoology**?

- A) Theophrastus
- B) Aristotle
- C) Carl Linnaeus
- D) Robert Hooke

Answer: Aristotle

4. The branch of biology that deals with the study of insects is:

- A) Ichthyology
- B) Herpetology
- C) Ornithology
- D) Entomology

Answer: Entomology

5. The percentage of oxygen by mass in the human body is approximately:

- A) 18%
- B) 65%
- C) 10%
- D) 3%

Answer: 65%

6. Which of the following is **NOT** a characteristic of life?

- A) Metabolism
- B) Homeostasis
- C) Crystallization
- D) Reproduction

Answer: Crystallization

7. Who formulated the Cell Theory along with Theodor Schwann?

- A) Rudolf Virchow
- B) Matthias Schleiden
- C) Robert Hooke
- D) Antonie van Leeuwenhoek

Answer: Matthias Schleiden

8. The basic structural and functional unit of life is the:

- A) Tissue
- B) Organ
- C) Cell
- D) Organelle

Answer: Cell

9. The study of fossils is known as:

- A) Taxonomy
- B) Palaeontology
- C) Embryology
- D) Histology

Answer: Palaeontology

10. Which bioelement accounts for about 18% of human body mass?

- A) Hydrogen
- B) Nitrogen
- C) Carbon
- D) Oxygen

Answer: Carbon

11. A group of similar cells performing a specific function is called:

- A) Organ
- B) Organ system
- C) Tissue
- D) Organism

Answer: Tissue

12. Who first observed and described living microorganisms?

- A) Robert Hooke
- B) Carl Linnaeus
- C) Antonie van Leeuwenhoek
- D) Louis Pasteur

Answer: Antonie van Leeuwenhoek

13. The branch of biology that studies the interactions between organisms and their environment is:

- A) Genetics
- B) Ecology
- C) Physiology
- D) Embryology

Answer: Ecology

14. "Omnis cellula e cellula" was stated by:

- A) Rudolf Virchow
- B) Charles Darwin
- C) Louis Pasteur
- D) Gregor Mendel

Answer: Rudolf Virchow

Chapter 2

Cell Biology

Introduction

Cell biology (cytology) is the interdisciplinary study of cell structure, function, and behavior, integrating microscopy, biochemistry, genetics, and computational biology. The **cell** is the smallest unit capable of performing all activities associated with life and serves as the fundamental **structural and functional unit** of all living organisms. While the components of a cell cannot survive independently, the cell operates as an integrated system, and groups of cells form tissues, organs, and organisms, demonstrating biological complexity.

Cell Theory

The **Cell Theory** is the foundational principle of modern biology, formalized by **Matthias Schleiden** (1838), **Theodor Schwann** (1839), and **Rudolf Virchow** (1855). **August Weismann** later contributed the concept of common descent.

Salient Features of Modern Cell Theory:

- All living organisms are composed of one or more cells.
- The cell is the basic unit of structure, function, and organization in all organisms.
- All cells arise from pre-existing cells through cell division (**Principle of Biogenesis**).
- Cells contain hereditary material (DNA) passed from parent to daughter cells.
- All cells are fundamentally similar in chemical composition and core metabolic processes.
- The total activity of an organism is the sum of the activities of its independent cells.

Historical Development and Validation of Cell Theory

- **Robert Hooke (1665)**: Coined the term "cell" after observing the honeycomb structure of dead cork tissue under a primitive microscope. He observed cell walls, not living content.
- **Anton van Leeuwenhoek (1670s)**: First to observe and describe living, moving cells ("animalcules") like bacteria, protozoa, and spermatozoa using superior, self-made lenses.
- **Robert Brown (1831)**: Discovered the **nucleus** within plant cells, identifying a constant internal structure.
- **Matthias Schleiden (1838) & Theodor Schwann (1839)**: Formally proposed the Cell Theory. Schleiden concluded plants are composed of cells, and Schwann extended this to animals, stating cells are the universal building blocks of life.
- **Rudolf Virchow (1855)**: Articulated "**Omnis cellula e cellula**" ("All cells come from cells"), refuting the theory of spontaneous generation for cellular life.
- **Louis Pasteur (1862)**: Provided definitive experimental proof for biogenesis using his famous swan-neck flask experiment, demonstrating that microorganisms arise only from pre-existing ones.
- **August Weismann (1880)**: Added that all living cells trace their ancestry back to ancient cells, establishing the principle of **common descent**, supported by universal biochemical similarities.

Advanced Microscopy: Principles and Techniques

Microscopy overcomes the limited **resolution** of the human eye (~0.1 mm). **Resolution (Resolving Power)** is the minimum distance between two points that can be distinguished as separate. **Magnification** is the increase in an object's apparent size. High magnification without sufficient resolution results in empty magnification (a blurry, enlarged image).

Light Microscopy (LM)

Utilizes visible light (400-700 nm wavelength) and a system of glass lenses.

- **Maximum Theoretical Resolution**: ~200 nm (0.2 μm), limited by the wavelength of light (Abbe's Law).
- **Maximum Useful Magnification**: ~1000X to 1500X.
- **Key Techniques**:

- **Bright-field Microscopy:** Specimen is illuminated from below. Requires staining for contrast, which usually kills cells.
- **Phase-contrast & DIC (Differential Interference Contrast):** Enhance contrast in **unstained, living cells** by converting differences in refractive index and thickness into variations in brightness.
- **Dark-field Microscopy:** Illuminates the specimen at an angle; only light scattered by the specimen enters the lens, creating a bright image on a dark background. Excellent for observing motility.
- **Fluorescence Microscopy:** Uses specific **fluorophores** (e.g., GFP, fluorescent dyes, antibody tags) that absorb high-energy light and emit lower-energy light. Allows localization of specific molecules. **Confocal microscopy** uses a laser and pinhole to create sharp, optical sections and 3D reconstructions, eliminating out-of-focus blur.
- **Super-resolution Microscopy** (e.g., STED, PALM, STORM): Breaks the diffraction limit of light, achieving resolutions down to 10-20 nm, allowing visualization of single molecules.

Electron Microscopy (EM)

Uses a beam of electrons (wavelength ~0.005 nm) focused by electromagnetic lenses, offering vastly superior resolution.

- **Maximum Resolution:** ~0.2 nm (theoretical) to 2 nm (practical for biological specimens).
- **Key Techniques:**
 - **Transmission EM (TEM):** Electrons pass through an ultra-thin section (50-100 nm) of a specimen stained with heavy metals (e.g., lead, uranium). Reveals **internal ultrastructure** (organelle details, macromolecular complexes).
 - **Scanning EM (SEM):** An electron beam scans the surface of a specimen coated with a thin metal film. Detects emitted secondary electrons to produce a detailed, **3D-like topographical image** of surfaces.
 - **Cryo-Electron Microscopy (Cryo-EM):** Specimens are rapidly frozen (vitrified) in liquid ethane, preserving them in a near-native, hydrated state without chemical fixation. Enables atomic-level structure determination of proteins and complexes (Nobel Prize 2017).

Light vs. Electron Microscope

Feature	Light Microscope (LM)	Electron Microscope (EM)
Illuminating Source	Visible Light	Beam of Electrons
Max Magnification	~1500X	Up to 1,000,000X+
Max Resolution	~200-250 nm	~0.2 nm
Specimen Status	Live or dead	Always dead, fixed, dehydrated
Image Color	Colored (if stained)	Monochrome (false color added digitally)
Vacuum Required	No	Yes
Primary Use	Living cells, tissues, large organelles	Ultrastructure, viruses, macromolecules

Techniques used in Cell Biology

- **Cell Fractionation & Centrifugation:** Separates cellular components based on size and density. Tissue is **homogenized** and then spun at high speeds in a **centrifuge**. **Differential centrifugation** yields pellets enriched with specific organelles (e.g., nuclei, mitochondria). **Density gradient centrifugation** provides finer separation.
- **Differential Staining:** Uses specific dyes (e.g., haematoxylin for nuclei, eosin for cytoplasm) to contrast different cellular structures.
- **Tissue Culture:** Growth of cells or tissues on a sterile, defined nutrient medium. Used for cloning, studying cell behavior, and cancer research.

- **Chromatography:** Separates chemical mixtures (e.g., pigments, proteins) based on differential solubility in mobile and stationary phases.
- **Electrophoresis:** Separates charged molecules (DNA, RNA, proteins) based on size and charge in a gel under an electric field.
- **Spectrophotometry:** Measures the absorption of specific light wavelengths by a compound, used to identify substances and determine concentrations.
- **Micrometry:** Measures microscopic objects using an **ocular micrometer** calibrated with a **stage micrometer**.

M K P R E P A R A T I O N S

Universal Features of All Cells

All cells, prokaryotic and eukaryotic, share four fundamental organizational features:

1. **Plasma Membrane:** A **phospholipid bilayer** that forms a selective barrier, regulating exchange and maintaining homeostasis.
2. **Cytoplasm:** The semi-fluid interior (**cytosol**) containing ions, metabolites, and organelles.
3. **Genetic Material:** **DNA** (deoxyribonucleic acid) as the hereditary information.
4. **Ribosomes:** **Ribonucleoprotein** complexes (rRNA + protein) that synthesize proteins.


Prokaryotic vs. Eukaryotic Cells: A Fundamental Divide

Life is classified into three domains: **Bacteria** and **Archaea** (prokaryotes), and **Eukarya** (eukaryotes: Protists, Fungi, Plants, Animals).


Differences: Bacteria vs. Archaea

Feature	Bacteria	Archaea
Cell Wall	Contains peptidoglycan	No peptidoglycan; diverse composition
Membrane Lipids	Ester-linked fatty acids to glycerol	Ether-linked hydrocarbons; can form lipid monolayers for thermal stability
Genetic Machinery	Distinct	More similar to eukaryotes (e.g., RNA polymerase, transcription factors)
Flagella	Bacterial Flagellum: Hollow filament of flagellin; rotary motor powered by proton (H^+) gradient.	Archaeum: Structurally distinct, related to type IV pili; powered by ATP hydrolysis . (Example of convergent evolution)


Prokaryotic cell



Plant Cell



Animal Cell



Prokaryotic vs. Eukaryotic Cells

Feature	Prokaryotic Cell (Bacteria/Archaea)	Eukaryotic Cell (Protists, Fungi, Plants, Animals)
Nucleus	Absent. DNA in a nucleoid region (not membrane-bound).	Present. DNA enclosed within a double-membrane nuclear envelope .
Membrane-Bound Organelles	Absent. (e.g., No mitochondria, ER, Golgi).	Present. Extensive compartmentalization (e.g., ER, Golgi, lysosomes, peroxisomes).
Cell Size	Generally small (0.5 – 5.0 μm).	Generally larger (10 – 100 μm).
Cytoskeleton	Primitive. Contains homologs of actin (MreB) and tubulin (FtsZ) for shape and division, but not a complex network.	Present. Complex, dynamic network of microtubules, microfilaments, and intermediate filaments.
Ribosomes	70S (50S + 30S subunits).	80S in cytosol (60S + 40S). 70S in mitochondria & chloroplasts.
Cell Division	Binary Fission. Simple division after DNA replication.	Mitosis (for growth/repair) or Meiosis (for gamete formation).
DNA Form	Single, circular chromosome. May have small circular plasmids.	Multiple, linear chromosomes complexed with histone proteins to form chromatin.
Flagella Structure	Simple, composed of flagellin protein; rotates like a propeller driven by a proton gradient.	Complex, composed of microtubules in a " 9+2 " axoneme pattern; bends via ATP-driven dynein motor proteins.
Cell Wall Composition	Bacteria: Peptidoglycan (sugar-polymer cross-linked by peptides). Archaea: Varied (pseudopeptidoglycan, polysaccharides, glycoprotein S-layers).	Plants & Algae: Cellulose. Fungi: Chitin. Animals: No cell wall.

The Eukaryotic Cell: Organelle Structure and Function

The Nucleus: Command Center

- **Discovery:** Robert Brown (1831).
- **Structure:**
 - **Nuclear Envelope:** Double phospholipid bilayer (inner and outer nuclear membrane). The outer membrane is continuous with the **Rough ER** and is studded with ribosomes. The **perinuclear space** lies between the two membranes.
 - **Nuclear Pore Complexes (NPCs):** Massive, octagonal protein structures spanning the envelope. They regulate **active, signal-dependent transport** of macromolecules (e.g., mRNAs out, ribosomal proteins in) while allowing passive diffusion of small molecules.
 - **Nuclear Lamina:** A meshwork of **intermediate filament proteins (lamins A, B, C)** lining the inner nuclear membrane. Provides structural support, organizes chromatin, and disassembles/reassembles during mitosis.
 - **Nucleoplasm:** The semi-fluid matrix within the nucleus.
 - **Chromatin:** The DNA-protein complex. **Euchromatin** is decondensed, genetically active. **Heterochromatin** is highly condensed, transcriptionally silent. Histone modifications and DNA methylation regulate this **epigenetic** state.

- **Nucleolus:** A dense, **membrane-less organelle** formed via liquid-liquid phase separation. Site of **rRNA transcription** by RNA Pol I and **assembly of ribosomal subunits** (60S and 40S). Larger in cells active in protein synthesis.
- **Function:** Stores genetic material (DNA), coordinates cellular activities (via gene expression regulation), site of DNA replication and transcription (RNA synthesis), and ribosome biogenesis.

Ribosomes: Protein Synthesis Machinery

- **Discovery:** George Palade (1950s) using EM.
- **Structure:** Composed of **rRNA (ribosomal RNA)** and proteins. A **ribonucleoprotein** complex with two subunits that assemble only during translation. Eukaryotic ribosomes are **80S** (comprising a large 60S and small 40S subunit).
- **Types & Location:**
 - **Free Ribosomes:** Suspended in cytosol. Synthesize soluble proteins that function within the cytosol (e.g., glycolytic enzymes, cytoskeletal proteins).
 - **Bound Ribosomes:** Attached to the cytosolic face of the Rough ER (or outer nuclear membrane). Synthesize proteins destined for secretion, incorporation into membranes, or targeted to specific organelles (lysosomes, Golgi).
- **Polyribosomes (Polysomes):** A single mRNA molecule being simultaneously translated by multiple ribosomes, greatly increasing the efficiency of protein production.

The Endomembrane System: Synthesis, Trafficking, and Degradation

A functionally interconnected network of organelles that includes the nuclear envelope, ER, Golgi, lysosomes, vesicles, vacuoles, and plasma membrane. Materials shuttle between components via **transport vesicles**.

1. Endoplasmic Reticulum (ER):

- **Discovery:** Keith Porter (1945) via EM.
- **Structure:** An extensive network of membranous tubules and flattened sacs called **cisternae**. The internal compartment is the **ER lumen** or cisternal space.
- **Rough ER (RER):** Studded with **bound ribosomes**. Appears "rough" in EM.
 - **Functions:** Synthesis of secretory, membrane-bound, and organellar proteins. Initial **post-translational modification** (e.g., **N-linked glycosylation**). Proper protein folding facilitated by **chaperones** (e.g., BiP). Site of **phospholipid synthesis** for membrane expansion.
- **Smooth ER (SER):** Lacks ribosomes; more tubular.
 - **Functions:** **Synthesis of lipids** (steroid hormones, phospholipids, oils). **Detoxification** of drugs and poisons (liver hepatocytes have abundant SER containing cytochrome P450 enzymes). **Carbohydrate metabolism** (glycogen breakdown in liver). **Calcium ion (Ca²⁺) storage** (crucial for muscle contraction signaling; in muscle cells, SER is specialized as the **sarcoplasmic reticulum**).

2. Golgi Apparatus / Golgi Complex:

- **Discovery:** Camillo Golgi (1898) using a silver-staining technique.
- **Structure:** A stack of 4-8 flattened, curved cisternae with distinct polarity: **cis face** (forming or receiving face, near ER), **medial cisternae**, and **trans face** (maturing or shipping face). The **trans-Golgi network (TGN)** is a complex of vesicles and tubules for final sorting.
- **Function:** The "post office" and "processing plant" of the cell.
 - **Modification:** Further processes products from ER (e.g., modifies carbohydrate chains on glycoproteins, sulfation).
 - **Synthesis:** Manufactures some polysaccharides (e.g., pectins and hemicellulose for plant cell walls).

- **Sorting, Tagging & Packaging:** Adds molecular "zip codes" (e.g., mannose-6-phosphate for lysosomes) and packages materials into distinct vesicles for delivery to the plasma membrane (secretion), lysosomes, or back to the ER.
- **Vesicle Transport Models: Vesicular Transport Model** (vesicles shuttle between static cisternae) and **Cisternal Maturation Model** (cisternae themselves mature and move from *cis* to *trans* while carrying cargo; supported by modern evidence).

3. Lysosomes:

- **Discovery:** Christian de Duve (1955) via biochemical fractionation; coined the term.
- **Structure:** Spherical, single-membrane-bound vesicles (~0.1-1.2 μm). Maintain an internal acidic pH of ~4.5-5.0 via **V-type H⁺ ATPase pumps** in their membrane.
- **Content:** Contain ~60 different types of **acid hydrolases** (proteases, nucleases, lipases, glycosidases) all optimally active at low pH. The membrane contains highly glycosylated proteins to protect it from self-digestion.
- **Biogenesis:** Enzymes are synthesized in the RER, tagged with mannose-6-phosphate in the *cis*-Golgi, recognized by receptors in the *trans*-Golgi, and packaged into clathrin-coated vesicles that bud to become **primary lysosomes**.
- **Functions:**
 - **Phagocytosis/Heterophagy:** Fuse with phagosomes (containing engulfed bacteria/food) to form **secondary lysosomes** or **phagolysosomes**.
 - **Autophagy:** Engulf and digest damaged or obsolete organelles (**autophagosomes** fuse with lysosomes). A critical recycling and quality control mechanism.
 - **Autolysis:** Rupture of lysosomes leading to total cell destruction (e.g., during metamorphosis, programmed cell death).
- **Clinical Relevance: Lysosomal Storage Diseases** are caused by inherited deficiencies of specific lysosomal enzymes, leading to substrate accumulation. Examples: **Tay-Sachs disease** (hexosaminidase A deficiency, GM2 ganglioside accumulation), **Gaucher's disease** (glucocerebrosidase deficiency).

4. Vacuoles:


- **Structure:** Large, membrane-bound sacs. The membrane is called the **tonoplast** in plant cells.
- **Types:**
 - **Central Vacuole (Plant Cells):** Can occupy up to 90% of cell volume. Contains **cell sap** (water, ions, sugars, pigments like anthocyanins, and toxins). Maintains **turgor pressure** for structural support.
 - **Food Vacuoles:** Form by phagocytosis in protists (e.g., amoeba) and some animal cells.
 - **Contractile Vacuoles:** Found in freshwater protists (e.g., *Paramecium*). Actively pump out excess water for **osmoregulation**.

Energy-Transducing Organelles

1. Mitochondria:


- **Discovery:** Observed in the 19th century; named "bioblasts" by Altmann (1890), term "mitochondrion" coined by Benda (1898).
- **Structure:** Double-membrane bound, dynamic organelles that fuse and divide.
 - **Outer Membrane:** Smooth, contains **porin** proteins, making it permeable to small molecules.
 - **Intermembrane Space:** Between the two membranes.
 - **Inner Membrane:** Highly folded into **cristae**, dramatically increasing surface area. Contains proteins for the **Electron Transport Chain (ETC)** and **ATP synthase (F₀F₁ complex)**. Impermeable to ions; requires specific transporters.

- **Matrix:** Innermost compartment. Contains mitochondrial DNA (mtDNA, circular), 70S ribosomes, granules (Ca^{2+} , phosphate), and enzymes for the **Krebs (TCA) Cycle** and **β -oxidation of fatty acids**.
- **Function:** Site of **aerobic cellular respiration** and **ATP synthesis via oxidative phosphorylation** (Chemiosmotic Theory, Peter Mitchell). Also involved in **apoptosis** (by releasing cytochrome c), **calcium homeostasis**, and **heme synthesis**.



2. Chloroplasts (& other Plastids):

- **Discovery:** Described in plant cells in the 19th century.
- **Structure (Chloroplast):** Double-membrane bound.
 - **Intermembrane Space.**
 - **Stroma:** Fluid-filled interior analogous to mitochondrial matrix. Contains chloroplast DNA (cpDNA), 70S ribosomes, enzymes for the **Calvin Cycle (Carbon Fixation)**, and starch granules.
 - **Thylakoid System:** Internal membrane system. **Thylakoids** are flattened, interconnected sacs; stacks are called **grana** (singular: granum). Thylakoid membranes contain chlorophyll and the protein complexes for the **light-dependent reactions** of photosynthesis.
- **Plastid Diversity:** Plastids are a family of interrelated organelles that can interconvert.
 - **Proplastids:** Undifferentiated precursors in meristems.
 - **Chloroplasts:** Green, photosynthetic.
 - **Chromoplasts:** Contain carotenoid pigments (red, orange, yellow); found in fruits, flowers, roots (carrot).
 - **Leucoplasts:** Colorless; for storage: **Amyloplasts** (starch), **Elaioplasts** (lipids/oils), **Proteinoplasts** (proteins).




Mitochondria and Chloroplasts

Feature	Mitochondrion	Chloroplast
Primary Function	ATP synthesis (Cellular Respiration)	Photosynthesis (ATP & sugar synthesis)
Pigments	Absent	Present (Chlorophyll, Carotenoids)
Energy Source	Oxidation of organic molecules (e.g., glucose)	Sunlight (photons)
Major Processes	Glycolysis (cytosol), Krebs cycle, ETC, Oxidative Phosphorylation	Light Reactions, Calvin Cycle (Carbon Fixation)
Internal Membranes	Cristae (infoldings of inner membrane)	Thylakoids (organized into grana)
Main Product	ATP, CO ₂ , H ₂ O	Glucose, O ₂ , ATP (for plant cell use)
Evolutionary Origin	Alpha-proteobacteria	Cyanobacteria

Endosymbiotic Theory: Strong evidence indicates mitochondria and chloroplasts evolved from engulfed prokaryotes (α -proteobacterium and cyanobacterium, respectively). Evidence: double membranes, own circular DNA, 70S ribosomes, autonomous division.

3. Peroxisomes (Microbodies):

- **Discovery:** Christian de Duve (1960s).
- **Structure:** Small, spherical, single-membrane organelles. **Not part of the endomembrane system;** they can form *de novo* from the ER and divide by fission.
- **Content:** Contain **oxidative enzymes** (e.g., **catalase**, urate oxidase, D-amino acid oxidase). Catalase is a hallmark enzyme, constituting up to 15% of total peroxisomal protein.
- **Functions:**
 - **Detoxification:** Break down toxic **hydrogen peroxide (H₂O₂)** (produced by oxidases) into water and oxygen (**catalase reaction: 2H₂O₂ → 2H₂O + O₂**).
 - **β -Oxidation:** Breakdown of **very long-chain fatty acids (VLCFAs)**, a process distinct from mitochondrial β -oxidation.
 - **Metabolism of purines** and polyamines.
 - **Photorespiration** in plant leaf cells (involves a coordinated cycle between chloroplasts, peroxisomes, and mitochondria).
 - **Glyoxysomes:** Specialized peroxisomes in germinating seeds that house the **glyoxylate cycle**, converting stored fats into carbohydrates.




The Cytoskeleton:

A dynamic, three-dimensional network of protein filaments that provides structural support, enables movement (cell and intracellular), and organizes cellular components.

Cytoskeletal Components


Feature	Microtubules	Microfilaments (Actin Filaments)	Intermediate Filaments
Protein Subunit	α -tubulin & β -tubulin heterodimers	Globular Actin (G-actin) monomers	Fibrous proteins (keratin, vimentin, desmin, lamins, neurofilaments)
Diameter	~25 nm (thickest)	~7 nm (thinnest)	~8-12 nm (intermediate)
Structure	Hollow cylinder of 13 protofilaments. Polar (+ end fast-growing, - end anchored).	Two intertwined helical strands. Polar (+ end barbed, - end pointed).	Rope-like, fibrous proteins coiled together. Non-polar , more stable.
Motor Proteins	Kinesins (move toward + end, usually outward), Dyneins (move toward - end, usually inward).	Myosins (move toward the + end of actin).	None known.
Key Functions	<ul style="list-style-type: none"> - Cell shape (resist compression). - Intracellular transport highways. - Form mitotic/meiotic spindle. - Core of cilia & flagella (axoneme: 9+2 array). - Chromosome movement. 	<ul style="list-style-type: none"> - Cell shape (cortical network, resist tension). - Muscle contraction (with myosin). - Cell motility (amoeboid movement, lamellipodia). - Cytokinesis (formation of contractile ring). - Cytoplasmic streaming (cyclosis) in plant cells. - Core of microvilli. 	<ul style="list-style-type: none"> - Mechanical strength; resist tension (tissue integrity). - Anchor organelles (e.g., nucleus via nuclear lamina). - Permanent structural framework (cell and tissue-specific).

Drug Targets	Colchicine, Vinblastine (depolymerize), Taxol (stabilizes).	Cytochalasin D (depolymerizes), Phalloidin (stabilizes).	No specific drugs.
---------------------	---	--	--------------------



Centrosome & Centrioles:


- **Centrosome:** The primary **Microtubule Organizing Center (MTOC)** in animal cells. Consists of two **centrioles** surrounded by **pericentriolar material (PCM)**, which contains γ -tubulin ring complexes for nucleating microtubule growth.
- **Centrioles:** A pair of cylindrical structures arranged at right angles. Each has a "**9 + 0**" arrangement of **microtubule triplets**. They replicate during the S phase. Function: organize the mitotic spindle poles. **Basal bodies**, which anchor cilia/flagella, are structurally identical to centrioles. Higher plants lack centrioles.



Cilia & Flagella:

- **Structure:** Membrane-bound projections with a core **axoneme**. The axoneme has a "**9 + 2**" **pattern**: nine outer doublet microtubules surrounding a central pair of singlet microtubules. Connected by **nexin links**. Outer doublets have **dynein arms** (motor proteins).

- **Mechanism of Movement: Dynein "walking"** on adjacent microtubule doublets causes sliding. Nexin links convert this sliding into coordinated bending of the entire structure. Requires ATP.
- **Differences: Cilia** are numerous, short (5-10 μm), and beat in coordinated waves (e.g., respiratory epithelium). **Flagella** are typically 1-2 per cell, long (up to 200 μm), and exhibit undulatory or whip-like motion (e.g., sperm).
- **Primary Cilium:** A single, non-motile sensory organelle on most vertebrate cells. Has a "9+0" axoneme. Acts as a **cellular antenna** for chemical and mechanical signals (e.g., Hedgehog signaling pathway).




Plasma Membrane: Structure and Transport

The Fluid Mosaic Model (Singer & Nicolson, 1972)

The membrane is a **fluid bilayer** of phospholipids with a mosaic of embedded proteins.

- **Phospholipids:** Amphipathic molecules forming the bilayer barrier.
- **Cholesterol:** Modulates **membrane fluidity** (restrains movement at high temps, prevents tight packing at low temps). Reduces permeability to small polar molecules.
- **Membrane Proteins:**
 - **Integral/Transmembrane Proteins:** Span the bilayer (e.g., channels, carriers, pumps, receptors). Often have hydrophobic α -helices.
 - **Peripheral Proteins:** Loosely bound to membrane surface, often to integral proteins or lipid heads (e.g., some cytoskeletal elements, signaling proteins).
- **Glycocalyx:** Carbohydrate coat on the extracellular surface from **glycolipids** and **glycoproteins**. Functions in cell recognition (e.g., ABO blood groups), adhesion, and protection.



Membrane Transport Mechanisms


Mechanism	Requires Protein?	Requires ATP?	Direction	Example
Simple Diffusion	No	No	High → Low Concentration	O ₂ , CO ₂ , lipids, ethanol
Facilitated Diffusion	Yes (Channel or Carrier)	No	High → Low Concentration	Glucose (via GLUT carriers), ions (K ⁺ channels), water (aquaporins)
Osmosis	Yes (Aquaporins) or No	No	Low Solute → High Solute (Water moves)	Water across any cell membrane
Primary Active Transport	Yes (Pump)	Yes	Against Concentration Gradient	Na⁺/K⁺ ATPase Pump, H⁺ Pump (V-ATPase), Ca²⁺ Pump
Secondary Active Transport	Yes (Symporter/Antiporter)	Indirect (uses ion gradient)	Against Gradient (coupled with <i>downhill</i> movement)	Na⁺/Glucose Symporter (intestine), Na⁺/Ca²⁺ Antiporter

Bulk Transport (Requires ATP and membrane trafficking machinery):

- **Endocytosis:** Import. **Phagocytosis** ("cell eating" of solids), **Pinocytosis** ("cell drinking" of fluid), **Receptor-Mediated Endocytosis** (highly specific, e.g., LDL-cholesterol uptake via clathrin-coated pits).
- **Exocytosis:** Export. Vesicles from the Golgi fuse with the plasma membrane to release contents (e.g., neurotransmitters, hormones, digestive enzymes).

Plant Cell Walls

- **Primary Cell Wall:** Flexible, thin. Contains **cellulose microfibrils** embedded in a matrix of **hemicellulose**, **pectin**, and structural glycoproteins.
- **Middle Lamella:** Outermost layer, rich in **pectin**, glues adjacent plant cells together.
- **Secondary Cell Wall:** Deposited inside the primary wall in some cells (e.g., xylem vessels, fibers). Contains more cellulose and is impregnated with **lignin** for extreme strength and waterproofing.
- **Plasmodesmata:** Cytoplasmic channels lined with plasma membrane that traverse cell walls, connecting the cytosol of adjacent cells. Contain a central **desmotubule** (derived from ER). Allow transport of water, small solutes, proteins, and even RNA (**symplastic transport**).



Extracellular Matrix (ECM) of Animal Cells

A complex meshwork secreted by the cells.

- **Components:** **Collagen** (provides tensile strength), **Elastin** (elasticity), **Proteoglycans** (protein cores with long GAG chains; form a hydrated gel resisting compression), **Glycoproteins** (e.g., **Fibronectin**, Laminin - facilitate adhesion).
- **Integrins:** Transmembrane receptor proteins. Their extracellular domains bind to ECM components (e.g., fibronectin); their intracellular domains link to the **cytoskeleton** (via talin, vinculin) and signaling molecules. Mediate **mechanotransduction** (converting mechanical force into biochemical signals) and **outside-in/inside-out signaling**.

Major Animal Cell Junctions

Junction Type	Main Proteins	Cytoskeletal Link	Primary Function
Tight Junctions	Claudins, Occludins	Actin filaments	Seal adjacent epithelial cells into a leak-proof sheet. Create apical/basolateral polarity.
Adherens Junctions	Cadherins (Classical)	Actin filaments (via α/β -catenins)	Strong adhesion, mechanical coupling, and signaling between cells.
Desmosomes	Desmoglein, Desmocollin (Cadherin family)	Intermediate filaments (Keratin)	Provide strong, spot-welded adhesion, especially in tissues under mechanical stress (skin, heart).
Hemidesmosomes	Integrins ($\alpha6\beta4$)	Intermediate filaments (Keratin)	Anchor epithelial cells to the basal lamina (a specialized ECM), not to another cell.
Gap Junctions	Connexins (form hexameric connexons)	None	Provide direct cytoplasmic continuity for passage of ions, second messengers (cAMP, Ca^{2+} , IP_3), and small metabolites (< ~1 kDa). Crucial for electrical coupling in cardiac/smooth muscle and rapid communication.

Stem Cells

Undifferentiated cells with **self-renewal** and **differentiation** potential.

- **Potency Classifications:**
 - **Totipotent:** Can form all cell types + extraembryonic tissues (zygote).
 - **Pluripotent:** Can form all embryonic cell types (**Embryonic Stem Cells - ESCs**).
 - **Multipotent:** Can form multiple, related lineages (e.g., **Hematopoietic Stem Cells**).
 - **Unipotent:** Produce one cell type (e.g., epidermal stem cells).
- **Induced Pluripotent Stem Cells (iPSCs):** Adult somatic cells reprogrammed to an embryonic-like state. Advantages: patient-specific, avoids ethical issues of ESCs. Disadvantages: risk of tumorigenesis, low efficiency.

Eukaryotic Cell Components

Structure	Main Function	Key Features (Animal Cell Focus)
Nucleus	Genetic control center, RNA synthesis	Double membrane, pores, chromatin, nucleolus
Ribosomes	Protein synthesis (translation)	rRNA & protein, free or bound to RER
Rough ER	Protein synthesis & processing, membrane factory	Ribosome-studded, continuous with nuclear envelope
Smooth ER	Lipid synthesis, detoxification, Ca^{2+} storage	No ribosomes, tubular structure

Golgi Apparatus	Modification, sorting, packaging of macromolecules	Cis/trans polarity, vesicle budding
Lysosomes	Intracellular digestion, autophagy, recycling	Acidic pH, hydrolytic enzymes
Vacuoles	Storage, digestion, water balance (plants)	Large central vacuole in plants
Mitochondria	ATP production (cellular respiration)	Double membrane, cristae, own DNA, dynamic
Peroxisomes	Fatty acid oxidation, detoxification (H ₂ O ₂ metabolism)	Single membrane, oxidative enzymes
Cytoskeleton	Structure, motility, transport, organization	Microtubules, microfilaments, intermediate filaments
Centrioles	Microtubule organization (cell division)	9x3 microtubule array, part of centrosome
Cilia/Flagella	Motility, fluid movement, sensory reception	9+2 microtubule array, dynein motor
Plasma Membrane	Selective barrier, cell signaling, adhesion	Phospholipid bilayer with proteins, glycocalyx
ECM	Support, adhesion, movement, signaling	Collagen, proteoglycans, fibronectin-integrin links
Cell Junctions	Intercellular connection & communication	Tight junctions, desmosomes, gap junctions

Comparison of Animal and Plant Cells

Animal and plant cells are both eukaryotic but exhibit key structural and functional differences.

Animal Cell

- Lacks a cell wall; enclosed only by a **plasma membrane**.
- Plastids (e.g., chloroplasts) are absent.
- Contains **centrosomes** with centrioles.
- Lysosomes are present.
- Multiple, small, temporary vacuoles.
- Stores carbohydrates as **glycogen**.
- Typically smaller and irregular in shape.
- Nucleus is usually central.
- Cytokinesis occurs by inward furrowing.
- Can perform phagocytosis.

Plant Cell

- Possesses a rigid **cell wall** (primarily cellulose) outside the plasma membrane.
- Contains plastids like **chloroplasts** (for photosynthesis) and chromoplasts.
- Lacks centrioles; uses **polar caps** for spindle formation.
- Lysosomes are generally absent (functions carried out by vacuole and other organelles).
- Has a large, central **vacuole** for storage, turgor pressure, and waste management.
- Stores carbohydrates as **starch**.
- Typically larger and has a fixed, rectangular shape.
- Nucleus is often pushed to the periphery by the central vacuole.
- Cytokinesis occurs by the formation of a **cell plate** growing outward.
- Does not perform phagocytosis.

Animal vs. Plant Cell

Feature	Animal Cell	Plant Cell
Cell Wall	Absent	Present (Cellulose)
Plastids	Absent	Present (Chloroplasts, etc.)



MK PREPARATIONS



Centrioles	Present	Absent
Vacuoles	Small, numerous	Single, large central vacuole
Lysosomes	Present	Usually absent
Shape	Irregular, round	Fixed, rectangular
Storage Product	Glycogen	Starch
Mode of Nutrition	Heterotrophic	Autotrophic (primarily)

2. Cell Biology

M
K

P
R
E
P
A
R
A
T
I
O
N
S



Cell Biology: One Liner

- Cells are the **basic unit of structure and function** of all living organisms.
- All organisms are made of **cells**, which exist as **single-celled** or **multicellular** entities.
- A cell is the **smallest unit** and **building block of life**.
- Cells exist in various **shapes and sizes** related to their specific functions.
- The **seven properties of life** are movement, respiration, homeostasis, growth, reproduction, excretion, and nutrition.
- **Movement** refers to external or internal motion by living things.
- **Respiration** is the use of environmental substances by living things to make energy.
- **Homeostasis** is the maintenance of a stable internal environment by living things.
- **Growth** is the ability of living things to change shape or size.
- **Reproduction** is the ability of living things to produce offspring sexually or asexually.
- **Excretion** is the removal of waste products by living things.
- **Nutrition** involves the exchange of materials and gases with the environment by living things.
- **Robert Hooke** (1665) first observed and named "**cells**" while examining cork under a microscope.
- **Matthias Schleiden** (1838), a German botanist, concluded all **plant tissues** are composed of cells.
- **Theodor Schwann** (1839), a German zoologist, concluded all **animal tissues** are also composed of cells.
- **Rudolf Virchow** (1855) stated that new cells arise only from **pre-existing cells**.
- The **cell theory** is a fundamental unifying concept in biology.
- The first tenet of cell theory is that the **cell is the fundamental unit of structure and function** in living things.
- The second tenet of cell theory is that **all organisms are composed of one or more cells**.
- The third tenet of cell theory is that **cells arise from pre-existing cells through cellular division**.
- **Validation of cell theory** includes evidence that cells removed from tissues can survive independently for short periods.
- **Validation of cell theory** includes the fact that nothing smaller than a cell can live independently.
- **Validation of cell theory** includes experiments by **Francesco Redi** and **Louis Pasteur** showing cells cannot arise in sterile conditions.
- **Viruses** are a major exception to cell theory as they are **acellular** and lack cell machinery.
- **Viroid's and prions** are also exceptions to the cell theory, behaving like viruses.
- **Mature Red Blood Cells (RBCs)** and **sieve tube cells** lack nuclei and cannot divide, forming an exception.
- **Bacteria and cyanobacteria** lack a well-organized nucleus, with DNA lying in direct contact with cytoplasm, forming an exception.
- **Coenocytic hyphae** of fungi and cells of some algae are **multinucleate**, forming an exception.
- **Protozoans** are **acellular** as their body is not divided into cells, forming an exception.
- **Microscopy** is the technique used to view objects that cannot be seen by the naked eye.
- A **light microscope (LM)** uses visible light to illuminate specimens in a two-lens (compound) system.
- The two lenses in a light microscope are the **ocular lens (eyepiece)** and the **objective lens**.
- The **working principle** of a light microscope involves illumination, image formation by the objective lens, and final magnification by the eyepiece.
- **Magnification** is the capacity of an optical instrument to increase the apparent size of an object.
- Magnification power is represented by the symbol "**X**" (times).
- **Total magnification** is calculated by multiplying the objective lens value by the ocular lens value.

- The **maximum magnification** of a standard light microscope is approximately **1500X**.
- **Resolution (resolving power)** is a measure of a microscope's ability to distinguish between two close points.
- The **resolution of the naked eye** is about **0.1 mm**.
- The **resolving power of a light microscope** is about **250 nm**.
- **Magnification** makes small objects appear larger, while **resolution** is the ability to distinguish between two objects.
- An **electron microscope** uses a beam of accelerated electrons as an illuminating source.
- The **working principle** of an electron microscope involves focusing an electron beam on a thin specimen and forming an image based on electron scattering.
- The **resolving power of an electron microscope** is about **0.2 nm**.
- An **electron microscope** can achieve magnifications between **1 and 50 million times**.
- A **light microscope** can view **both live and dead specimens**.
- An **electron microscope** can view **only dead and dried specimens**.
- In a light microscope, image formation depends on **light absorption**.
- In an electron microscope, image formation depends on **electron scattering**.
- An **electron microscope** requires a **vacuum** for operation, while a light microscope does not.
- Images from a light microscope are usually **colored**, while electron microscope images are **black and white**.
- The **ultrastructure** of a cell refers to its fine structure as seen with an electron microscope.
- The **Svedberg unit (S)** measures the sedimentation rate of a particle during centrifugation.
- The **Svedberg unit** indicates the **size** of a molecule, with larger molecules having larger S values.
- The two major types of electron microscopes are the **Transmission Electron Microscope (TEM)** and the **Scanning Electron Microscope (SEM)**.
- **TEM** produces an image by detecting **transmitted electrons** through a specimen.
- **SEM** produces an image by detecting **secondary or backscattered electrons** from a specimen's surface.
- The **plasma (cell) membrane** is the outer living boundary of the cell.
- A **cell wall** is an extracellular component formed exterior to the plasma membrane in some cells.
- **Cell walls** are present in plant cells, prokaryotes, fungi, and some protists, but **not in animal cells**.
- The plant cell wall is an **extracellular structure** that protects the cell, maintains shape, and prevents excessive water uptake.
- The plant cell wall is **permeable**, allowing free passage of water and dissolved materials.
- The plant cell wall consists of three main layers: **primary cell wall, middle lamella, and secondary cell wall**.
- The **primary cell wall** is thin, flexible, and develops in newly growing cells.
- The primary cell wall is composed of **cellulose microfibrils** in a matrix of **hemicellulose and pectin**.
- Cellulose microfibrils show a **crisscross arrangement**, providing great strength.
- The **secondary cell wall** is thick, rigid, and formed inside the primary wall in some cells (e.g., sclerenchyma).
- The secondary cell wall contains **cellulose, hemicellulose, lignin, inorganic salts, and waxes**.
- **Lignin** in the secondary cell wall provides rigidity and mechanical support.
- The **middle lamella** is a sticky, gel-like layer composed of **pectin and magnesium/calcium salts** that holds adjacent plant cells together.
- The **plasma membrane** is found in all living prokaryotic and eukaryotic cells.
- Chemically, the plasma membrane consists of **proteins (60-80%), lipids (20-40%), and a small quantity of carbohydrates**.

- The **fluid mosaic model** describes the plasma membrane as a **phospholipid bilayer** with embedded or attached proteins.
- The phospholipid bilayer is arranged with **hydrophobic tails facing inward** and **hydrophilic heads facing outward**.
- **Cholesterol** molecules are wedged into the phospholipid bilayer, affecting membrane fluidity.
- The plasma membrane is **asymmetrical**, meaning its two surfaces are not identical.
- **Membrane proteins** can drift sideways within the fluid bilayer.
- Carbohydrates on the membrane are attached to proteins (**glycoproteins**) or lipids (**glycolipids**), mainly on the outer surface.
- **Cytoskeleton filaments** are present on the inner surface of the plasma membrane for support.
- Plasma membrane proteins function as **transport channels or carriers, enzymes, receptors, or antigens**.
- **Glycolipids and glycoproteins** often act as **cell surface markers** for cell-to-cell recognition.
- The plasma membrane regulates the cell's interaction with its environment by **controlling material transport**.
- Techniques to study cell membranes include **atomic force microscopy, super-resolution fluorescent microscopy, and Scanning Electron Microscopy (SEM)**.
- **Cell signaling** is the process by which cells communicate with other cells or with the external environment.
- Signaling molecules are called **ligands**, which can be proteins, lipids, gases, or other molecules.
- Proteins that respond to ligands are called **receptors**.
- The first step in cell signaling is **reception**, where a signal molecule binds to a specific receptor.
- The second step in cell signaling is **transduction**, converting the extracellular signal into an intracellular form.
- The final step in cell signaling is a specific **cellular response**, like enzyme activation or gene expression.
- A **signal transduction pathway** is the series of steps converting a cell surface signal into a cellular response.
- **Protein/Peptide hormones** are water-soluble and act as **first messengers**, binding to receptors on the target cell's plasma membrane.
- Protein signaling often involves the production of a **second messenger** (e.g., cAMP) inside the cell.
- **Steroid hormones** are small, hydrophobic, and lipophilic, allowing them to diffuse directly through the plasma membrane.
- Steroid hormones bind to **internal receptors** in the cytoplasm or nucleus, forming a **hormone-receptor complex**.
- The activated steroid hormone-receptor complex acts as a **transcription factor**, binding to DNA to regulate gene expression.
- **Membrane transport** controls the movement of solutes like ions and small molecules through biological membranes.
- The four primary transport mechanisms are **simple diffusion, facilitated diffusion, osmosis, and active transport**.
- **Simple diffusion** is the passive movement of molecules from an area of **higher concentration to lower concentration** down a concentration gradient.
- Small, nonpolar molecules like **oxygen, carbon dioxide, and water** can cross the membrane via simple diffusion.
- **Facilitated diffusion** is the passive movement of molecules **down their concentration gradient** with the assistance of transport proteins.
- The two types of transport proteins in facilitated diffusion are **channel proteins and carrier proteins**.

- **Channel proteins** form hydrophilic pores for charged molecules or ions to pass through.
- **Gated channel proteins** can open or close to regulate substance passage.
- **Carrier proteins** bind to specific molecules, change shape, and release them on the other side of the membrane.
- **Osmosis** is the diffusion of **water** across a **semipermeable membrane** from a region of low solute concentration to high solute concentration.
- A **semipermeable membrane** allows solvent (water) molecules to pass but blocks larger solute molecules.
- **Active transport** moves molecules or ions **against their concentration gradient** from low to high concentration, requiring energy (ATP).
- **Primary active transport** directly uses **ATP** as an energy source (e.g., Sodium-Potassium pump).
- The **Sodium-Potassium pump (Na⁺/K⁺ ATPase)** actively transports **3 Na⁺ ions out** and **2 K⁺ ions into** the cell.
- **Secondary active transport** uses the energy from an **ion's electrochemical gradient** (often Na⁺ or H⁺) to transport another molecule against its gradient.
- **Endocytosis** is the bulk transport of materials **into the cell** via vesicle formation from the plasma membrane.
- **Phagocytosis** ("cell eating") is a type of endocytosis where large particles or cells are engulfed.
- **Pinocytosis** ("cell drinking") is a type of endocytosis where extracellular fluid and dissolved solutes are taken in.
- **Receptor-mediated endocytosis** is a specific form of pinocytosis where ligands bind to receptors in **coated pits** (e.g., cholesterol uptake).
- **Exocytosis** is the bulk transport of materials **out of the cell** via vesicle fusion with the plasma membrane.
- The living matter of a cell is called **protoplasm**.
- In eukaryotic cells, protoplasm is divided into **cytoplasm** and **nucleus**.
- **Cytoplasm** is the region between the nuclear membrane and the plasma membrane.
- **Cytosol** is the water-soluble, liquid matrix of the cytoplasm surrounding organelles.
- The cytoplasm acts as a **site of metabolism** and a **storehouse** for the cell.
- **Cell organelles** are discrete, membrane-bound (mostly) structures within eukaryotic cells with specific functions.
- The **Endoplasmic Reticulum (ER)** is an extensive network of membranous tubules and sacs (cisternae).
- The ER lumen (cisternal space) is the internal compartment separated from the cytosol.
- **Rough ER (RER)** has ribosomes attached to its cytoplasmic surface, giving it a "rough" appearance.
- **Rough ER** functions in the synthesis of **secretory proteins, membrane proteins, and organelle proteins**.
- Newly synthesized proteins in the RER can be modified by **glycosylation**.
- **Smooth ER (SER)** lacks ribosomes and appears smooth.
- **Smooth ER** functions in **lipid synthesis** (steroids, phospholipids), **detoxification** of drugs, and **calcium ion (Ca²⁺) storage**.
- In muscle cells, the specialized SER is called the **sarcoplasmic reticulum**.
- **Ribosomes** are non-membranous, granular organelles composed of **rRNA and protein** (ribonucleoprotein).
- Ribosomes are the site of **protein synthesis (translation)**.
- **Eukaryotic ribosomes** are **80S**, composed of a large 60S subunit and a small 40S subunit.
- **Prokaryotic ribosomes** are **70S**, composed of a large 50S and a small 30S subunit.

- **Polysomes (polyribosomes)** are chains of multiple ribosomes translating a single mRNA molecule simultaneously.
- The **Golgi complex (Golgi apparatus)** consists of a stack of flattened membrane sacs called **cisternae**.
- The Golgi has a **cis face** (receiving side near ER) and a **trans face** (shipping side).
- The Golgi complex functions in the **modification, sorting, packaging, and distribution** of cell products (e.g., glycoproteins).
- The Golgi is involved in the synthesis of some **polysaccharides** (e.g., for plant cell walls).
- **Vesicles** are small, membrane-bound sacs that transport materials within the cell.
- **Lysosomes** are single-membrane-bound vesicles containing **hydrolytic (digestive) enzymes** active at acidic pH.
- Lysosomal enzymes are synthesized in the **RER** and processed in the **Golgi apparatus**.
- **Primary lysosomes** are newly formed lysosomes before they fuse with other vesicles.
- Lysosomes fuse with food vacuoles (**phagocytosis**) or damaged organelles (**autophagy**) to form **secondary lysosomes** for digestion.
- Lysosomes are involved in **autolysis** (cell self-digestion).
- **Peroxisomes** are single-membrane-bound vesicles containing oxidative enzymes like **catalase and peroxidase**.
- Peroxisomes are involved in **detoxification**, breaking down hydrogen peroxide (**H₂O₂**) into water and oxygen.
- In plant cells, peroxisomes are involved in **photorespiration**.
- **Glyoxysomes** are specialized peroxisomes in oil seed plants that contain enzymes for the **glyoxylate cycle**, converting fats to carbohydrates.
- **Vacuoles** are large vesicles originating from the ER, Golgi, and plasma membrane.
- In animal cells, **food vacuoles** form by phagocytosis.
- **Contractile vacuoles** in freshwater protists pump excess water out for **osmoregulation**.
- Plant cells have a large **central vacuole** containing **cell sap** (water, ions, pigments).
- The central vacuole maintains **turgor pressure** for mechanical support in plant cells.
- The vacuole membrane is called the **tonoplast** in plant cells.
- **Mitochondria** are double-membrane-bound organelles found in all eukaryotic cells.
- The mitochondrial **outer membrane** is smooth and contains **porins**, making it permeable to small molecules.
- The mitochondrial **inner membrane** is highly folded into **cristae** to increase surface area.
- The inner membrane contains proteins for the **Electron Transport Chain (ETC)** and **ATP synthase (FO-F1 particles)**.
- The space between the two mitochondrial membranes is the **intermembrane space**.
- The innermost compartment is the **mitochondrial matrix**, containing enzymes, mitochondrial DNA (mtDNA), and 70S ribosomes.
- Mitochondria are the sites of **aerobic cellular respiration** and **ATP production** via oxidative phosphorylation.
- Mitochondria are **semi-autonomous** organelles capable of self-replication by fission.
- **Plastids** are double-membrane-bound organelles in plant and algal cells, classified by pigment presence.
- **Proplastids** are young, undifferentiated plastids.
- **Leucoplasts** are colorless plastids for storage (e.g., amyloplasts store starch).
- **Chromoplasts** contain pigments other than green (e.g., carotenoids).
- **Chloroplasts** are green plastids that perform **photosynthesis**.
- The chloroplast is bounded by a double membrane envelope.
- The internal fluid-filled space of the chloroplast is the **stroma**, containing enzymes, chloroplast DNA (cpDNA), and 70S ribosomes.

- The **thylakoid** system inside chloroplasts consists of flattened membranous sacs.
- Stacks of thylakoids are called **grana** (singular: granum).
- Thylakoid membranes contain **chlorophyll** and the machinery for the **light-dependent reactions** of photosynthesis.
- The **stroma** is the site of the **light-independent reactions (Calvin cycle)** of photosynthesis.
- **Centrioles** are non-membranous, rod-shaped organelles found in pairs near the nucleus in animal cells.
- Each centriole is composed of **nine triplets of microtubules** arranged in a circle.
- Centrioles help organize the **microtubules of the mitotic spindle** during cell division.
- Centrioles also give rise to the **basal bodies** of cilia and flagella.
- The **cytoskeleton** is a network of protein filaments providing structural support, enabling movement, and organizing the cytoplasm.
- The three types of cytoskeletal fibers are **microfilaments, microtubules, and intermediate filaments**.
- **Microfilaments (Actin filaments)** are the thinnest filaments (~7 nm), composed of **actin** protein.
- Microfilaments are involved in **cell movement, muscle contraction, cytokinesis, and cytoplasmic streaming**.
- **Microtubules** are the thickest filaments (~25 nm), hollow tubes composed of **α and β -tubulin** dimers.
- Microtubules function in **cell shape maintenance, intracellular transport, and forming the mitotic spindle and cilia/flagella cores**.
- **Intermediate filaments** (8-12 nm) are composed of various fibrous proteins (e.g., keratin, vimentin, lamins).
- Intermediate filaments provide **mechanical strength and structural stability** to cells and tissues.
- The **nucleus** is the most prominent, membrane-bound organelle housing the cell's genetic material.
- The nucleus is bounded by a double membrane called the **nuclear envelope**.
- The **nuclear envelope** is perforated by **nuclear pores** that regulate molecular traffic between the nucleus and cytoplasm.
- The **nuclear lamina** is a dense network of intermediate filaments (lamins) lining the inner nuclear membrane, providing structural support.
- The fluid inside the nucleus is the **nucleoplasm**.
- The **nucleolus** is a dense, non-membrane-bound region within the nucleus where **rRNA is transcribed and ribosomal subunits are assembled**.
- **Chromatin** is the complex of DNA and proteins (histones) found within the nucleus.
- During cell division, chromatin condenses to form visible **chromosomes**.
- A duplicated chromosome consists of two identical **sister chromatids** joined at the **centromere**.
- **Prokaryotic cells** (Bacteria and Archaea) lack a membrane-bound nucleus and other membrane-bound organelles.
- **Eukaryotic cells** (Protists, Fungi, Plants, Animals) possess a membrane-bound nucleus and various membrane-bound organelles.
- In prokaryotic cells, DNA is located in a region called the **nucleoid**.
- Prokaryotic ribosomes are **70S**, while eukaryotic cytoplasmic ribosomes are **80S**.
- Prokaryotes divide by **binary fission**, while eukaryotes divide by **mitosis** (somatic cells) or **meiosis** (germ cells).
- **Stem cells** are undifferentiated cells with the potential to develop into many different specialized cell types.
- The key properties of stem cells are **self-renewal** and **differentiation**.
- **Adult (somatic) stem cells** are multipotent, giving rise to a limited range of cell types within a particular tissue.

- **Embryonic Stem Cells (ESCs)** are pluripotent, capable of forming all cell types of the body.
- **Induced Pluripotent Stem Cells (iPSCs)** are adult cells reprogrammed in the lab to an embryonic-like pluripotent state.
- **Totipotent** stem cells (e.g., zygote) can form all cell types plus extraembryonic tissues.
- **Pluripotent** stem cells can form all embryonic cell types.
- **Multipotent** stem cells can form multiple, closely related cell types.
- **Oligopotent** stem cells can differentiate into only a few cell types.
- **Unipotent** stem cells can produce only one cell type but can self-renew.
- **Advantages of using stem cells** include their potential in regenerative medicine for treating diseases like spinal cord injuries, diabetes, and Parkinson's.
- **Advantages of iPSCs** include avoiding ethical issues with ESCs and potential for patient-specific therapies.
- **Disadvantages of iPSCs** include the risk of tumorigenesis (cancer) due to the methods used for reprogramming (e.g., retroviruses).
- **Cell division** is essential for the continuity of life, based on the reproduction of cells.
- The two main types of cell division are **mitosis** (produces identical somatic cells) and **meiosis** (produces gametes/spores with half the chromosomes).
- The **cell cycle** is the series of stages a cell passes through to grow and divide.
- The two main phases of the cell cycle are **interphase** (growth and DNA replication) and the **mitotic (M) phase** (division).
- Interphase consists of **G1 phase** (first gap/growth), **S phase** (DNA synthesis), and **G2 phase** (second gap/preparation for division).
- DNA replication occurs only during the **S phase** of interphase.
- Cell cycle checkpoints (G1, G2, M) are control mechanisms that ensure proper progression and integrity of the cell cycle.
- **Mitosis** is the division of the nucleus resulting in two genetically identical daughter nuclei.
- For description, mitosis is divided into five phases: **prophase, prometaphase, metaphase, anaphase, and telophase**.
- In **prophase**, chromosomes condense, the nucleolus disappears, and the mitotic spindle begins to form.
- In **prometaphase**, the nuclear envelope breaks down, and spindle microtubules attach to chromosome kinetochores.
- In **metaphase**, chromosomes align at the cell's equator (**metaphase plate**).
- In **anaphase**, sister chromatids separate and move to opposite poles as individual chromosomes.
- In **telophase**, chromosomes de-condense, and new nuclear envelopes form around the two daughter nuclei.
- **Cytokinesis** is the division of the cytoplasm, usually following telophase.
- In animal cells, cytokinesis occurs via the formation of a **cleavage furrow**.
- In plant cells, cytokinesis occurs via the formation of a **cell plate**.
- **Significance of mitosis** includes growth, tissue repair, asexual reproduction, and maintaining genetic consistency.
- **Meiosis** is a specialized cell division that reduces the chromosome number by half, producing four haploid cells.
- Meiosis consists of **two consecutive divisions: Meiosis I** (reductional) and **Meiosis II** (equational).
- **Prophase I** of meiosis is lengthy and involves **synapsis** (pairing of homologous chromosomes) and **crossing over** (genetic recombination).
- Paired homologous chromosomes during prophase I form a **tetrad (bivalent)** consisting of four chromatids.
- The site of crossing over is visible as a **chiasma** (plural: chiasmata).



MK PREPARATIONS



- In **Metaphase I**, tetrads align at the metaphase plate, with homologous chromosomes facing opposite poles.
- In **Anaphase I**, homologous chromosomes separate and move to opposite poles (sister chromatids remain attached).
- **Telophase I and cytokinesis** result in two haploid cells, each with duplicated chromosomes.
- **Meiosis II** resembles a mitotic division, separating sister chromatids.
- **Significance of meiosis** includes producing genetic variation through crossing over and independent assortment, and maintaining a constant chromosome number across generations.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

2. Cell Biology



Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. Which scientist first coined the term "cell" after observing cork under a microscope?

- A) Anton van Leeuwenhoek
- B) Robert Brown
- C) Matthias Schleiden
- D) Robert Hooke

Answer: Robert Hooke

2. The principle "Omnis cellula e cellula" (all cells come from cells) was proposed by:

- A) Rudolf Virchow
- B) Louis Pasteur
- C) Theodor Schwann
- D) August Weismann

Answer: Rudolf Virchow

3. Which of the following is NOT a tenet of modern cell theory?

- A) All living organisms are composed of one or more cells.
- B) All cells arise from pre-existing cells.
- C) All cells contain a membrane-bound nucleus.
- D) Cells contain hereditary material (DNA) passed to daughter cells.

Answer: All cells contain a membrane-bound nucleus.

4. The maximum theoretical resolution of a standard light microscope is approximately:

- A) 0.2 nm
- B) 200 nm
- C) 2 μ m
- D) 0.2 mm

Answer: 200 nm

5. Which microscopy technique enhances contrast in unstained, living cells by converting differences in refractive index into brightness variations?

- A) Bright-field microscopy
- B) Dark-field microscopy
- C) Phase-contrast microscopy
- D) Fluorescence microscopy

Answer: Phase-contrast microscopy

6. Transmission Electron Microscopy (TEM) is primarily used to study:

- A) Surface topography of specimens
- B) Internal ultrastructure of thin sections
- C) Living cellular processes
- D) Fluorescently tagged proteins

Answer: Internal ultrastructure of thin sections

7. Cryo-electron microscopy (cryo-EM) is notable for:

- A) Using heavy metal stains to enhance contrast
- B) Visualizing specimens in a near-native, hydrated state
- C) Being suitable only for live cell imaging
- D) Having a resolution limit of about 200 nm

Answer: Visualizing specimens in a near-native, hydrated state

8. Cell fractionation and differential centrifugation separate cellular components based on:

- A) Color and shape
- B) Electrical charge
- C) Size and density
- D) Magnetic properties

Answer: Size and density

9. All cells, prokaryotic and eukaryotic, share which of the following features?

- A) Membrane-bound organelles
- B) A cell wall made of peptidoglycan
- C) A phospholipid bilayer plasma membrane
- D) A well-defined nucleus

Answer: A phospholipid bilayer plasma membrane

10. In prokaryotic cells, the region where DNA is located is called the:

- A) Nucleus
- B) Nucleolus
- C) Nucleoid
- D) Nuclear envelope

Answer: Nucleoid

11. Which of the following is a key structural difference between bacterial and archaeal cell membranes?

- A) Bacteria have ester-linked lipids; Archaea have ether-linked lipids.
- B) Archaea have peptidoglycan in their cell walls; Bacteria do not.
- C) Bacteria have a nucleus; Archaea do not.
- D) Archaea have 80S ribosomes; Bacteria have 70S ribosomes.

Answer: Bacteria have ester-linked lipids; Archaea have ether-linked lipids.

12. Eukaryotic ribosomes in the cytosol are designated as:

- A) 70S
- B) 80S

C) 60S

D) 50S

Answer: 80S

13. Which organelle is responsible for the synthesis of ribosomal RNA (rRNA) and assembly of ribosomal subunits?

A) Nucleus

B) Nucleolus

C) Rough ER

D) Golgi apparatus

Answer: Nucleolus

14. Nuclear Pore Complexes (NPCs) function primarily to:

A) Synthesize DNA

B) Regulate active transport of macromolecules between nucleus and cytoplasm

C) Provide structural support to the nuclear envelope

D) Digest damaged organelles

Answer: Regulate active transport of macromolecules between nucleus and cytoplasm

15. Ribosomes that are attached to the cytosolic face of the Rough ER synthesize proteins destined for:

A) Use within the cytosol

B) Secretion, membranes, or specific organelles

C) The mitochondrial matrix

D) The nucleoplasm

Answer: Secretion, membranes, or specific organelles

16. The function of the Smooth Endoplasmic Reticulum (SER) includes all EXCEPT:

A) Lipid and steroid hormone synthesis

B) Detoxification of drugs and poisons

C) Calcium ion storage

D) Synthesis of secretory proteins

Answer: Synthesis of secretory proteins

17. In muscle cells, the specialized form of smooth ER that stores calcium ions is called the:

A) Golgi apparatus

B) Sarcoplasmic reticulum

C) Peroxisome

D) Lysosome

Answer: Sarcoplasmic reticulum

18. The "cis" face of the Golgi apparatus is typically oriented:

A) Toward the plasma membrane

B) Away from the ER

C) Near the ER, receiving transport vesicles

D) Toward the nucleus

Answer: Near the ER, receiving transport vesicles

19. Lysosomes maintain an internal acidic pH of about 4.5-5.0 using:

A) Sodium-potassium pumps

B) V-type H⁺ ATPase pumps

C) Proton channels

D) Calcium pumps

Answer: V-type H⁺ ATPase pumps

20. Autophagy refers to the process where lysosomes:

A) Digest materials taken in by phagocytosis

B) Fuse with the plasma membrane for exocytosis

C) Digest the cell's own damaged organelles

D) Break down very long-chain fatty acids

Answer: Digest the cell's own damaged organelles

21. Tay-Sachs disease is an example of a lysosomal storage disorder caused by a deficiency in:

A) Glucocerebrosidase

B) Hexosaminidase A

C) Catalase

D) Cytochrome P450

Answer: Hexosaminidase A

22. The large central vacuole in plant cells is primarily involved in all EXCEPT:

A) Storage of cell sap

B) Maintenance of turgor pressure

C) Photosynthesis

D) Waste management

Answer: Photosynthesis

23. The infoldings of the inner mitochondrial membrane are called:

A) Cristae

B) Thylakoids

C) Cisternae

D) Grana

Answer: Cristae

24. According to the Endosymbiotic Theory, mitochondria are thought to have evolved from:

A) Cyanobacteria

B) Alpha-proteobacteria

C) Archaea

D) Eukaryotic nuclei

Answer: Alpha-proteobacteria

25. Which organelle contains catalase and is involved in the breakdown of hydrogen peroxide (H₂O₂)?

- A) Peroxisome
- B) Lysosome
- C) Glyoxysome
- D) Golgi apparatus

Answer: Peroxisome

26. Glyoxysomes, specialized peroxisomes in germinating seeds, are involved in:

- A) The glyoxylate cycle to convert fats to carbohydrates
- B) The Calvin cycle for carbon fixation
- C) The Krebs cycle for ATP production
- D) Photorespiration

Answer: The glyoxylate cycle to convert fats to carbohydrates

27. The thickest cytoskeletal filaments, composed of tubulin dimers, are:

- A) Microfilaments
- B) Intermediate filaments
- C) Microtubules
- D) Keratin filaments

Answer: Microtubules

28. Which motor protein is associated with movement toward the minus end of microtubules?

- A) Kinesin
- B) Myosin
- C) Dynein
- D) Actin

Answer: Dynein

29. The "9+2" arrangement of microtubule doublets is characteristic of the core structure of:

- A) Centrioles
- B) The nuclear lamina
- C) Cilia and flagella (axoneme)
- D) Microvilli

Answer: Cilia and flagella (axoneme)

30. The primary function of the centrosome in animal cells is to:

- A) Synthesize ATP
- B) Act as the Microtubule Organizing Center (MTOC)
- C) Digest macromolecules
- D) Store calcium ions

Answer: Act as the Microtubule Organizing Center (MTOC)

31. According to the Fluid Mosaic Model, the plasma membrane is composed of:

- A) A static bilayer of proteins with embedded lipids
- B) A fluid bilayer of phospholipids with embedded proteins
- C) A rigid layer of cellulose with protein channels
- D) A single layer of phospholipids coated with carbohydrates

Answer: A fluid bilayer of phospholipids with embedded proteins

32. Cholesterol in the plasma membrane functions to:

- A) Act as a receptor for hormones
- B) Modulate membrane fluidity and stability
- C) Facilitate active transport of ions
- D) Synthesize ATP

Answer: Modulate membrane fluidity and stability

33. The movement of water across a selectively permeable membrane from a region of low solute concentration to high solute concentration is called:

- A) Simple diffusion
- B) Facilitated diffusion
- C) Osmosis
- D) Active transport

Answer: Osmosis

34. The Na⁺/K⁺ ATPase pump is an example of:

- A) Simple diffusion
- B) Facilitated diffusion
- C) Primary active transport
- D) Secondary active transport

Answer: Primary active transport

35. Receptor-mediated endocytosis often involves the formation of:

- A) Food vacuoles
- B) Contractile vacuoles
- C) Clathrin-coated pits
- D) Autophagosomes

Answer: Clathrin-coated pits

36. The primary cell wall of plants is composed mainly of:

- A) Cellulose microfibrils in a matrix of hemicellulose and pectin
- B) Chitin and proteins
- C) Peptidoglycan
- D) Lignin and suberin

M
K
P
R
E
P
A
R
A
T
I
O
N
S

Answer: Cellulose microfibrils in a matrix of hemicellulose and pectin

37. Plasmodesmata in plant cells function to:

- A) Provide tensile strength
- B) Connect the cytoplasm of adjacent cells for communication
- C) Anchor the cell to the extracellular matrix
- D) Seal cells into leak-proof sheets

Answer: Connect the cytoplasm of adjacent cells for communication

38. Which component of the animal extracellular matrix (ECM) provides tensile strength?

- A) Elastin
- B) Proteoglycans
- C) Collagen
- D) Fibronectin

Answer: Collagen

39. Transmembrane proteins that link the ECM to the intracellular cytoskeleton are called:

- A) Integrins
- B) Cadherins
- C) Connexins
- D) Claudins

Answer: Integrins

40. Tight junctions in animal epithelia primarily function to:

- A) Allow passage of ions and small molecules
- B) Provide strong mechanical adhesion between cells
- C) Create a seal preventing leakage between cells
- D) Anchor cells to the basal lamina

Answer: Create a seal preventing leakage between cells

41. Gap junctions are composed of proteins called:

- A) Integrins
- B) Cadherins
- C) Connexins
- D) Keratins

Answer: Connexins

42. Stem cells that can give rise to all cell types of an organism, including extraembryonic tissues, are termed:

- A) Pluripotent
- B) Multipotent
- C) Totipotent

D) Unipotent

Answer: Totipotent

43. Induced Pluripotent Stem Cells (iPSCs) are created by:

- A) Fertilization of an egg cell
- B) Reprogramming adult somatic cells
- C) Isolating cells from the inner cell mass of a blastocyst
- D) Fusing two different cell types

Answer: Reprogramming adult somatic cells

44. Which of the following is a major DISADVANTAGE of using iPSCs?

- A) They are ethically controversial.
- B) They cannot differentiate into many cell types.
- C) They carry a risk of tumorigenesis due to reprogramming methods.
- D) They are difficult to culture in the lab.

Answer: They carry a risk of tumorigenesis due to reprogramming methods.

45. A major structural difference between plant and animal cells is that plant cells have:

- A) Centrioles
- B) A cell wall and chloroplasts
- C) Lysosomes
- D) A small, numerous vacuoles

Answer: A cell wall and chloroplasts

46. Plant cells typically store carbohydrates in the form of:

- A) Glycogen
- B) Starch
- C) Cellulose
- D) Chitin

Answer: Starch

47. Animal cells typically store carbohydrates in the form of:

- A) Glycogen
- B) Starch
- C) Cellulose
- D) Sucrose

Answer: Glycogen

48. Cytokinesis in plant cells occurs through the formation of a:

- A) Cleavage furrow
- B) Contractile ring of actin and myosin
- C) Cell plate
- D) Septum

Answer: Cell plate

49. Which of the following is generally absent in plant cells?

- A) Mitochondria
- B) Centrioles
- C) Peroxisomes
- D) Ribosomes

Answer: Centrioles

50. The resolution of the naked human eye is approximately:

- A) 0.1 mm
- B) 1 μm
- C) 0.2 nm
- D) 10 nm

Answer: 0.1 mm

51. The Svedberg unit (S) is a measure of a particle's:

- A) Mass
- B) Charge
- C) Sedimentation rate
- D) Density

Answer: Sedimentation rate

52. In a prokaryotic cell, the genetic material is typically:

- A) Multiple linear chromosomes with histones
- B) A single circular chromosome, sometimes with plasmids
- C) Packaged within a double-membrane nucleus
- D) Composed of RNA only

Answer: A single circular chromosome, sometimes with plasmids

53. Binary fission is the method of cell division in:

- A) Animal cells
- B) Plant cells
- C) Prokaryotic cells
- D) Fungal cells

Answer: Prokaryotic cells

54. The outer membrane of mitochondria contains large channel-forming proteins called:

- A) ATP synthases
- B) Porins
- C) Cytochromes
- D) Integrins

Answer: Porins

55. The mitochondrial matrix contains all EXCEPT:

- A) Circular mitochondrial DNA (mtDNA)
- B) 70S ribosomes
- C) Enzymes for the Krebs cycle
- D) Thylakoid membranes

Answer: Thylakoid membranes

56. Thylakoid stacks (grana) are found within which organelle?

- A) Mitochondrion
- B) Chloroplast
- C) Peroxisome
- D) Lysosome

Answer: Chloroplast

57. The light-independent reactions of photosynthesis (Calvin cycle) occur in the:

- A) Thylakoid lumen
- B) Stroma of the chloroplast
- C) Mitochondrial matrix
- D) Cytosol

Answer: Stroma of the chloroplast

58. Which of the following plastids is responsible for photosynthesis?

- A) Chromoplast
- B) Leucoplast
- C) Chloroplast
- D) Amyloplast

Answer: Chloroplast

59. Microfilaments (actin filaments) are primarily involved in:

- A) Organizing the mitotic spindle
- B) Intracellular transport along microtubules
- C) Cell shape, muscle contraction, and cytokinesis
- D) Providing permanent mechanical strength to tissues

Answer: Cell shape, muscle contraction, and cytokinesis

60. Cytochalasin D is a drug that affects the cytoskeleton by:

- A) Stabilizing microtubules
- B) Depolymerizing microfilaments
- C) Disrupting intermediate filaments
- D) Inhibiting dynein motor proteins

Answer: Depolymerizing microfilaments

61. Taxol (paclitaxel) is a drug that acts by:

- A) Depolymerizing microtubules
- B) Stabilizing microtubules, inhibiting their disassembly
- C) Blocking actin polymerization
- D) Inhibiting myosin ATPase

Answer: Stabilizing microtubules, inhibiting their disassembly

62. The nuclear lamina is composed of:

- A) Microtubules
- B) Actin filaments
- C) Intermediate filament proteins (lamins)

M
K
P
R
E
P
A
R
A
T
I
O
N
S

D) Collagen fibers

Answer: Intermediate filament proteins (lamins)

63. Chromatin that is highly condensed and transcriptionally silent is termed:

- A) Euchromatin
- B) Heterochromatin
- C) Nucleosome
- D) Chromosome

Answer: Heterochromatin

64. The endomembrane system does NOT include:

- A) The nuclear envelope
- B) Mitochondria
- C) The Golgi apparatus
- D) Lysosomes

Answer: Mitochondria

65. N-linked glycosylation of proteins begins in the:

- A) Golgi apparatus
- B) Smooth ER
- C) Rough ER
- D) Cytosol

Answer: Rough ER

66. The cisternal maturation model describes a process occurring in the:

- A) Nucleus
- B) Mitochondria
- C) Golgi apparatus
- D) Peroxisomes

Answer: Golgi apparatus

67. Primary lysosomes are formed by budding from the:

- A) Rough ER
- B) Smooth ER
- C) *trans*-Golgi network
- D) Plasma membrane

Answer: *trans*-Golgi network

68. Contractile vacuoles in freshwater protists function primarily for:

- A) Intracellular digestion
- B) Photosynthesis
- C) Osmoregulation
- D) Lipid synthesis

Answer: Osmoregulation

69. The tonoplast is the membrane surrounding the:

- A) Nucleus
- B) Chloroplast
- C) Central vacuole in plant cells

D) Lysosome

Answer: Central vacuole in plant cells

70. The F₀F₁ complex (ATP synthase) is located in the:

- A) Outer mitochondrial membrane
- B) Inner mitochondrial membrane
- C) Mitochondrial matrix
- D) Intermembrane space

Answer: Inner mitochondrial membrane

71. Beta-oxidation of very long-chain fatty acids occurs in:

- A) Mitochondria only
- B) Peroxisomes only
- C) Both mitochondria and peroxisomes
- D) The smooth ER

Answer: Both mitochondria and peroxisomes

72. Which cytoskeletal element is non-polar and provides high tensile strength?

- A) Microtubules
- B) Microfilaments
- C) Intermediate filaments
- D) All of the above

Answer: Intermediate filaments

73. The basal body of a cilium has a microtubule arrangement of:

- A) 9+2
- B) 9+0
- C) 9+1
- D) 8+2

Answer: 9+0

74. The movement of dynein arms walking along adjacent microtubule doublets causes:

- A) Sliding of microtubules, converted to bending by nexin links
- B) Contraction of the basal body
- C) Depolymerization of the axoneme
- D) Formation of the nuclear lamina

Answer: Sliding of microtubules, converted to bending by nexin links

75. Pinocytosis refers to:

- A) Cellular "eating" of large particles
- B) Cellular "drinking" of extracellular fluid
- C) Receptor-mediated uptake of specific ligands
- D) Export of materials via vesicles

Answer: Cellular "drinking" of extracellular fluid

76. Secondary active transport uses the energy from:

- A) ATP hydrolysis directly
- B) Light energy

M
K
P
R
E
P
A
R
A
T
I
O
N
S

C) An electrochemical gradient of another ion (e.g., Na⁺)
D) GTP hydrolysis
Answer: An electrochemical gradient of another ion (e.g., Na⁺)

77. Aquaporins are channel proteins that facilitate the transport of:

- A) Sodium ions
- B) Glucose
- C) Water
- D) Potassium ions

Answer: Water

78. The middle lamella of plant cell walls is rich in:

- A) Cellulose
- B) Lignin
- C) Pectin
- D) Chitin

Answer: Pectin

79. Lignin is a complex polymer that impregnates the _____ of some plant cells, providing rigidity.

- A) Primary cell wall
- B) Secondary cell wall
- C) Plasma membrane
- D) Middle lamella

Answer: Secondary cell wall

80. Hemidesmosomes function to:

- A) Seal adjacent epithelial cells tightly
- B) Anchor epithelial cells to the basal lamina (ECM)
- C) Provide cytoplasmic channels for communication
- D) Facilitate cell-cell adhesion via cadherins

Answer: Anchor epithelial cells to the basal lamina (ECM)

81. Desmosomes are linked intracellularly to:

- A) Microtubules
- B) Microfilaments (actin)
- C) Intermediate filaments (e.g., keratin)
- D) The nuclear lamina

Answer: Intermediate filaments (e.g., keratin)

82. A major function of the glycocalyx is:

- A) ATP synthesis
- B) Cell recognition and adhesion
- C) DNA replication
- D) Protein synthesis

Answer: Cell recognition and adhesion

83. Integrins are involved in:

- A) Mechanotransduction and outside-in

signaling

B) Tight sealing of epithelial layers

C) Forming pores for intercellular communication

D) Digestion of extracellular debris

Answer: Mechanotransduction and outside-in signaling

84. Which of the following is an acellular entity and an exception to cell theory?

- A) Bacterium
- B) Yeast cell
- C) Virus
- D) Red blood cell

Answer: Virus

85. Mature human red blood cells are an exception to cell theory because they:

- A) Lack a nucleus and cannot divide
- B) Have multiple nuclei
- C) Are surrounded by a cell wall
- D) Contain chloroplasts

Answer: Lack a nucleus and cannot divide

86. The maximum useful magnification of a standard light microscope is about:

- A) 100X
- B) 1000-1500X
- C) 10,000X
- D) 1,000,000X

Answer: 1000-1500X

87. In an electron microscope, image formation depends on:

- A) Light absorption
- B) Electron scattering
- C) Refractive index differences
- D) Fluorescence emission

Answer: Electron scattering

88. Super-resolution microscopy techniques (e.g., STED, PALM) can achieve resolutions down to:

- A) 200 nm
- B) 10-20 nm
- C) 2 μ m
- D) 0.2 nm

Answer: 10-20 nm

89. Which of the following techniques separates charged molecules like DNA or proteins based on size and charge?

- A) Chromatography
- B) Electrophoresis
- C) Spectrophotometry

D) Micrometry

Answer: Electrophoresis

90. Tissue culture involves:

- A) Separating organelles by centrifugation
- B) Growing cells on a sterile nutrient medium
- C) Staining tissues for microscopic examination
- D) Measuring light absorption by compounds

Answer: Growing cells on a sterile nutrient medium

91. The bacterial flagellum is powered by:

- A) ATP hydrolysis
- B) A proton (H⁺) gradient and rotary motor
- C) Dynein motor proteins
- D) Myosin-actin interactions

Answer: A proton (H⁺) gradient and rotary motor

92. The archaellum in Archaea is structurally distinct from the bacterial flagellum and is powered by:

- A) Proton motive force
- B) ATP hydrolysis
- C) Sodium ion gradient
- D) Light energy

Answer: ATP hydrolysis

93. Which domain of life includes organisms with cells that often have ether-linked lipids in their membranes?

- A) Bacteria
- B) Archaea
- C) Eukarya
- D) Both Bacteria and Archaea

Answer: Archaea

94. Peptidoglycan is a key component of the cell wall in:

- A) Plants
- B) Fungi
- C) Bacteria
- D) Animals

Answer: Bacteria

95. The cell wall of fungi is primarily composed of:

- A) Cellulose
- B) Chitin
- C) Peptidoglycan
- D) Silica

Answer: Chitin

96. Which of the following is NOT a function of the plasma membrane?

- A) Selective barrier for transport
- B) Cell signaling via receptors

C) Synthesis of ribosomal RNA

D) Cell adhesion and recognition

Answer: Synthesis of ribosomal RNA

97. In the Fluid Mosaic Model, membrane proteins are described as:

- A) Static and immobile
- B) Able to drift laterally in the fluid lipid bilayer
- C) Only located on the external surface
- D) Covalently bound to phospholipids

Answer: Able to drift laterally in the fluid lipid bilayer

98. Facilitated diffusion differs from simple diffusion in that it:

- A) Requires energy from ATP
- B) Moves molecules against their concentration gradient
- C) Uses specific transport proteins (channels or carriers)
- D) Is only for non-polar molecules

Answer: Uses specific transport proteins (channels or carriers)

99. The Sodium-Glucose Symporter in the intestine is an example of:

- A) Simple diffusion
- B) Facilitated diffusion
- C) Primary active transport
- D) Secondary active transport

Answer: Secondary active transport

100. Exocytosis is the process by which cells:

- A) Take in large particles
- B) Secrete macromolecules by vesicle fusion with the plasma membrane
- C) Import fluids and solutes
- D) Degrade internal organelles

Answer: Secrete macromolecules by vesicle fusion with the plasma membrane

101. Polysomes (polyribosomes) are:

- A) Multiple ribosomes translating a single mRNA simultaneously
- B) Aggregates of ribosomal RNA in the nucleolus
- C) Vesicles containing multiple enzymes
- D) Structures within the Golgi apparatus

Answer: Multiple ribosomes translating a single mRNA simultaneously

102. The cell theory was formulated primarily based on the work of:

- A) Hooke, Leeuwenhoek, and Pasteur
- B) Schleiden, Schwann, and Virchow
- C) Brown, Palade, and de Duve

D) Singer, Nicolson, and Mitchell

Answer: Schleiden, Schwann, and Virchow

103. Louis Pasteur's swan-neck flask experiment provided evidence against:

- A) Cell theory
- B) The germ theory of disease
- C) Spontaneous generation for microorganisms
- D) The theory of evolution

Answer: Spontaneous generation for microorganisms

104. August Weismann contributed the concept of _____ to cell theory.

- A) Biogenesis
- B) Common descent
- C) Spontaneous generation
- D) Vitalism

Answer: Common descent

105. Dark-field microscopy is particularly useful for observing:

- A) Internal organelle details
- B) The surface of metal-coated specimens
- C) Motility of living cells
- D) Fluorescently labeled proteins

Answer: Motility of living cells

106. Confocal microscopy improves image clarity by:

- A) Using a laser and pinhole to eliminate out-of-focus light
- B) Employing electrons with very short wavelengths
- C) Staining with heavy metals
- D) Converting phase differences to contrast

Answer: Using a laser and pinhole to eliminate out-of-focus light

107. Differential interference contrast (DIC) microscopy creates images that appear:

- A) Bright on a dark background
- B) Almost three-dimensional
- C) Only in black and white
- D) Based on fluorescence emission

Answer: Almost three-dimensional

108. Scanning Electron Microscopy (SEM) produces images that are:

- A) Two-dimensional internal cross-sections
- B) Three-dimensional topographical views of surfaces
- C) Limited to a resolution of 200 nm
- D) Only of living specimens

Answer: Three-dimensional topographical views of surfaces

109. Which organelle is considered the "powerhouse" of the cell?

- A) Nucleus
- B) Chloroplast
- C) Mitochondrion
- D) Lysosome

Answer: Mitochondrion

110. Which organelle is the site of photosynthesis?

- A) Mitochondrion
- B) Chloroplast
- C) Peroxisome
- D) Vacuole

Answer: Chloroplast

111. The stroma is the internal fluid-filled space of the:

- A) Mitochondrion
- B) Chloroplast
- C) Nucleus
- D) Lysosome

Answer: Chloroplast

112. The inner mitochondrial membrane has a high surface area due to:

- A) Thylakoids
- B) Cristae
- C) Cisternae
- D) Grana

Answer: Cristae

113. Which of the following is a semi-autonomous organelle with its own DNA?

- A) Lysosome
- B) Peroxisome
- C) Golgi apparatus
- D) Mitochondrion

Answer: Mitochondrion

114. The primary cilium functions mainly as a:

- A) Motile structure for fluid movement
- B) Sensory "antenna" for signal reception
- C) Site of protein synthesis
- D) Digestive compartment

Answer: Sensory "antenna" for signal reception

115. Proteins destined for secretion are synthesized by ribosomes on the:

- A) Free ribosomes in the cytosol
- B) Rough ER
- C) Smooth ER
- D) Nuclear envelope

Answer: Rough ER

116. The signal hypothesis explains how proteins destined for the endomembrane system are targeted to the:

- A) Mitochondria
- B) Cytosol
- C) ER
- D) Nucleus

Answer: ER

117. Which of the following modifications of proteins occurs in the Golgi apparatus?

- A) N-linked glycosylation initiation
- B) Further modification of carbohydrate chains (e.g., sulfation)
- C) Initial folding assisted by chaperones like BiP
- D) Transcription of mRNA

Answer: Further modification of carbohydrate chains (e.g., sulfation)

118. Mannose-6-phosphate is a molecular tag that directs proteins to:

- A) The nucleus
- B) Lysosomes
- C) Mitochondria
- D) The plasma membrane for secretion

Answer: Lysosomes

119. Autolysis refers to:

- A) Digestion of foreign bacteria
- B) Digestion of the cell's own organelles
- C) Complete self-digestion of a cell by lysosomal rupture
- D) Breakdown of fatty acids

Answer: Complete self-digestion of a cell by lysosomal rupture

120. Peroxisomes are NOT involved in:

- A) Detoxification of H₂O₂
- B) Beta-oxidation of very long-chain fatty acids
- C) The Krebs cycle
- D) Photorespiration in plant cells

Answer: The Krebs cycle

121. Which of the following structures is part of the cytoskeleton?

- A) Ribosome
- B) Peroxisome
- C) Microtubule
- D) Nucleolus

Answer: Microtubule

122. Kinesin motor proteins generally move cargo toward the _____ end of microtubules.

- A) Plus (growing)
- B) Minus
- C) Both directions equally

D) Neither; they move on microfilaments

Answer: Plus (growing)

123. Myosin motor proteins are associated with which cytoskeletal element?

- A) Microtubules
- B) Microfilaments (actin)
- C) Intermediate filaments
- D) Nuclear lamina

Answer: Microfilaments (actin)

124. Cytoplasmic streaming (cyclosis) in plant cells is driven by interactions between:

- A) Microtubules and kinesin
- B) Actin and myosin
- C) Intermediate filaments and dynein
- D) Tubulin and dynein

Answer: Actin and myosin

125. During cell division, the mitotic spindle is primarily composed of:

- A) Microfilaments
- B) Intermediate filaments
- C) Microtubules
- D) Collagen fibers

Answer: Microtubules

126. The drug colchicine inhibits microtubule polymerization and would thus disrupt:

- A) Muscle contraction
- B) Cytoplasmic streaming
- C) Mitotic spindle formation
- D) Nuclear envelope structure

Answer: Mitotic spindle formation

127. Phalloidin stabilizes microfilaments, while cytochalasin disrupts them. Both affect processes like:

- A) Chromosome separation
- B) Ciliary beating
- C) Cell shape changes and motility
- D) Nuclear transport

Answer: Cell shape changes and motility

128. Which type of cell junction allows for the direct passage of ions and small molecules between animal cells?

- A) Tight junction
- B) Adherens junction
- C) Desmosome
- D) Gap junction

Answer: Gap junction

129. Adherens junctions are linked intracellularly to:

- A) Intermediate filaments
- B) Microfilaments (actin) via catenins

C) Microtubules

D) The nuclear lamina

Answer: Microfilaments (actin) via catenins

130. The extracellular matrix (ECM) of animal cells resists compression largely due to:

A) Collagen fibers

B) Elastin

C) Proteoglycans

D) Fibronectin

Answer: Proteoglycans

131. Which of the following is a major component of plant cell walls but NOT animal ECM?

A) Collagen

B) Cellulose

C) Fibronectin

D) Integrins

Answer: Cellulose

132. A cell placed in a hypertonic solution will likely:

A) Swell and burst

B) Shrink (plasmolyze if plant cell)

C) Remain unchanged

D) Divide rapidly

Answer: Shrink (plasmolyze if plant cell)

133. The movement of oxygen across the plasma membrane occurs via:

A) Simple diffusion

B) Facilitated diffusion

C) Active transport

D) Endocytosis

Answer: Simple diffusion

134. The GLUT transporter that moves glucose into cells is an example of:

A) A channel protein

B) A carrier protein for facilitated diffusion

C) An active transport pump

D) A symporter

Answer: A carrier protein for facilitated diffusion

135. Vesicles budding from the Golgi apparatus can be targeted to different locations based on:

A) Their lipid composition only

B) Molecular "zip codes" (e.g., protein tags)

C) Their size alone

D) Random diffusion

Answer: Molecular "zip codes" (e.g., protein tags)

136. Which organelle is primarily responsible for the synthesis of phospholipids for membrane formation?

A) Rough ER

B) Smooth ER

C) Golgi apparatus

D) Mitochondrion

Answer: Smooth ER

137. The enzymes of the electron transport chain are located in the:

A) Outer mitochondrial membrane

B) Inner mitochondrial membrane

C) Mitochondrial matrix

D) Intermembrane space

Answer: Inner mitochondrial membrane

138. The Calvin cycle fixes carbon dioxide into sugar using ATP and NADPH produced in the:

A) Light-dependent reactions in thylakoids

B) Krebs cycle in the matrix

C) Glyoxylate cycle in glyoxysomes

D) Electron transport chain in the inner membrane

Answer: Light-dependent reactions in thylakoids

139. Coenocytic organisms have bodies that are:

A) Made of a single cell

B) Not divided into separate cells (multinucleate)

C) Prokaryotic in nature

D) Lacking a nucleus

Answer: Not divided into separate cells (multinucleate)

140. The "cell sap" found in the central vacuole of plant cells contains:

A) Hydrolytic enzymes

B) Chlorophyll

C) Water, ions, sugars, pigments

D) Digestive enzymes for autophagy

Answer: Water, ions, sugars, pigments

Chapter 3

Cell Cycle

Introduction to Cell Division

Cell division is the fundamental process where a parent cell divides into two or more daughter cells.

Essential for:

- Growth & Development
- Tissue Repair & Renewal
- Reproduction (asexual & sexual)

Two main types in eukaryotes:

- Mitosis: Produces genetically identical somatic cells
- Meiosis: Produces genetically diverse gametes


Eukaryotic Chromosome Structure & Packaging

Chromosome Composition

- Made of **chromatin** = DNA + proteins (histones & non-histones)
- **Histones** (H2A, H2B, H3, H4): Positively charged proteins for DNA wrapping
- **Non-histone proteins**: Structural & regulatory functions

Levels of Chromatin Packaging

Level	Structure	Description
1°	Nucleosome	146 bp DNA wrapped around histone octamer (2 each: H2A, H2B, H3, H4)
2°	30-nm Fiber	Nucleosomes packed with linker histone H1
3°	Looped Domains	30-nm fibers form loops attached to protein scaffold
4°	Metaphase Chromosome	Maximum condensation via condensin proteins



Chromosome Terminology

- **Diploid (2n)**: Two sets of chromosomes (human somatic cells: 2n=46)



Cell Cycle: One Liners

- The ordered sequence of events from one cell division to the next is the **cell cycle**.
- The **mitotic (M) phase** alternates with the much longer **interphase**.
- **Interphase** accounts for about 90% of the cell cycle and is a period of growth and preparation.
- Interphase is divided into the **G₁ phase** (first gap, cell growth), **S phase** (synthesis, DNA replication), and **G₂ phase** (second gap, preparation for division).
- The **M phase** includes **mitosis** (nuclear division) and **cytokinesis** (cytoplasmic division).
- Cells that are not dividing exit the cycle into a nondividing state called the **G₀ phase**.
- In humans, the entire cell cycle may take about 24 hours, with M phase lasting less than an hour.
- The **S phase** may occupy 10-12 hours, while **G₁** is typically the most variable in length.
- A cell's DNA is called its **genome**; eukaryotic genomes consist of multiple DNA molecules.
- DNA is packaged with proteins into structures called **chromosomes**.
- The entire complex of DNA and proteins is called **chromatin**.
- Each eukaryotic species has a characteristic number of chromosomes in its somatic cells.
- **Human somatic cells** are **diploid (2n)**, with **46 chromosomes** (two sets of 23).
- **Human gametes** are **haploid (n)**, with **23 chromosomes** (one set).
- The two chromosomes of a pair are called **homologous chromosomes (homologs)**; one from each parent.
- **Sex chromosomes** (X and Y) determine sex; the others are **autosomes**.
- Before division, each chromosome is **duplicated** and consists of two identical **sister chromatids**.
- Sister chromatids are joined copies of the original chromosome, held together by **cohesin** proteins.
- The region where sister chromatids are most closely attached is the **centromere**.
- A **kinetochore** is a protein structure that assembles on the centromere and attaches to spindle microtubules.
- When sister chromatids separate, they are considered individual **daughter chromosomes**.
- **Mitosis** is the division of the nucleus, resulting in two genetically identical daughter nuclei.
- It is conventionally broken down into **prophase, prometaphase, metaphase, anaphase, and telophase**.
- **Prophase**: Chromatin condenses into visible chromosomes, nucleoli disappear, mitotic spindle begins to form, centrosomes move apart.
- In animal cells, the **centrosome** organizes the spindle; it contains two **centrioles**.
- Radial arrays of microtubules extending from centrosomes are called **asters**.
- **Prometaphase**: The nuclear envelope fragments, spindle microtubules invade the nuclear area, kinetochores form, and chromosomes are captured by microtubules.
- **Metaphase**: Chromosomes align at the **metaphase plate**, an imaginary plane equidistant between the spindle poles.
- **Anaphase**: Sister chromatids separate (cohesins are cleaved) and move to opposite poles as individual chromosomes; the cell elongates.
- **Telophase**: Daughter nuclei form, nuclear envelopes re-form, chromosomes decondense, nucleoli reappear.
- **Cytokinesis** in animal cells occurs via a **cleavage furrow** created by a **contractile ring** of actin and myosin.
- **Cytokinesis** in plant cells occurs via **cell plate** formation from Golgi-derived vesicles.
- The **mitotic spindle** is a structure made of microtubules and associated proteins that controls chromosome movement.
- Spindle microtubules **polymerize** (elongate) and **depolymerize** (shorten) by gaining or losing tubulin subunits.
- **Kinetochore microtubules** attach to kinetochores and move chromosomes.

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. What is the primary purpose of mitosis in multicellular organisms?

- A) To produce genetically diverse gametes
- B) To reduce the chromosome number by half
- C) To promote growth, tissue repair, and asexual reproduction
- D) To allow for genetic recombination through crossing over

Answer: To promote growth, tissue repair, and asexual reproduction

2. During which phase of the cell cycle does DNA replication occur?

- A) G₁ phase
- B) G₂ phase
- C) M phase
- D) S phase

Answer: S phase

3. Which protein forms a contractile ring during cytokinesis in animal cells?

- A) Tubulin
- B) Actin
- C) Keratin
- D) Collagen

Answer: Actin

4. What is the term for the paired maternal and paternal chromosomes that carry the same genes?

- A) Sister chromatids
- B) Homologous chromosomes
- C) Sex chromosomes
- D) Recombinant chromosomes

Answer: Homologous chromosomes

5. The point of constriction on a chromosome that holds sister chromatids together is called the:

- A) Kinetochore
- B) Telomere
- C) Centromere
- D) Centrosome

Answer: Centromere

6. Which of the following is a key feature of prophase I of meiosis that does NOT occur in mitosis?

- A) Chromosome condensation
- B) Breakdown of the nuclear envelope
- C) Synapsis and crossing over
- D) Formation of the mitotic spindle

Answer: Synapsis and crossing over

7. What is the haploid (n) number of chromosomes in a human somatic cell?

- A) 23
- B) 46
- C) 92
- D) 2

Answer: 23

8. The enzyme that adds DNA sequences to telomeres to counteract shortening is:

- A) DNA polymerase
- B) Telomerase
- C) Ligase
- D) Primase

Answer: Telomerase

9. Which checkpoint ensures that all chromosomes are properly attached to the spindle before anaphase?

- A) G₁/S checkpoint
- B) G₂/M checkpoint
- C) Spindle assembly checkpoint
- D) Post-replication checkpoint

Answer: Spindle assembly checkpoint

10. What is the correct order of the mitotic phases?

- A) Prophase, Metaphase, Anaphase, Telophase, Prometaphase
- B) Prophase, Prometaphase, Metaphase, Anaphase, Telophase
- C) Interphase, Prophase, Metaphase, Anaphase, Telophase
- D) Prophase, Anaphase, Metaphase, Telophase, Cytokinesis

Answer: Prophase, Prometaphase, Metaphase, Anaphase, Telophase

11. Binary fission in bacteria is most similar to which eukaryotic process?

- A) Mitosis
- B) Meiosis I
- C) Meiosis II
- D) Cytokinesis

Answer: Mitosis

12. Which protein complex holds sister chromatids together after DNA replication?

- A) Condensin
- B) Cohesin
- C) Kinetochore
- D) Securin

Answer: Cohesin



Chapter 4

Biological Molecules & Enzymes

Biochemistry is the branch of science that explores the **chemical processes and substances occurring within living organisms**. It serves as the bridge between biology and chemistry, explaining life at a molecular level.

Foundational Principles of Biochemistry

- Cellular Basis:** All biochemical processes occur within or are mediated by cells.
- Hierarchy of Structure:** Atoms → Small Molecules → Monomers → Polymers → Supramolecular Complexes → Organelles → Cells.
- Structure-Function Relationship:** The three-dimensional shape of a biomolecule (its **conformation**) is directly linked to its biological function. Denaturation (loss of shape) leads to loss of function.
- Metabolism:** Living organisms transform energy and matter through a vast network of chemical reactions (pathways).
- Homeostasis:** Biochemical systems are tightly regulated to maintain a stable internal environment despite external changes.
- Information Flow:** Genetic information flows from **DNA → RNA → Protein** (The Central Dogma). This information directs all cellular activities.

Metabolism

Metabolism is the **totality of an organism's chemical reactions**. It is an emergent property of life that manages the material and energy resources of the cell through intricate, enzyme-catalyzed pathways.

Two Complementary Sides of Metabolism

Aspect	Catabolism	Anabolism (Biosynthesis)
Core Concept	Breakdown pathways.	Build-up pathways.
Energy Relationship	Exergonic: Releases energy. Some is captured as ATP; the rest is released as heat.	Endergonic: Consumes energy. Driven by ATP hydrolysis.
Redox Relationship	Oxidative: Releases electrons, often captured by carriers like NAD ⁺ (forming NADH).	Reductive: Consumes electrons, often from carriers like NADPH.
Carbon Flow	Complex molecules (carbs, fats) → Smaller, simpler molecules (CO ₂ , lactate, ethanol).	Simple precursors (amino acids, sugars) → Complex macromolecules (proteins, polysaccharides).
Primary Goal	1. Generate usable energy (ATP, reducing power). 2. Create precursor metabolites for biosynthesis.	1. Synthesize cellular components for growth and repair. 2. Store energy for later use.
Examples	Glycolysis, Krebs (TCA) Cycle, β-Oxidation of fats, Cellular Respiration.	Protein synthesis, Glycogenesis, DNA Replication, Gluconeogenesis.

Metabolic Concepts

- Metabolic Pathway:** A series of linked, enzyme-catalyzed reactions where the product of one reaction becomes the substrate for the next.
- ATP (Adenosine Triphosphate):** The universal "**energy currency**" of the cell. Energy from catabolism is used to phosphorylate ADP into ATP. Anabolic processes hydrolyze ATP back to ADP + Pi, releasing energy to drive reactions.
- Redox Coenzymes:** Key electron carriers.

M
K
P
R
E
P
A
R
A
T
I
O
N
S



M
K

P
R
E
P
A
R
A
T
I
O
N
S

- **NAD⁺/NADH & FAD/FADH₂:** Primarily involved in **catabolic** reactions, carrying electrons to the electron transport chain for ATP synthesis.
- **NADP⁺/NADPH:** Primarily involved in **anabolic** reactions (e.g., fatty acid synthesis, photosynthesis), providing reducing power for biosynthesis.
- **Regulation:** Metabolic pathways are precisely controlled via:
 - **Allosteric Regulation:** A molecule binds at a site other than the active site, changing enzyme activity (e.g., feedback inhibition).
 - **Covalent Modification:** Reversible addition/removal of chemical groups (e.g., phosphorylation).
 - **Compartmentalization:** Separating opposing pathways into different organelles (e.g., fatty acid breakdown in mitochondria vs. synthesis in cytoplasm).
- **Metabolic Disorders:** Diseases resulting from defects in enzymes or pathways (e.g., Phenylketonuria (PKU), Diabetes Mellitus, Mitochondrial diseases).

Laws of Thermodynamics in Biochemistry

Thermodynamics governs **energy transformations** in biochemical systems, determining whether reactions can occur spontaneously.

The First Law: Conservation of Energy

- **Statement:** Energy can be **neither created nor destroyed**, only converted from one form to another.
- **In Biochemistry:** The total energy of a cell and its surroundings is constant. Chemical potential energy stored in nutrients (glucose, fats) is converted into other forms:
 - **Work:** Mechanical (muscle contraction), transport (active transport), chemical (biosynthesis).
 - **Heat:** A byproduct of inefficient energy transfers; used to maintain body temperature in endotherms.
 - **Storage:** In energy-rich bonds of ATP or as chemical bonds in macromolecules.

The Second Law: Increasing Entropy

- **Statement:** In any spontaneous process, the **total entropy (disorder) of the universe always increases**.
- **Key Concepts:**
 - **Entropy (S):** A measure of randomness or disorder. Systems tend toward greater disorder.
 - **Spontaneous Process:** One that can occur without an ongoing input of external energy (does **not** imply it will be fast; kinetics vs. thermodynamics).
- **In Biochemistry:**
 - Living organisms are **highly ordered (low entropy) systems**. They maintain this order by **constantly increasing the entropy of their surroundings**.
 - They do this by releasing heat and simple waste products (like CO₂ and H₂O) into the environment, which are more disordered than the complex nutrients consumed.
 - **Local order is purchased at the cost of universal disorder.**

The Central Concept: Gibbs Free Energy (G)

Gibbs Free Energy (ΔG) combines the first and second laws into a single, practical measure for predicting reaction spontaneity **at constant temperature and pressure**.

- **Equation:** $\Delta G = \Delta H - T\Delta S$
 - ΔG : Change in free energy (usable energy).
 - ΔH : Change in enthalpy (total heat content; bond energy).
 - T : Absolute Temperature (in Kelvin).
 - ΔS : Change in entropy.
- **Interpreting ΔG :**



- $\Delta G < 0$ (**Negative**): The reaction is **exergonic** and **spontaneous**. It releases free energy (e.g., ATP hydrolysis, fuel oxidation).
- $\Delta G > 0$ (**Positive**): The reaction is **endergonic** and **non-spontaneous**. It requires an input of free energy (e.g., protein synthesis, gluconeogenesis).
- $\Delta G = 0$: The reaction is at **equilibrium**; no net change occurs.

Biochemical Applications of Thermodynamics

1. **ATP as an Energy Coupler:** An endergonic reaction ($\Delta G > 0$) can be driven by coupling it to the highly exergonic hydrolysis of ATP ($\Delta G \ll 0$), making the **net ΔG for the coupled process negative**.
2. **Kinetics vs. Thermodynamics:** A negative ΔG means a reaction *can* happen, but **not** how *fast* it will happen. **Enzymes** are biological catalysts that **lower the activation energy (E_a)**, speeding up thermodynamically favorable reactions to biologically useful rates.
3. **Redox Reactions:** The transfer of electrons from a reducing agent (electron donor) to an oxidizing agent (electron acceptor) releases free energy (exergonic). This energy is harnessed in the electron transport chain to pump protons and synthesize ATP.

Importance of Water

Water is the universal solvent of life and the most abundant molecule in living cells, constituting 70-90% of cell mass. Its unique chemical and physical properties are not merely a background environment but are **central to the structure, function, and very existence** of all biomolecules and biochemical processes.

I. Key Properties of Water that Dictate its Biological Role

A. Polarity and Hydrogen Bonding

- **Molecular Structure:** A water molecule (H_2O) has a bent geometry with oxygen at the center. Oxygen is more electronegative than hydrogen, creating a **polar covalent bond**.
- **Dipole Moment:** This results in a partial negative charge (δ^-) on the oxygen and partial positive charges (δ^+) on the hydrogens.
- **Hydrogen Bonds:** The δ^+ hydrogen of one water molecule is strongly attracted to the δ^- oxygen of another. Each water molecule can form up to **four hydrogen bonds** in a tetrahedral arrangement.
 - **Consequence:** This extensive, dynamic H-bonding network is responsible for water's high cohesiveness, surface tension, and unique thermal properties.

II. The Multifaceted Roles of Water in Biological Systems

A. The Solvent of Life (The Universal Aqueous Medium)

Water's polarity makes it an excellent solvent for other polar and charged molecules (**hydrophilic** substances).

- **Mechanism:** Water molecules surround solutes, forming **hydration shells**. Positive ions (cations) are surrounded by water's δ^- oxygen; negative ions (anions) by the δ^+ hydrogens.
- **Biological Impact:**
 - **Dissolves & Transports:** Enables the dissolution and circulation of nutrients (glucose, amino acids, ions), gases (O_2 , CO_2), and waste products throughout organisms (blood, sap, cytosol).
 - **Facilitates Reactions:** Brings reactants together in solution, allowing metabolic reactions to occur.

B. Thermal Regulator

Water has an exceptionally **high specific heat capacity**, **high heat of vaporization**, and **high heat of fusion**.

- **High Specific Heat:** It takes a large amount of heat to raise water's temperature. This means water **buffers cells and organisms against rapid temperature changes**, maintaining a stable internal environment for enzyme function.



- **Specific Heat Capacity of Water:** The amount of heat energy (in joules) required to raise the temperature of 1 gram of a substance by 1 degree Celsius. Water has an exceptionally **high specific heat capacity** of **4.184 J/g·°C** (or 1 cal/g·°C). This is among the highest of all common liquids.
- **Heat of Vaporization of Water:** The amount of heat energy required to convert 1 gram of a liquid into vapor at its boiling point without a temperature change. Water has an exceptionally **high heat of vaporization** of **~2260 J/g** (or ~540 cal/g). This is remarkably high compared to most other liquids. Evaporation of water (e.g., sweating, panting) requires significant heat energy, providing a powerful **cooling mechanism** for organisms.
- **Biological Impact:** Essential for **homeostasis** (temperature regulation) and allows life to exist in varied climates.

C. Reactant and Product in Biochemical Reactions

Water is a direct participant in key metabolic reactions.

- **Hydrolysis Reactions:** Water is used to **break** bonds, with its -H and -OH added across the broken bond. Crucial for:
 - Digestion of macromolecules (proteins → amino acids, polysaccharides → sugars).
 - Breaking down ATP ($\text{ATP} + \text{H}_2\text{O} \rightarrow \text{ADP} + \text{P}_i + \text{energy}$).
- **Dehydration Synthesis (Condensation) Reactions:** The reverse of hydrolysis. Biomolecules are **synthesized** when monomers join, releasing a water molecule. Essential for building proteins, nucleic acids, and polysaccharides.

D. Structural Role through Hydrophobic Interactions

Water's interaction with **nonpolar (hydrophobic)** molecules is equally critical.

- **The Hydrophobic Effect:** Nonpolar molecules (e.g., lipids, hydrophobic amino acid side chains) cannot form H-bonds with water. To minimize disruption of the water's H-bonding network, water molecules reorganize into a more ordered "cage" around them, which is thermodynamically unfavorable.
- **Biological Impact:** This effect is the **primary driving force** for:
 - **Membrane Formation:** Lipids spontaneously aggregate to form bilayers and micelles, minimizing contact with water.
 - **Protein Folding:** Hydrophobic amino acids are driven to the interior of the protein, defining its 3D structure.
 - **DNA Base Stacking:** Stabilizes the double helix.

E. Cohesion, Adhesion, and Surface Tension

- **Cohesion:** Hydrogen bonding between water molecules creates high **cohesive forces**.
- **Adhesion:** Water's polarity allows it to **adhere** to other polar surfaces (e.g., plant cell walls).
- **Biological Impact:**
 - **Capillary Action:** Cohesion and adhesion together allow water to move against gravity in narrow tubes (e.g., xylem vessels in plants).
 - **Surface Tension:** Creates a habitat (e.g., for some insects) and is important in lung alveoli.

F. Ionization and pH

Water undergoes slight autoionization: $\text{H}_2\text{O} \rightleftharpoons \text{H}^+ + \text{OH}^-$.

- **The Foundation of pH:** The concentration of H^+ (protons) defines the acidity or basicity of a solution. The **pH scale** is central to biochemistry.
- **Biological Impact:** Enzymes and biomolecules are exquisitely sensitive to pH. Water is the medium that allows the establishment of **proton gradients**, which are crucial for:
 - **ATP Synthesis:** The proton motive force drives ATP synthase in mitochondria and chloroplasts.
 - **Cellular Respiration & Photosynthesis:** Chemiosmosis depends on gradients established across membranes.

Property of Water	Chemical Basis	Biological Consequence
Excellent Solvent	Polarity & H-bonding	Dissolves & transports nutrients/wastes; medium for reactions.
High Heat Capacity	Extensive H-bonding	Buffers temperature, maintains homeostasis.
High Heat of Vaporization	H-bonding must be broken	Effective evaporative cooling (sweating).
Cohesion/Adhesion	H-bonding	Capillary action in plants; surface tension.
Participates in Reactions	Polar, can be split	Reactant in hydrolysis ; product in dehydration synthesis .
Density of Ice < Liquid	H-bonds form a crystalline lattice	Ice floats, insulating aquatic life below.

The Importance of Carbon: The Element of Life

Carbon (C) is the fundamental element of all known life on Earth. Its unique chemical properties make it the indispensable backbone for the structure and function of every biological molecule. Biochemistry is, in essence, the chemistry of carbon compounds.

1. Core Reasons for Carbon's Central Role

Property of Carbon	Consequence for Biochemistry
Tetravalency (4 Covalent Bonds)	A carbon atom can form four stable covalent bonds simultaneously. This allows it to link to other carbons and various other atoms (H, O, N, S, P), creating complex, multi-dimensional molecular skeletons.
Bond Strength & Stability	C-C bonds and C-H bonds are very strong, providing stability to large organic molecules. However, they are not <i>unbreakably</i> strong, allowing for controlled reactions in metabolism.
Versatility in Bonding	Carbon can form single (σ) , double (π) , and triple bonds with itself and other elements. This creates a vast diversity of molecular shapes and functional groups (e.g., alkenes, carbonyls).
Ability to Form Long Chains & Rings (Catenation)	Carbon atoms bond together to form stable, extended chains (linear or branched) and cyclic/ring structures . This is the foundation for macromolecule diversity.
Moderate Electronegativity	Carbon is neither strongly electropositive nor electronegative (Pauling EN = 2.55). This makes its bonds with other biogenic elements (H, O, N) relatively non-polar or only moderately polar , preventing molecules from being too ionic or reactive in water.

2. Key Chemical Concepts Derived from Carbon's Properties

- **Molecular Diversity & Complexity:** The combination of tetravalency and catenation leads to an almost infinite number of possible carbon-based (**organic**) compounds. This provides the **structural foundation** for the immense variety of biomolecules.
- **Isomerism:** Carbon's tetrahedral geometry enables the existence of **isomers**—molecules with the same molecular formula but different structures.
 - **Structural Isomers:** Different bonding arrangements (e.g., glucose vs. fructose).
 - **Stereoisomers (Enantiomers):** Mirror-image molecules crucial in biology. Life almost exclusively uses **L-amino acids** and **D-sugars**, demonstrating carbon-based **molecular chirality**.

- Functional Groups:** Specific, reactive clusters of atoms (often involving carbon bonded to O, N, S, P) attached to carbon skeletons. They determine a molecule's **chemical properties and reactivity**. Common biochemical functional groups include:
 - Hydroxyl (-OH):** Alcohols, sugars (polar, forms H-bonds).
 - Carbonyl (>C=O):** Aldehydes & ketones (present in sugars).
 - Carboxyl (-COOH):** Organic acids, amino acids (acidic, donates H⁺).
 - Amino (-NH₂):** Amines, amino acids (basic, accepts H⁺).
 - Phosphate (-OPO₃²⁻):** Nucleotides, ATP (energetic, acidic).
 - Sulfhydryl (-SH):** Thiols, cysteine (forms disulfide bonds in proteins).

3. Role of Carbon in the Four Major Classes of Biomolecules

Biomolecule Class	Carbon's Role	Key Examples
1. Carbohydrates	Forms the sugar backbone . Carbon atoms in chains or rings bonded to -OH groups. Provide energy (fuel) , structural support , and cellular recognition.	Glucose (C₆H₁₂O₆): Primary energy molecule. Cellulose: Structural polymer in plants (β-glucose chains).
2. Lipids	Forms the hydrocarbon chains (fatty acids) and steroid ring systems . Carbon's non-polar C-C and C-H bonds create hydrophobic properties essential for membranes and energy storage.	Fatty Acids: Long C-H chains (e.g., palmitic acid). Cholesterol: Four fused carbon rings.
3. Proteins	Forms the amino acid backbone . Every amino acid has a central α-carbon bonded to an amino group, a carboxyl group, and a variable side chain (R group). Carbon diversity in R groups dictates protein function.	All 20 Amino Acids: Differ in their R groups attached to the α-carbon. Polypeptide Chain: A polymer of amino acids linked by peptide (C-N) bonds.
4. Nucleic Acids	Forms the pentose sugar (ribose/deoxyribose) and the nitrogen bases (purines, pyrimidines) in nucleotides. The genetic code is stored in sequences of carbon-based nucleotides.	DNA Nucleotides: Deoxyribose (C5 sugar) + base (A,T,G,C). ATP: The primary "energy currency" of the cell is a carbon-based nucleotide.

4. Carbon in Metabolism and the Energy Cycle

- Carbon as an Energy Carrier:** The chemical energy in food is stored in the bonds of carbon-based molecules (e.g., C-C and C-H bonds in glucose and fats). **Cellular respiration** is fundamentally the **controlled, stepwise oxidation of carbon** (from C-H/C-C bonds to C=O bonds in CO₂), releasing energy captured as ATP.
- The Carbon Cycle:** Carbon atoms are **recycled** through the biosphere. **Autotrophs** (plants) fix inorganic carbon (CO₂) into organic molecules (e.g., glucose via photosynthesis). **Heterotrophs** (animals) consume and metabolize these organic molecules, returning CO₂ to the atmosphere. This biogeochemical cycle is powered by carbon's chemistry.

The Centrality of Carbon in Biochemistry

Aspect	Explanation	Biological Implication
Structural Backbone	Forms stable chains & rings.	Creates diverse skeletons for all biomolecules.



Functional Diversity	Binds to key biogenic elements (H,O,N,P,S).	Enables creation of functional groups that define reactivity.
Isomerism & Chirality	Tetrahedral geometry allows 3D variation.	Basis for molecular specificity (e.g., enzyme-substrate, receptor-ligand).
Energy Transactions	Stable C-C/C-H bonds store energy; oxidation releases it.	Carbon compounds are the universal fuel and energy currency of life.
Information Storage	Forms stable, sequenceable polymers (nucleotides).	Genetic information (DNA/RNA) is stored in carbon-based code.

M
K
P
R
E
P
A
R
A
T
I
O
N
S

4. Biological Molecules & Enzymes

Biological Molecules

Carbohydrates

Carbohydrates are **polyhydroxy aldehydes or ketones**, or substances that hydrolyze to yield such compounds. They are the most abundant biomolecules on Earth and serve as central molecules in biochemical pathways. Biochemically, they are defined by their empirical formula $(CH_2O)_n$, where $n \geq 3$, though derivatives may contain nitrogen, sulfur, or phosphorus.

2. Sources & Biochemical Origin

Anabolism (Synthesis):

- **Photosynthesis (Calvin Cycle):** Occurs in the stroma of chloroplasts. Uses ATP and NADPH from light reactions to fix CO_2 into **3-phosphoglycerate (3-PG)**, which is converted to triose phosphates and eventually to glucose-6-phosphate and starch. The key enzyme is **Ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO)**.
- **Gluconeogenesis:** The 11-step pathway (largely a reversal of glycolysis with three bypass reactions) that synthesizes glucose from non-carbohydrate precursors.
- **Glycogenesis:** The synthesis of glycogen from glucose-1-phosphate in liver and muscle cells, regulated by **glycogen synthase** and branching enzyme.

Catabolism (Breakdown) & Dietary Sources:

- **Digestion:** Enzymes like **α -amylase** (saliva/pancreas), **sucrase-isomaltase**, **lactase**, and **maltase** hydrolyze dietary polysaccharides and disaccharides into absorbable monosaccharides (glucose, fructose, galactose).
- **Metabolic Pathways:**
 - **Glycolysis:** The 10-step cytosolic pathway converting **glucose to pyruvate**, yielding 2 ATP and 2 NADH per glucose.

3. Properties (Biochemical Perspective)

Structural & Isomeric Properties:

- **Chirality & Stereoisomerism:**
 - **Enantiomers:** Non-superimposable mirror images (D- vs. L-). Biochemistry almost exclusively uses **D-sugars**.
 - **Diastereomers:** Non-mirror image stereoisomers. **Epimers** are a special subclass differing at only one chiral center (e.g., glucose and galactose are C-4 epimers).
- **Anomerism:** A type of diastereomerism specific to the **anomeric carbon** (C-1 in aldoses, C-2 in ketoses) upon ring formation. **α -anomer** has the -OH group *trans* to the CH_2OH group (axial in glucose). **β -anomer** has it *cis* (equatorial in glucose). Interconversion in solution is **mutarotation**.
- **Ring Structures:** Predominant forms are **pyranose** (6-membered, chair conformation) and **furanose** (5-membered). Chair conformation stability is governed by sterics: bulky groups (like -OH) prefer equatorial positions.

Chemical Reactivity:

- **The Anomeric Carbon:** The most reactive center in a sugar. It is involved in:

- **Glycosidic Bond Formation:** Creates acetal/ketal linkages with -OH groups of other molecules.
- **Reducing Ability:** The free anomeric carbon in linear form reduces metal ions ($\text{Cu}^{2+} \rightarrow \text{Cu}^+$ in Benedict's test). Sucrose is non-reducing because both anomeric carbons are involved in its glycosidic bond.

Physical Properties:

- **Solubility & Osmolarity:** High solubility and low molecular weight of monosaccharides create high osmotic pressure, which is why cells store glucose as large, insoluble polymers (glycogen, starch).

4. Classification & Structural Biochemistry

I. Monosaccharides

Monosaccharides are the simplest carbohydrate units that cannot be hydrolyzed into smaller carbohydrate molecules. They are the monomers from which all more complex carbohydrates (disaccharides, oligosaccharides, polysaccharides) are built.

- **Empirical Formula:** Typically $(\text{CH}_2\text{O})_n$, where $n = 3-9$ (most commonly 3, 5, or 6).
- **Functional Groups:** They are **polyhydroxy aldehydes** (aldoses) or **polyhydroxy ketones** (ketoses).
- **Biochemical Role:** Serve as:
 - Primary fuel molecules (e.g., glucose).
 - Metabolic intermediates (e.g., fructose-1,6-bisphosphate).
 - Building blocks for polymers (e.g., glycogen, cellulose, chitin).
 - Precursors for other biomolecules (e.g., ribose for nucleic acids).

Classification of Monosaccharides

Monosaccharides are classified based on three key features:

A. By the Number of Carbon Atoms

Carbon Count	Name	Examples (Biologically Important)
3	Triose	Glyceraldehyde (aldotriose), Dihydroxyacetone (ketotriose)
4	Tetrose	Erythrose, Threose (intermediates in PPP)
5	Pentose	Ribose (RNA), 2-Deoxyribose (DNA), Ribulose , Xylulose (PPP)
6	Hexose	Glucose , Galactose , Mannose , Fructose
7	Heptose	Sedoheptulose (intermediate in PPP and Calvin cycle)
9	Nonose	Neuraminic acid (precursor of sialic acids)

B. By the Nature of the Carbonyl Group

1. **Aldoses:** Contain an **aldehyde group** ($-\text{CHO}$) at carbon 1 (C-1).
 - General formula: $\text{H}-(\text{CHOH})_n-\text{CHO}$
 - Examples: Glyceraldehyde, Ribose, Glucose, Galactose.
2. **Ketoses:** Contain a **ketone group** ($>\text{C}=\text{O}$) at carbon 2 (C-2), typically.
 - General formula: $\text{H}-(\text{CHOH})_n-\text{CO}-(\text{CHOH})_m-\text{H}$
 - Examples: Dihydroxyacetone, Ribulose, Fructose.

Conformational Structures

- **Pyranose Rings:** Adopt **chair conformations** (${}^1\text{C}_4$ or ${}^4\text{C}_1$).
 - ${}^1\text{C}_4$: The anomeric carbon (C-1) is axial.
 - ${}^4\text{C}_1$: The anomeric carbon is equatorial (more stable for D-sugars as bulky groups prefer equatorial positions).
- **Furanose Rings:** Adopt **envelope or twist conformations**.

Physical & Chemical Properties

Physical Properties

- **State:** Colorless, crystalline solids.
- **Solubility:** Highly soluble in water due to extensive hydrogen bonding via -OH groups.

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

Page 8 | 59

- **Sweetness:** Varies; fructose > sucrose > glucose > galactose > maltose > lactose.
- **Optical Activity:** All monosaccharides (except dihydroxyacetone) are optically active, rotating plane-polarized light. Specific rotation ($[\alpha]_D$) is a characteristic property (e.g., D-glucose = $+52.7^\circ$ at equilibrium).

Chemical Properties

1. **Reducing Sugars:** All monosaccharides are **reducing sugars** because their anomeric carbon is free (in equilibrium with the open-chain aldehyde or ketone form). They can reduce agents like:
 - **Fehling's/Benedict's Reagent** ($\text{Cu}^{2+} \rightarrow \text{Cu}^+$, forming a brick-red precipitate).
 - **Tollens' Reagent** ($\text{Ag}^+ \rightarrow \text{Ag}^0$, silver mirror).
2. **Esterification:** -OH groups can form esters with acids (e.g., phosphate, sulfate, acetate).
3. **Glycoside Formation:** The anomeric -OH reacts with an alcohol or amine of another molecule, forming a **glycosidic bond** and releasing water. This is the fundamental reaction for forming disaccharides and polysaccharides.
4. **Oxidation & Reduction:** As described in derivatives (to acids or alcohols).

Monosaccharides at a Glance

Name	Type	Significant Isomerism	Primary Biological Role
D-Glucose	Aldohexose	C-2 epimer of Mannose; C-4 epimer of Galactose	Central metabolic fuel; monomer of major polysaccharides.
D-Fructose	Ketohexose	Anomer of Glucose (in sucrose)	Dietary sugar; very sweet; metabolized in liver.
D-Galactose	Aldohexose	C-4 epimer of Glucose	Component of lactose and many glycoconjugates.
D-Mannose	Aldohexose	C-2 epimer of Glucose	Component of N-linked glycans.
D-Ribose	Aldopentose	-	Backbone of RNA and nucleotide cofactors (ATP, NADH).
2-Deoxy-D-Ribose	Aldopentose	Lacks O at C-2 vs. Ribose	Backbone of DNA.

II. Disaccharides

Disaccharides

1. Introduction & Definition

Disaccharides are carbohydrates formed when **two monosaccharide units** are joined by a **glycosidic bond** through a **dehydration (condensation) reaction**. They are the simplest type of **oligosaccharide**.

- **General Reaction:**
 $\text{Monosaccharide-OH} + \text{HO-Monosaccharide} \rightarrow \text{Disaccharide} + \text{H}_2\text{O}$
- **Molecular Formula:** Typically $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ (for two hexoses minus one water).

2. Formation & Structure: The Glycosidic Bond




A. Nature of the Glycosidic Bond

- **Formation:** A covalent bond between the **anomeric carbon** (C-1 of an aldose or C-2 of a ketose) of one monosaccharide and a **hydroxyl group** (-OH) of another.

B. Reducing vs. Non-Reducing Disaccharides

This classification is **fundamental** and depends on whether the anomeric carbon of the second monosaccharide is free.

- **Reducing Disaccharides:**





M
K

P
R
E
P
A
R
A
T
I
O
N
S

- **Feature:** One anomeric carbon is involved in the glycosidic bond; the other remains **free** and can interconvert between α and β forms (mutarotation).
- **Chemical Behavior:** They exhibit **reducing properties** (give positive Benedict's/Fehling's test) because the free anomeric carbon can open to reveal an aldehyde or ketone group.
- **Examples:** Maltose, Lactose, Cellobiose.
- **Non-Reducing Disaccharides:**
 - **Feature:** **Both anomeric carbons** are involved in the glycosidic bond (a "head-to-head" linkage).
 - **Chemical Behavior:** **No free anomeric carbon**, therefore **no mutarotation** and **no reducing properties** (negative Benedict's test).
 - **Example:** Sucrose

Key Disaccharides

Disaccharide	Monosaccharide Components	Glycosidic Bond	Reducing?	Major Source / Role	Digestive Enzyme
Sucrose	Glucose + Fructose	$\alpha(1\rightarrow2\beta)$	No	Table sugar; Transport in plants	Sucrase
Lactose	Galactose + Glucose	$\beta(1\rightarrow4)$	Yes	Milk sugar	Lactase
Maltose	Glucose + Glucose	$\alpha(1\rightarrow4)$	Yes	Starch digestion; Germinating grains	Maltase
Cellobiose	Glucose + Glucose	$\beta(1\rightarrow4)$	Yes	Cellulose hydrolysis (not digestible by humans)	(Cellulase - absent)
Trehalose	Glucose + Glucose	$\alpha(1\rightarrow1)\alpha$	No	Insect blood sugar; Stress protectant in fungi	Trehalase

III. Oligosaccharides

3-20 monosaccharide units. **N-linked glycans** are attached via the amide nitrogen of Asn in the sequon **Asn-X-Ser/Thr**. Synthesis involves a **dolichol phosphate** lipid carrier in the ER.

IV. Polysaccharides (Glycans)

Polysaccharides are high-molecular-weight carbohydrates composed of **long chains of monosaccharide units** (typically >10) linked together by **glycosidic bonds**. They are also called **glycans**.

- **General Formula:** $(C_6H_{10}O_5)_n$ where n = number of monosaccharide units (ranges from hundreds to tens of thousands).
- **Nature:** They are **polymers** and constitute one of the main classes of biological macromolecules.
- **Biochemical Role:** Primarily serve as:
 - **Storage forms of energy** (e.g., starch, glycogen).
 - **Structural materials** (e.g., cellulose, chitin).
 - **Key components of cell recognition and signaling systems** (e.g., glycosaminoglycans in extracellular matrix).

Classification of Polysaccharides

Polysaccharides are classified based on composition, structure, and function.

A. Based on Monosaccharide Composition

1. **Homopolysaccharides (Homoglycans):** Composed of only **one type** of monosaccharide unit.
 - *Examples:* Starch, Glycogen, Cellulose (all of glucose), Chitin (of N-acetylglucosamine).
2. **Heteropolysaccharides (Heteroglycans):** Composed of **two or more different** types of monosaccharides or their derivatives.
 - *Examples:* Hyaluronic acid, Heparin, Peptidoglycan.

B. Based on Biological Function

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091



M
K

P
R
E
P
A
R
A
T
I
O
N
S

- Storage Polysaccharides:** Compact, often branched structures that are hydrolyzed to release sugar monomers when energy is needed. They are **osmotically inert**.
 - Examples:* Starch (plants), Glycogen (animals).
- Structural Polysaccharides:** Typically form fibrous, water-insoluble structures that provide mechanical support and protection.
 - Examples:* Cellulose (plants), Chitin (arthropods/fungi), Peptidoglycan (bacteria).

C. Based on Chain Structure

- Linear (Unbranched) Polysaccharides:** The monosaccharide units are linked in a continuous straight chain.
 - Examples:* Cellulose, Amylose, Chitin.
- Branched Polysaccharides:** The main chain has side chains attached at branching points.
 - Examples:* Amylopectin, Glycogen, Glycosaminoglycans (GAGs).

Structural Features & Chemical Properties

A. General Structural Features

- Glycosidic Linkages:** The type (α or β) and position (e.g., 1 \rightarrow 4, 1 \rightarrow 6) of the glycosidic bonds determine the 3D conformation and biological properties.
- Molecular Weight:** Very high (from $\sim 10^4$ to $>10^7$ Daltons).
- Supramolecular Organization:** Many structural polysaccharides form **fibers** or **sheets** through extensive intermolecular hydrogen bonding.

B. Physical Properties

- Solubility:** Varies widely.
 - Storage polysaccharides** (glycogen, amylopectin) are **soluble in water** due to their highly branched, open structure.
 - Structural polysaccharides** (cellulose, chitin) are **insoluble** due to tight packing and extensive H-bonding.
- Taste:** **Tasteless** (non-sweet) due to their large size preventing interaction with sweet taste receptors.
- Optical Activity:** Exhibit optical activity but no mutarotation (anomeric carbons are locked in glycosidic bonds).
- Reducing Properties:** Generally **non-reducing**. The number of free anomeric carbons at the ends of long chains is negligible.

C. Chemical Properties

- Hydrolysis:** Can be broken down completely to their constituent monosaccharides by:
 - Acid Hydrolysis:** Boiling with strong mineral acids (e.g., 6M HCl).
 - Enzymatic Hydrolysis:** Specific enzymes (amylases, cellulases, lysozyme).
- Iodine Test:** Forms characteristic-colored complexes with iodine due to the trapping of iodine molecules within the helical structure of the polysaccharide.
 - Amylose:** Deep blue.
 - Amylopectin/Glycogen:** Reddish-purple to brown.
 - Cellulose/Chitin:** No color (no helix formation).

Polysaccharides at a Glance

Polysaccharide	Type	Monomer(s)	Key Linkage(s)	Main Function	Distinctive Feature
Starch	Storage (Plant)	α -D-Glucose	$\alpha(1\rightarrow4)$ linear; $\alpha(1\rightarrow6)$ branches	Energy reserve	Iodine: Blue (amylose), Purple (amylopectin)
Glycogen	Storage (Animal)	α -D-Glucose	$\alpha(1\rightarrow4)$ linear; $\alpha(1\rightarrow6)$	Energy reserve (liver, muscle)	Most branched; soluble; iodine: reddish-brown

			branches (frequent)		
Cellulose	Structural (Plant)	β -D-Glucose	$\beta(1\rightarrow4)$	Mechanical strength (cell walls)	Linear, H-bonded microfibrils; indigestible
Chitin	Structural (Animal/Fungal)	N-Acetyl- β -D-Glucosamine	$\beta(1\rightarrow4)$	Exoskeletons, fungal cell walls	Analogous to cellulose but with NAG; very tough

Lipids

Lipids are a heterogeneous group of organic compounds that are:

- **Insoluble in water** (hydrophobic)
- **Soluble in organic solvents** (ether, chloroform, benzene, acetone)
- Characterized by their **hydrophobic nature** due to predominantly **nonpolar hydrocarbon chains**

General Formula: Predominantly composed of **carbon, hydrogen, and oxygen** (some contain phosphorus, nitrogen, sulfur)

B. Biological Importance and Functions

1. **Energy Storage:** Most efficient energy reserves (9 kcal/g vs. 4 kcal/g for carbs/proteins)
2. **Structural Components:** Major constituents of biological membranes (phospholipids, cholesterol)
3. **Insulation and Protection:** Thermal insulation (subcutaneous fat), organ cushioning
4. **Hormone Precursors:** Steroid hormones, prostaglandins
5. **Vitamin Carriers:** Fat-soluble vitamins (A, D, E, K)
6. **Signaling Molecules:** Eicosanoids, phosphatidylinositol derivatives
7. **Electron Carriers:** Coenzyme Q in ETC
8. **Enzyme Cofactors:** Vitamin K in blood clotting

C. General Characteristics

- **Amphipathic nature:** Most biological lipids have both hydrophobic and hydrophilic regions
- **Diverse structures:** Range from simple hydrocarbon chains to complex ring systems
- **Varied melting points:** Depending on chain length and saturation

PROPERTIES OF LIPIDS

A. Physical Properties

1. **Solubility:**
 - Insoluble in water (hydrophobic)
 - Soluble in nonpolar organic solvents (ether, chloroform, benzene)
 - Amphipathic lipids form micelles/bilayers in aqueous solutions
2. **Melting Points:**
 - **Saturated fats:** Higher melting points (solid at room temp)
 - **Unsaturated fats:** Lower melting points (liquid at room temp)
 - **Chain length effect:** Longer chains = higher melting points
3. **Specific Gravity:** Less than water (0.8-0.9 g/mL) - float on water
4. **Optical Activity:** Some lipids are optically active (glycerol derivatives with asymmetric carbons)
5. **Surface Activity:** Amphipathic lipids reduce surface tension (emulsifiers)

B. Chemical Properties

1. **Hydrolysis:**
 - Acid hydrolysis: Triglycerides \rightarrow Glycerol + Fatty acids



- Alkaline hydrolysis (saponification): Triglycerides → Glycerol + Soap
- Enzymatic hydrolysis (lipases): Digestion of dietary fats
- 2. **Saponification:**
 - Reaction with alkali to form soap
 - **Saponification number:** mg KOH required to saponify 1g fat
 - Indicator of average chain length (higher number = shorter chains)
- 3. **Hydrogenation:**
 - Addition of H₂ to double bonds (unsaturated → saturated)
 - Used in margarine production
 - May produce trans fats (partial hydrogenation)
- 4. **Halogenation:**
 - Addition of halogens (I₂, Br₂) to double bonds
 - **Iodine number:** grams of iodine absorbed by 100g fat
 - Measures degree of unsaturation
- 5. **Rancidity:**
 - **Hydrolytic rancidity:** Lipase action releases fatty acids
 - **Oxidative rancidity:** Auto-oxidation of unsaturated fats
 - Initiation: Free radical formation
 - Propagation: Chain reaction
 - Termination: Antioxidants stop reaction
 - **Prevention:** Antioxidants (vitamin E, BHT, BHA), refrigeration, nitrogen packaging

4. Classification Of Lipids

A. Simple Lipids

Simple lipids are esters of **fatty acids** with various **alcohols** that yield **only two types of products** upon hydrolysis:

1. Fatty acids
2. Alcohols

General Formula: Alcohol + Fatty acid(s) → Ester + H₂O


B. Key Characteristics

1. **Composition:** Contain only **carbon, hydrogen, and oxygen**
2. **No additional groups:** Unlike complex lipids, lack phosphate, carbohydrate, or nitrogenous groups
3. **Hydrophobic:** Primarily nonpolar due to long hydrocarbon chains
4. **Energy-rich:** High caloric value (9 kcal/g)

TRIACYLGLYCEROLS (TRIGLYCERIDES) - FATS AND OILS

A. Chemical Structure

- **Backbone:** Glycerol (a trihydroxy alcohol, C₃H₈O₃)
- **Ester linkage:** Three fatty acids attached via ester bonds
- **General structure:**



B. Types of Triacylglycerols

1. Based on Fatty Acid Composition

a) Simple Triglycerides

- All three fatty acids are identical
- **Examples:**



- **Tristearin:** 3 stearic acids (C18:0)
- **Tripalmitin:** 3 palmitic acids (C16:0)
- **Triolein:** 3 oleic acids (C18:1)

- **Rare in nature:** Most natural fats are mixed

2. Based on Physical State at Room Temperature

a) Fats

- **Solid or semi-solid** at room temperature (20-25°C)
- **High proportion of saturated fatty acids**
- **Animal origin** (mostly): Lard, tallow, butter
- **Plant exceptions:** Cocoa butter, coconut oil (saturated plant fats)

b) Oils

- **Liquid** at room temperature
- **High proportion of unsaturated fatty acids**
- **Plant origin** (mostly): Olive oil, sunflower oil, corn oil

Natural Sources and Distribution

1. Animal Fats

Source	Major Fatty Acids	Characteristics	Uses
Butter	C4:0, C16:0, C18:0, C18:1	80% fat, cholesterol, vitamin A	Cooking, baking
Lard	C16:0, C18:0, C18:1	Pig fat, semi-solid	Pastry, frying
Tallow	C16:0, C18:0, C18:1	Beef/mutton fat, solid	Soap, candles, cooking
Fish oils	C20:5, C22:6 (ω -3)	Liquid, highly unsaturated	Supplements, margarine

2. Plant Oils

Source	Major Fatty Acids	Characteristics	Uses
Olive oil	C18:1 (55-83%), C16:0	Monounsaturated, stable	Cooking, salad dressings
Coconut oil	C12:0 (45%), C14:0	Saturated, medium-chain	Cooking, cosmetics
Palm oil	C16:0 (44%), C18:1	Saturated, semi-solid	Margarine, cooking oil
Sunflower oil	C18:2 (68%), C18:1	Polyunsaturated (ω -6)	Cooking oil, margarine
Flaxseed oil	C18:3 (57%, ω -3)	Highly unsaturated, oxidizes easily	Supplements, paints

Waxes

A. Definition and Structure

- **Esters of long-chain fatty acids with long-chain alcohols**
- **General formula:** R-CO-O-R'
 - R-COOH: Fatty acid (C14-C36)
 - R'-OH: Alcohol (C16-C30)
- **No glycerol backbone**

B. Chemical Composition

1. Fatty Acid Components

- Usually saturated or monounsaturated
- **Even-numbered:** C14-C36
- **Common:** Palmitic (C16:0), stearic (C18:0), oleic (C18:1)
- **Hydroxy fatty acids** in some waxes

2. Alcohol Components

- **Monohydric alcohols:** Straight-chain, saturated
- **Diols:** Some plant waxes
- **Sterols:** Cholesterol in animal waxes
- **Common:** Cetyl alcohol (C16), stearyl alcohol (C18)



M
K

P
R
E
P
A
R
A
T
I
O
N
S

C. Physical Properties

1. **Consistency:** Hard, pliable, non-greasy
2. **Melting point:** 60-100°C (higher than fats)
3. **Solubility:** Insoluble in water, soluble in organic solvents
4. **Hydrophobicity:** Excellent water repellency
5. **Luster:** Naturally shiny appearance

D. Biological Functions in Nature

1. **Waterproofing:** Cuticle of plants, animal fur/feathers
2. **Protection:** Against UV, pathogens, physical damage
3. **Structural:** Beeswax in honeycomb construction
4. **Energy storage:** Some marine organisms
5. **Buoyancy:** Spermaceti in sperm whales

Fatty Acids

Common Name	Systematic Name	Shorthand	Structure Summary
Palmitic	Hexadecanoic	C16:0	Saturated, 16C
Stearic	Octadecanoic	C18:0	Saturated, 18C
Oleic	<i>cis</i> -9-Octadecenoic	C18:1 Δ^9 or ω -9	Monounsaturated, <i>cis</i>
Linoleic	<i>cis,cis</i> -9,12-Octadecadienoic	C18:2 $\Delta^{9,12}$ or ω -6	Dienoic, methylene-interrupted
α -Linolenic	<i>all-cis</i> -9,12,15-Octadecatrienoic	C18:3 $\Delta^{9,12,15}$ or ω -3	Trienoic, ω -3

D. Physical Properties Relationship With Structure

1. Melting Point Trends

- Increasing Chain Length → Higher Melting Point
C12:0 (44°C) < C14:0 (58°C) < C16:0 (63°C) < C18:0 (70°C)
- Increasing Unsaturation → Lower Melting Point
C18:0 (70°C) > C18:1 (13°C) > C18:2 (-5°C) > C18:3 (-11°C)
- Cis vs. Trans Isomers
C18:1 *cis* (oleic): 13°C
C18:1 *trans* (elaidic): 45°C

2. Solubility

- **In water:** Decreases with chain length
 - C2:0 (acetic) - miscible
 - C4:0 (butyric) - soluble
 - C16:0 (palmitic) - insoluble
- **In organic solvents:** Increases with chain length

E. BIOCHEMICAL ROLES AND FUNCTIONS

1. Membrane Fluidity Regulation

- **Saturated fatty acids:** Decrease fluidity, increase packing
- **Unsaturated fatty acids:** Increase fluidity, create kinks
- **Homeoviscous adaptation:** Organisms adjust fatty acid composition to maintain membrane fluidity

2. Energy Storage and Metabolism

- **β -oxidation:** Major pathway for fatty acid catabolism
- **ATP yield:**
 - Palmitic acid (C16:0): 106 ATP
 - Stearic acid (C18:0): 120 ATP
 - Account for activation (-2 ATP) and transport costs

3. Signaling Molecules Precursors

- **Eicosanoids:** Derived from 20-carbon PUFAs
- **Second messengers:** Diacylglycerol (DAG)
- **Protein modification:** Palmitoylation (S-palmitoylation)

4. Gene Expression Regulation

- **PPARs activation:** Polyunsaturated fatty acids activate peroxisome proliferator-activated receptors
- **SREBP regulation:** Sterol regulatory element-binding proteins

B. Complex Lipids


Contain additional groups besides fatty acids and alcohol

1. Phospholipids


Contain phosphoric acid and often nitrogenous base

a) Glycerophospholipids (Phosphoglycerides)

- Glycerol + 2 fatty acids + phosphate + alcohol
- **Major types:**
 - **Phosphatidylcholine (Lecithin):** Choline as alcohol
 - Major membrane component, lung surfactant (dipalmitoyl lecithin)



(1) Phosphatidic acid



(2) Lecithin (phosphatidylcholine)

- **Phosphatidylethanolamine (Cephalin):** Ethanolamine as alcohol
 - Brain and nerve tissue
- **Phosphatidylserine:** Serine as alcohol
 - Important in apoptosis, blood clotting


C. Derived Lipids

Steroids


Basic Structure: Cyclopentanoperhydrophenanthrene nucleus (4 fused rings)

Major Classes:


- 1. Sterols:**
 - **Cholesterol:** Animal membranes, precursor to other steroids
 - **Phytosterols:** Plant sterols (β -sitosterol, campesterol)
 - **Ergosterol:** Fungal sterol, vitamin D precursor
- 2. Steroid Hormones:**
 - **Sex hormones:** Estrogens, androgens, progestins
 - **Adrenocortical hormones:** Glucocorticoids, mineralocorticoids
- 3. Bile Acids:**
 - Cholic acid, chenodeoxycholic acid
 - Emulsify dietary fats
- 4. Vitamin D:** Calcitriol (active form)



Steroid nucleus



Cholesterol



Ergosterol

Terpenes

- Isoprene (C_5H_8) units
- **Monoterpenes (C_{10}):** Essential oils (menthol, limonene)
- **Sesquiterpenes (C_{15}):** Abscisic acid
- **Diterpenes (C_{20}):** Phytol (chlorophyll), gibberellins

- **Triterpenes (C₃₀):** Squalene
- **Tetraterpenes (C₄₀):** Carotenoids
- **Polyterpenes:** Rubber

Lipid Classification Overview

Category	Subcategory	Components	Examples	Functions
SIMPLE	Neutral lipids	Glycerol + 3FA	Triacylglycerols	Energy storage
	Waxes	Long-chain alcohol + FA	Beeswax, lanolin	Protection, waterproofing
	COMPLEX	Phospholipids	Alcohol + FA + P + N-base	Phosphatidylcholine
Glycolipids		Sphingosine + FA + sugar	Cerebrosides	Cell recognition
Lipoproteins		Lipids + proteins	HDL, LDL	Lipid transport
DERIVED	Fatty acids	Hydrocarbon + COOH	Palmitate, oleate	Energy, precursors
	Steroids	Steroid nucleus	Cholesterol, hormones	Membrane, signaling
	Alcohols	Hydroxyl groups	Glycerol, cholesterol	Lipid backbones
MISC	Terpenes	Isoprene units	Carotenoids, rubber	Pigments, hormones
	Eicosanoids	Arachidonate derivatives	Prostaglandins	Local hormones

Proteins

- **Definition:** Proteins are large, complex macromolecules that are fundamental to all life forms. They are polymers, meaning they are built from long, unbranched chains of smaller molecular units.
- **Composition:** The monomeric units (building blocks) of proteins are **amino acids**. A typical protein is composed of **50 to over 2000 amino acids** linked together in a specific, genetically determined sequence.
- **Key Elements:** All proteins contain **Carbon (C), Hydrogen (H), Oxygen (O), and Nitrogen (N)**. Many also contain **Sulfur (S)**, found in certain amino acids like cysteine and methionine.
- **Central Role:** Proteins are the primary functional and structural molecules in cells, executing nearly every cellular task. They are not just a nutritional category but the "workhorses" of the cell.

2. Amino Acids

- **Role:** Amino acids are the **monomers** (singular units) that polymerize to form protein chains (polymers). There are 20 standard amino acids commonly found in proteins, each encoded by the genetic code.
- **Common Structure:** Despite their differences, all 20 standard amino acids share a common fundamental framework.


3. General Structure of Amino Acids

Every amino acid (except proline, which has a slight variation) has the same core structure, centered on the **alpha (α)-carbon**:

1. **Alpha Carbon (C_α):** A central carbon atom.
2. **Amino Group (NH₂ or NH₃⁺):** A basic (proton-accepting) functional group. At physiological pH (~7.4), it is typically protonated, carrying a positive charge (NH₃⁺).
3. **Carboxyl Group (COOH or COO⁻):** An acidic (proton-donating) functional group. At physiological pH, it is typically deprotonated, carrying a negative charge (COO⁻).
4. **Hydrogen Atom (H):** A single hydrogen atom bonded to the alpha carbon.

5. **Side Chain (R-group):** A variable chemical group that is different in each of the 20 amino acids. The identity of the R-group determines the unique chemical properties (size, charge, polarity, hydrophobicity) of each amino acid.


Generalized Structure:



- This structure makes the alpha carbon a **chiral center** (except in glycine, where R=H), giving rise to L- and D-isomers. **All amino acids in proteins are in the L-configuration.**
- At neutral pH, amino acids exist as **zwitterions**—molecules with both a positive and a negative charge, making them overall electrically neutral but polar.

Peptide Bond Formation

- **Process:** Peptide bonds are formed through a **condensation (dehydration) reaction.**
- **Mechanism:**
 1. The **carboxyl group (COOH)** of one amino acid reacts with the **amino group (NH₂)** of another amino acid.
 2. A molecule of water (**H₂O**) is removed.
 3. A covalent bond, the **peptide bond (CO-NH)**, is formed between the carbon of the first amino acid and the nitrogen of the second.




- **Result:** The linked molecules are called **amino acid residues**. Two linked residues form a **dipeptide**, three a **tripeptide**, and many form a **polypeptide** (a single, unbranched protein chain).

Levels of Protein Structure

Protein structure is hierarchical, organized into four levels of increasing complexity.

A. Primary Structure

- **Definition:** The **linear sequence of amino acid residues** in a polypeptide chain, read from N-terminus to C-terminus.
- **Bond Type:** Covalent **peptide bonds** (and sometimes covalent **disulfide bonds** between cysteine residues, which are technically part of the primary sequence but form during folding).
- **Significance:** **The primary structure is the most fundamental level. It dictates all higher levels of folding and, ultimately, the protein's final three-dimensional shape and function.** A change in even a single amino acid (a mutation) can drastically alter protein structure and cause disease (e.g., sickle cell anemia).




B. Secondary Structure

- **Definition:** Local, regular, repeating patterns of folding in segments of the polypeptide backbone, stabilized by **hydrogen bonds** between the backbone carbonyl (C=O) and amide (N-H) groups. *The R-groups/side chains are not involved in these bonds.*
- **Bond Type:** Primarily **hydrogen bonds**.
- **Major Types:**


1. Alpha Helix (α -helix):

- A right-handed coiled structure resembling a spring.
- Stabilized by **intra-chain hydrogen bonds** between the carbonyl oxygen of residue $*n*$ and the amide hydrogen of residue $*n+4*$ (four positions ahead in the sequence).
- Side chains point **outward** from the helical core.



2. Beta Pleated Sheet (β -sheet):

- Formed by stretches of the polypeptide chain (β -strands) lying side-by-side.
- Stabilized by **inter-chain hydrogen bonds** (or intra-chain in a hairpin turn) between neighboring strands.
- Can be **parallel** (adjacent strands run in the same N \rightarrow C direction) or **antiparallel** (adjacent strands run in opposite directions). Antiparallel sheets are generally more stable.
- Side chains alternate pointing **above and below** the plane of the sheet.



3. Beta Turns (β -turns) and Loops:

- Tight, 180° reverse turns that allow the polypeptide chain to change direction.
- Often connect successive strands of β -sheets.
- Loops are less regular, often flexible regions on the protein surface.

C. Tertiary Structure


- **Definition:** The overall, three-dimensional conformation of a single, entire polypeptide chain. It describes how all the secondary structure elements (helices, sheets, loops) fold and pack together in space to form a compact, functional unit.

- **Stabilizing Forces (from strongest to weakest):**
 1. **Covalent Bonds: Disulfide bridges** between the sulfur atoms of two cysteine residues (strong, but not all proteins have them).
 2. **Electrostatic (Ionic) Interactions:** Attraction between positively charged (e.g., Lys, Arg) and negatively charged (e.g., Asp, Glu) side chains. Also called **salt bridges**.
 3. **Hydrogen Bonds:** Between polar side chains and/or with the aqueous environment.
 4. **Van der Waals Forces:** Weak attractions between closely packed non-polar side chains.
 5. **Hydrophobic Effect: The major driving force for folding.** Nonpolar, hydrophobic side chains cluster together in the interior of the protein, away from water, while polar and charged side chains are typically on the surface.

- **Result:** Tertiary folding creates specific regions with unique chemistry called **active sites** (in enzymes) or **binding pockets**. The biologically active conformation is called the **native state**.

D. Quaternary Structure

- **Definition:** The association of two or more individual, folded polypeptide chains (called **subunits** or **protomers**) into a single, functional protein complex.
- **Key Point:** Not all proteins have quaternary structure (e.g., myoglobin is a single polypeptide). Proteins that do are called **multimeric** (dimer = 2 subunits, trimer = 3, tetramer = 4, etc.).
- **Stabilizing Forces:** The same non-covalent interactions as tertiary structure (hydrophobic effect, hydrogen bonds, ionic bonds, van der Waals). Sometimes also covalent **disulfide bonds** between subunits.
- **Significance:** Allows for **cooperativity** and **allosteric regulation**. Subunits often work together. A classic example is **hemoglobin**, a tetramer where binding of oxygen to one subunit increases the oxygen affinity of the other subunits.
- **Analogy:** Tertiary structure is a single engine part; quaternary structure is the assembly of many parts into a complete working engine.



4.12 : Major bonds in protein structure (A) Disulfide bond (B) Hydrogen bonds (C) Hydrophobic bonds (D) Electrostatic bond.

7. Functions of Proteins

Proteins are the most versatile macromolecules, performing an enormous range of functions essential for life. The specific function of a protein is a direct consequence of its unique three-dimensional structure.

A. Structural Proteins

- **Role:** Provide **mechanical support, strength, and integrity** to cells, tissues, and organisms. They create scaffolding and framework.
- **Key Characteristics:** Often form long fibers or sheets; are tough and durable.
- **Examples:**
 - **Collagen:** The most abundant protein in animals. A triple-helical fiber that provides tensile strength to skin, tendons, ligaments, bones, and connective tissues.
 - **Elastin:** Provides elasticity and recoil to tissues like skin, blood vessels, and lungs.



- **Keratin:** A tough, fibrous protein that forms the structural basis of hair, nails, feathers, horns, and the outer layer of skin.
- **Actin & Tubulin:** Globular proteins that polymerize into long filaments (**microfilaments** and **microtubules**). They form the **cytoskeleton**, which maintains cell shape, enables cell movement, and facilitates intracellular transport.

B. Enzymatic Proteins (Enzymes)

- **Role:** Act as **biological catalysts**. They dramatically **speed up (catalyze) specific chemical reactions** without being consumed in the process. Almost every metabolic reaction is facilitated by an enzyme.
- **Key Characteristics:** Highly specific for their **substrate** (the molecule they act upon). They lower the **activation energy** required for a reaction to proceed.
- **Mechanism:** Bind substrates at a specialized region called the **active site**. The "lock-and-key" and "induced fit" models describe this specificity.
- **Examples:**
 - **Amylase:** Catalyzes the breakdown of starch into sugars in digestion.
 - **ATP Synthase:** A complex membrane protein that synthesizes ATP, the cell's energy currency.
 - **DNA Polymerase:** Catalyzes the synthesis of new DNA strands during replication.

C. Transport Proteins

- **Role:** Bind and carry **specific atoms or molecules** from one location to another, either within bodily fluids or across cellular membranes.
- **Types and Examples:**
 1. **Transport in Blood/Bodily Fluids:**
 - **Hemoglobin:** The iron-containing protein in red blood cells that transports **oxygen** from lungs to tissues and carries some **carbon dioxide** back.
 - **Serum Albumin:** Transports fatty acids, hormones, and other substances in the blood.
 2. **Membrane Transport Proteins:** Embedded in cell membranes to move substances across.
 - **Channels & Carriers:** Facilitate the movement of ions (e.g., sodium, potassium) or small molecules (e.g., glucose) across the plasma membrane.
 - **Pumps (e.g., Sodium-Potassium Pump):** Use energy (ATP) to actively transport ions against their concentration gradient.
 3. **Intracellular Transport:** Motor proteins like **myosin, kinesin, and dynein** "walk" along cytoskeletal tracks to transport vesicles and organelles within the cell.

D. Hormonal Proteins (Protein/Peptide Hormones)

- **Role:** Function as **chemical messengers** in endocrine signaling. They are secreted by endocrine glands or tissues into the bloodstream and travel to **target cells** to regulate physiology, growth, and metabolism.
- **Key Characteristics:** They bind to specific **receptors** on the surface of (or inside) target cells, triggering a signal transduction cascade that alters cell activity.
- **Examples:**
 - **Insulin:** Secreted by the pancreas. Signals cells (especially liver, muscle, fat) to take up glucose from the blood, lowering blood sugar levels.
 - **Glucagon:** Also from the pancreas; has the opposite effect of insulin, stimulating the release of glucose into the blood.
 - **Growth Hormone (GH):** Secreted by the pituitary gland, stimulates growth, cell reproduction, and regeneration.
 - **Antidiuretic Hormone (ADH/Vasopressin):** Regulates water balance by signaling the kidneys to retain water.



M
K
P
R
E
P
A
R
A
T
I
O
N
S

Other Crucial Protein Functions (Implied by the outline's focus):

- **Defensive Proteins:** Antibodies (immunoglobulins) that recognize and neutralize pathogens.
- **Storage Proteins:** Store nutrients (e.g., ferritin stores iron; casein in milk stores amino acids).
- **Receptor Proteins:** Built into membranes to receive and transduce chemical signals (e.g., hormone receptors).
- **Motor/Contractile Proteins:** Enable movement (e.g., myosin and actin in muscle contraction).

Enzymes (Biological Catalysts)

1. Definition of Enzymes

Enzymes are highly specialized biological macromolecules (primarily proteins) that act as **catalysts** for biochemical reactions. A catalyst is a substance that **increases the rate of a chemical reaction** without itself being permanently altered or consumed in the process. Enzymes achieve this by **significantly lowering the activation energy** required for the reaction to proceed. They facilitate the conversion of substrates into products, enabling life-sustaining metabolic processes to occur at rates necessary for life, often millions of times faster than uncatalyzed reactions.

2. Nature of Enzymes (Protein Nature)

- **Primary Composition:** The vast majority of enzymes are **globular proteins**. This means they are composed of long chains of amino acids (polypeptides) folded into a specific, complex three-dimensional shape.
- **Key Characteristics Deriving from Protein Nature:**
 - **Specificity:** Their unique 3D structure creates an **active site**—a precise, often cleft or pocket-like region—that is complementary in shape, charge, and hydrophobic/hydrophilic character to a particular substrate or group of closely related substrates. This ensures precise control in metabolism.
 - **Denaturation:** Being proteins, enzymes are susceptible to **denaturation**. This is the irreversible loss of their specific 3D structure due to disruptive conditions (e.g., extreme heat, pH changes, heavy metals). Denaturation destroys the active site, rendering the enzyme permanently inactive.
 - **Cofactors:** Many enzymes require non-protein helper components to function.
 - **Cofactors:** Inorganic ions (e.g., Mg^{2+} , Zn^{2+} , Fe^{2+}/Fe^{3+}) that are tightly bound, often at the active site, and are essential for catalysis.
 - **Coenzymes:** Complex organic or metalloorganic molecules, often derived from vitamins (e.g., NAD^+ , FAD, Coenzyme A). They act as transient carriers of specific functional groups (e.g., electrons, acyl groups).
 - **Holoenzyme vs. Apoenzyme:** The complete, active enzyme complex (protein part + cofactor) is the **holoenzyme**. The protein component alone, which is inactive, is the **apoenzyme**.

Example: DNA Polymerase III (bacterial)

- Apoenzyme: Multiple polypeptide subunits
- Cofactor: Mg^{2+} ions
- Holoenzyme: Complete complex capable of DNA synthesis

Feature	Apoenzyme	Holoenzyme
Composition	Protein part only	Protein + cofactor
Catalytic Activity	Inactive	Active
Formation	Synthesized by ribosomes	Forms when cofactor binds
Specificity	Determines substrate specificity	Determines reaction specificity
Stability	May be unstable without cofactor	Generally more stable
Molecular Weight	Lower	Higher

Example	Pyruvate kinase without K^+/Mg^{2+}	Pyruvate kinase with K^+/Mg^{2+} ions
----------------	---------------------------------------	---

3. Mechanism of Enzyme Action


Enzymes catalyze reactions by providing an alternative, lower-energy pathway. The general sequence is:

- Binding:** The substrate(s) bind reversibly to the enzyme's active site, forming an **enzyme-substrate (ES) complex**. This binding is mediated by multiple weak, non-covalent interactions (hydrogen bonds, ionic bonds, van der Waals forces, hydrophobic interactions).
- Catalysis:** The precise environment of the active site promotes the chemical transformation of the bound substrate(s). Mechanisms include:
 - **Strain and Distortion:** Binding induces strain in the substrate, bending bonds to make them more reactive.
 - **Proximity and Orientation:** Reactants are held in the optimal spatial arrangement for reaction.
 - **Acid-Base Catalysis:** Amino acid side chains donate or accept protons.
 - **Covalent Catalysis:** Temporary formation of a covalent bond between the enzyme and substrate.
 - **Microenvironment:** The active site's unique polarity or hydrophobicity can stabilize transition states.
- Product Formation & Release:** The substrate is converted into product(s), forming an **enzyme-product (EP) complex**. The products, having lower affinity for the active site, are released. The enzyme is then free to bind another substrate molecule. This cycle is described by the **Michaelis-Menten model**.

4. Models of Enzyme-Substrate Specificity

Two primary models explain the specificity of enzyme-substrate binding:

- **Lock and Key Model (Emil Fischer, 1894):**
 - **Concept:** The enzyme's active site is viewed as a rigid, pre-formed **lock** with a fixed geometric shape. The substrate is the **key** that fits perfectly into this lock. Only the correct, complementary substrate can fit and be catalyzed.
 - **Limitation:** It fails to account for the dynamic flexibility of proteins and the observed stabilization of the transition state.
- **Induced Fit Model (Daniel Koshland, 1958):**
 - **Concept:** This is the currently accepted model. The active site is not a rigid structure. Upon initial, weak binding of the substrate, the enzyme undergoes a **conformational change** (a change in shape). This change **induces** a tighter fit around the substrate, like a handshake or a glove molding around a hand. This induced fit:
 1. Optimizes the alignment of catalytic groups.
 2. **Strains** the substrate towards its transition state, lowering the activation energy.
 3. Explains how enzymes can discriminate against similar molecules and perform sequential reactions.




5. Factors Affecting Enzyme Activity

Enzyme reaction rates are highly sensitive to environmental conditions due to their protein nature.


• A. Temperature:

- **General Effect:** Reaction rate increases with temperature (Q_{10} effect) as molecules have more kinetic energy and collide more frequently.
- **Optimum Temperature:** The temperature at which the enzyme operates at maximum velocity. For most human enzymes, this is $\sim 37^{\circ}\text{C}$.
- **Denaturation:** Above the optimum, increased thermal energy disrupts the weak bonds maintaining tertiary structure. The enzyme **denatures**, losing its shape and activity. The rate drops sharply. The curve is a bell shape.



• B. pH:


- **General Effect:** pH affects the ionization state of amino acid side chains (especially in the active site) and the substrate itself.
- **Optimum pH:** The pH at which the enzyme's active site and substrate have the ideal charge configuration for binding and catalysis. (e.g., Pepsin in stomach $\sim \text{pH } 2$, Trypsin in small intestine $\sim \text{pH } 7.5$).
- **Extreme pH:** Very high or low pH can cause **denaturation** by disrupting ionic and hydrogen bonds, or alter charges critical for catalysis, leading to loss of activity. The curve is bell-shaped.




• C. Substrate Concentration ([S]):

- **At Low [S]:** The reaction rate is directly proportional to [S]. Active sites are largely free, and the enzyme is not saturated. Rate increases linearly (first-order kinetics).
- **As [S] Increases:** The rate increases but begins to level off as more active sites become occupied.

- **At High [S]:** The rate plateaus, reaching **Maximum Velocity (V_{max})**. All active sites are continuously occupied (saturated), and the enzyme is working at its maximum capacity. The rate becomes zero-order (independent of [S]).
- **Michaelis Constant (K_m):** This is the substrate concentration at which the reaction rate is **half of V_{max}** . It is a measure of the **enzyme's affinity for its substrate**. A **low K_m** indicates high affinity (the enzyme reaches half V_{max} at a low [S]). A **high K_m** indicates low affinity.



- **D. Enzyme Concentration ([E]):**
 - **General Effect:** Provided there is an **excess of substrate**, the reaction rate is **directly proportional to the enzyme concentration**. Doubling [E] doubles the rate, as there are twice as many catalytic sites available. This relationship is linear until the substrate becomes limiting.



6. Importance of Enzymes

Enzymes are indispensable for all forms of life due to their roles in:

- **Metabolism:** They orchestrate every step of **catabolic pathways** (breaking down molecules to release energy, e.g., glycolysis) and **anabolic pathways** (building complex molecules using energy, e.g., protein synthesis).
- **Homeostasis:** Enzymes help maintain internal stability by regulating biochemical processes in response to cellular signals.
- **Digestion:** Hydrolytic enzymes (e.g., amylase, protease, lipase) break down macromolecules in food into absorbable monomers.
- **Cellular Signaling & Regulation:** Enzymes like kinases and phosphatases control processes by adding/removing phosphate groups, turning proteins on or off.



- **DNA Replication and Repair:** DNA polymerase, ligase, and nucleases are essential for copying and maintaining genetic integrity.
- **Defense:** Enzymes like lysozyme in tears degrade bacterial cell walls.
- **Detoxification:** Liver enzymes (e.g., cytochrome P450) modify toxins and drugs for excretion.
- **Industrial & Medical Applications:** Used in brewing, cheese-making, detergents (proteases, lipases), biofuel production, diagnostic assays, and as therapeutic agents.

M K P R E P A R A T I O N S

7. Enzyme Inhibition (Crucial for Regulation)

Inhibitors are molecules that decrease enzyme activity, providing critical control over metabolic pathways.

- **A. Reversible Inhibition:**
 - **Competitive Inhibition:**
 - **Mechanism:** Inhibitor (I) closely resembles the substrate and **competes for the active site**. It blocks substrate binding.
 - **Effect on Kinetics: Increases apparent K_m** (more substrate is needed to reach half V_{max}), but **V_{max} is unchanged** (with enough substrate, it can outcompete the inhibitor).
 - **Example:** Statin drugs competitively inhibit HMG-CoA reductase, a key enzyme in cholesterol synthesis.
 - **Non-Competitive Inhibition:**
 - **Mechanism:** Inhibitor binds to a site **other than the active site** (an allosteric site), causing a conformational change that deforms the active site and reduces its catalytic efficiency.
 - **Effect on Kinetics: V_{max} is decreased, but K_m remains unchanged** (affinity for substrate is unaltered; binding is unaffected).
 - **Uncompetitive Inhibition:**
 - **Mechanism:** Inhibitor binds only to the **Enzyme-Substrate (ES) complex**, locking it and preventing product release.
 - **Effect on Kinetics: Both V_{max} and apparent K_m are decreased.**
- **B. Irreversible Inhibition:**
 - **Mechanism:** Inhibitor forms strong **covalent bonds** with amino acid residues in the active site, permanently inactivating the enzyme. The enzyme must be degraded and replaced.
 - **Example:** Penicillin irreversibly inhibits transpeptidase, an enzyme essential for bacterial cell wall synthesis.

8. Regulation of Enzyme Activity

Cells precisely control enzyme activity to match metabolic demands.

- **Allosteric Regulation:** Effector molecules (activators or inhibitors) bind to **allosteric sites**, causing conformational changes that either enhance or reduce activity. Often seen in multi-subunit enzymes at key pathway junctions (e.g., feedback inhibition).
- **Covalent Modification:** Reversible addition/removal of chemical groups (e.g., phosphate, acetyl) by other enzymes. **Phosphorylation** (by kinases) is a major regulatory switch.
- **Zymogen (Proenzyme) Activation:** Enzymes are synthesized as inactive precursors to prevent damage inside producing cells. They are activated by proteolytic cleavage when needed (e.g., pepsinogen → pepsin in stomach; blood clotting cascade enzymes).
- **Compartmentalization:** Segregating enzymes and substrates into specific organelles (e.g., digestive enzymes in lysosomes) controls when/where reactions occur.
- **Genetic/Transcriptional Control:** Regulating the *amount* of enzyme produced by turning genes on or off (a slower, long-term adaptation).

Classification of Enzymes

Enzymes are systematically classified by the **Enzyme Commission (EC)**, an international organization, based on the **type of chemical reaction they catalyze**. Each enzyme is assigned a unique four-part **EC number** (e.g., EC 1.1.1.1 for Alcohol Dehydrogenase).

The six main classes are:

EC 1: Oxidoreductases

- **Reaction Catalyzed:** Catalyze **oxidation-reduction (redox) reactions**. Involve the transfer of electrons (often as hydride ions (H^-) or hydrogen atoms) from one molecule (the **reductant**, or electron donor) to another (the **oxidant**, or electron acceptor).
- **Typical Naming:** *Dehydrogenases* (remove hydrogens), *Reductases*, *Oxidases* (use O_2 as electron acceptor), *Peroxidases* (use H_2O_2), *Oxygenases* (incorporate oxygen into a substrate).
- **General Equation:** $AH_2 + B \rightarrow A + BH_2$ (where AH_2 is the electron donor).
- **Key Cofactors:** NAD^+ , $NADP^+$, FAD, FMN, metal ions (Fe, Cu).
- **Examples:**
 - **Cytochrome c Oxidase (EC 1.9.3.1):** Final electron acceptor in the electron transport chain.
 - **Alcohol Dehydrogenase (EC 1.1.1.1):** Oxidizes ethanol to acetaldehyde.
 - **Catalase (EC 1.11.1.6):** Breaks down hydrogen peroxide: $2H_2O_2 \rightarrow O_2 + 2H_2O$

EC 2: Transferases

- **Reaction Catalyzed:** Transfer a **specific functional group** (e.g., phosphate, methyl, amino, glycosyl) from a **donor molecule** to an **acceptor molecule**.
- **Typical Naming:** *Transaminase* (transfer amino group), *Kinase* (transfers phosphate from ATP), *Methyltransferase*.
- **General Equation:** $AX + B \rightarrow A + BX$
- **Examples:**
 - **Hexokinase (EC 2.7.1.1):** Transfers a phosphate from ATP to glucose.
 - **Alanine Transaminase (ALT, EC 2.6.1.2):** Transfers an amino group between alanine and α -ketoglutarate. A key liver enzyme.

EC 3: Hydrolases

- **Reaction Catalyzed:** Catalyze **hydrolysis reactions** – breaking bonds using water. They cleave substrates by adding H^+ and OH^- from water across the bond.
- **Typical Naming:** Often ends in *-ase* preceded by the substrate (e.g., *Protease*, *Lipase*, *Nuclease*, *Phosphatase*).
- **General Equation:** $A-B + H_2O \rightarrow A-H + B-OH$
- **Examples:**
 - **ATP Synthase (EC 3.6.3.14):** Actually catalyzes ATP hydrolysis in reverse to synthesize ATP. (Note: It is named for its reverse, hydrolytic reaction in classification).
 - **Trypsin (EC 3.4.21.4):** A protease that hydrolyzes peptide bonds.
 - **Acetylcholinesterase (EC 3.1.1.7):** Hydrolyzes the neurotransmitter acetylcholine.

EC 4: Lyases

- **Reaction Catalyzed:** Cleave **C-C, C-O, C-N, and other bonds by means other than hydrolysis or oxidation**. They often form a **new double bond** or a **new ring structure**, or they add groups *to* double bonds.
- **Key Feature:** They work in **reverse** as well, catalyzing the addition of groups to double bonds.
- **Typical Naming:** *Decarboxylase* (removes CO_2), *Dehydratase* (removes H_2O), *Aldolase* (reversibly cleaves aldols), *Synthase* (often used for lyases that form bonds).
- **General Equation:** $A-B \rightarrow A + B$ (non-hydrolytic, non-oxidative elimination).
- **Examples:**
 - **Pyruvate Decarboxylase (EC 4.1.1.1):** Removes CO_2 from pyruvate to form acetaldehyde.

- **Fumarase (EC 4.2.1.2):** Hydration of fumarate to malate (adds H₂O across a double bond).

EC 5: Isomerases

- **Reaction Catalyzed:** Catalyze **intramolecular rearrangements**; they change the structure of a molecule by moving functional groups *within* the molecule to form an **isomer**.
- **Types:** *Racemases* or *Epimerases* (change optical isomer), *Cis-trans Isomerases*, *Intramolecular Transferases* (mutases).
- **General Equation:** A → A' (isomer)
- **Examples:**
 - **Triosephosphate Isomerase (TIM, EC 5.3.1.1):** Rapidly converts dihydroxyacetone phosphate (a ketose) to glyceraldehyde 3-phosphate (an aldose). Crucial in glycolysis.
 - **Phosphoglucumutase (EC 5.4.2.2):** Converts glucose 1-phosphate to glucose 6-phosphate.

EC 6: Ligases (Synthetases)

- **Reaction Catalyzed:** Join two molecules together with **covalent bonds**, coupled with the **hydrolysis of a high-energy triphosphate** (usually ATP).
- **Crucial Point:** **Require energy input** (from nucleoside triphosphates like ATP).
- **Typical Naming:** *Synthetase* (this term is reserved for ligases; note: *synthase* is used for lyases).
- **General Equation:** A + B + ATP → A-B + ADP + Pi (or AMP + PPi)
- **Examples:**
 - **DNA Ligase (EC 6.5.1.1):** Seals nicks in DNA backbone during replication/repair.
 - **Acetyl-CoA Synthetase (EC 6.2.1.1):** Activates acetate: Acetate + CoA + ATP → Acetyl-CoA + AMP + PPi
 - **Aminoacyl-tRNA Synthetases (e.g., EC 6.1.1.-):** Attach amino acids to their corresponding tRNAs (essential for translation).

Summary Mnemonic: "Over The Hills, Lions Isolate Lunch"

- Oxidoreductases
- Transferases
- Hydrolases
- Lyases
- Isomerases
- Ligases

Each main class is further divided into subclasses and sub-subclasses based on more specific details of the reaction (e.g., the type of group transferred or the exact bond cleaved), leading to the full four-digit EC number.

EC Class	Class Name	Type of Reaction Catalyzed	Example (EC Number)
EC 1	Oxidoreductases	Oxidation-Reduction (electron transfer).	Alcohol Dehydrogenase (EC 1.1.1.1) - Oxidizes ethanol to acetaldehyde.
EC 2	Transferases	Group Transfer (moving functional groups).	Hexokinase (EC 2.7.1.1) - Transfers a phosphate from ATP to glucose.
EC 3	Hydrolases	Hydrolysis (bond cleavage with water).	Trypsin (EC 3.4.21.4) - Hydrolyzes peptide bonds in proteins.
EC 4	Lyases	Non-Hydrolytic Addition/Elimination (forming or breaking double bonds).	Pyruvate Decarboxylase (EC 4.1.1.1) - Removes CO ₂ from pyruvate.



M
K
P
R
E
P
A
R
A
T
I
O
N
S

EC 5	Isomerases	Isomerization (intramolecular or rearrangement).	Triosephosphate Isomerase (TIM) (EC 5.3.1.1) - Converts dihydroxyacetone phosphate to glyceraldehyde 3-phosphate.
EC 6	Ligases	Bond Formation Coupled to ATP Hydrolysis (joining molecules).	DNA Ligase (EC 6.5.1.1) - Joins DNA strands by forming a phosphodiester bond.

Nucleic Acids

Nucleic acids are large, complex macromolecules that are essential for all known forms of life. They are biopolymers composed of repeating monomeric units called **nucleotides**. Their primary function is to store, transmit, and express genetic information.

Types of Nucleic Acids

There are two main types of nucleic acids, which differ in their structure, sugar component, and specific roles within the cell:

Feature	DNA (Deoxyribonucleic Acid)	RNA (Ribonucleic Acid)
Full Name	Deoxyribonucleic Acid	Ribonucleic Acid
Sugar	Deoxyribose	Ribose
Bases	Adenine (A), Guanine (G), Cytosine (C), Thymine (T)	Adenine (A), Guanine (G), Cytosine (C), Uracil (U)
Structure	Typically double-stranded (double helix)	Typically single-stranded (but can fold)
Stability	More chemically stable	Less chemically stable
Primary Role	Long-term storage of genetic blueprints; hereditary material	Involved in the process of decoding DNA into proteins; various functional roles

- **DNA** serves as the permanent, archival repository of genetic instructions for an organism's development, functioning, and reproduction. It is located in the cell nucleus (and mitochondria/chloroplasts).
- **RNA** acts as a messenger and helper molecule, carrying out the instructions encoded in DNA. It is involved in protein synthesis (gene expression) and has other regulatory and catalytic functions.


Structure of Nucleotides

The nucleotide is the fundamental building block of all nucleic acids. Each nucleotide consists of three components:


1. **A Pentose Sugar:** This is a 5-carbon sugar.
 - In **DNA**, the sugar is **deoxyribose** (lacks an oxygen atom on the 2' carbon).
 - In **RNA**, the sugar is **ribose** (has a hydroxyl group on the 2' carbon).
2. **A Phosphate Group:** A phosphorus atom bonded to four oxygen atoms. This group is attached to the 5' carbon of the sugar. The phosphate group gives nucleic acids their acidic character and allows nucleotides to link together via **phosphodiester bonds**.

3. A **Nitrogenous Base**: A nitrogen-containing molecule attached to the 1' carbon of the sugar. There are two categories of bases:


- **Purines** (double-ring structures): **Adenine (A)** and **Guanine (G)**.
- **Pyrimidines** (single-ring structures): **Cytosine (C)**, **Thymine (T)** (found only in DNA), and **Uracil (U)** (found only in RNA).




Adenine (A)
(6-aminopurine)




Guanine (G)
(2-amino 6-oxypurine)





Cytosine (C)
(2-oxo 4-aminopyrimidine)



Thymine (T)
(2,4-dioxy-5 methylpyrimidine)



Uracil (U)
(2,4-dioxypyrimidine)



Structure of DNA

DNA's structure exists on multiple levels, culminating in its iconic three-dimensional form.

- **Primary Structure:** This is the linear sequence of nucleotides linked by phosphodiester bonds, forming a single polynucleotide strand. The sequence of bases (A, T, C, G) in this strand is the genetic code.
- **Secondary Structure: The Double Helix**
The famous **double helix** model was elucidated by James Watson and Francis Crick in 1953, based on the X-ray crystallography work of Rosalind Franklin. This model describes DNA's stable, three-dimensional shape and has several key features:
 1. **Two Strands:** DNA consists of **two** polynucleotide strands.
 2. **Anti-parallel Orientation:** The two strands run in opposite directions. One strand runs in the **5' → 3'** direction (from its phosphate end to its sugar hydroxyl end), while the complementary strand runs **3' → 5'**.
 3. **Sugar-Phosphate Backbone:** The backbones of the two strands coil around the outside of the molecule.
 4. **Bases Inward:** The nitrogenous bases point inward, toward the central axis of the helix, like the rungs of a twisted ladder.
 5. **Helical Twist:** The two strands twist around each other to form a right-handed spiral, completing one turn approximately every 10 base pairs.
- **Base Pairing (Complementary Base Pairing)**
The two strands of the helix are held together by specific hydrogen bonds between the inward-facing bases. This follows a strict rule known as **Chargaff's rules**:
 - **Adenine (A)** always pairs with **Thymine (T)** via **two** hydrogen bonds.
 - **Guanine (G)** always pairs with **Cytosine (C)** via **three** hydrogen bonds.

This specific **A-T / G-C pairing** is called **complementary base pairing**. It means the sequence of one strand automatically dictates the sequence of the other. For example, if one strand reads 5'-ATGCC-3', the complementary strand must read 3'-TACGG-5'.

Consequences of Base Pairing:

- **Stability:** The three hydrogen bonds in G-C pairs make regions of DNA with high G-C content slightly more stable than regions with high A-T content.
- **Replication:** Complementarity allows DNA to be replicated accurately; each strand serves as a template for the synthesis of a new complementary strand.
- **Information Storage:** Genetic information is stored in the *sequence* of bases, not in their chemical identity. Any sequence is possible within the pairing rules.

The DNA Double Helix (Watson-Crick Model, 1953)

Informed by Rosalind Franklin's X-ray diffraction data and Chargaff's rules.

Key Features

1. **Double Helix:** Two polynucleotide strands wind around a common axis in a right-handed spiral.
2. **Antiparallel:** One strand runs 5' → 3', the other runs 3' → 5'.
3. **Complementary Base Pairing:** Hydrogen bonds form between bases on opposite strands.
 - **Adenine (A) pairs with Thymine (T) via 2 hydrogen bonds.**
 - **Guanine (G) pairs with Cytosine (C) via 3 hydrogen bonds.**
 - This explains **Chargaff's Rules:** A=T and G=C.
4. **Base Stacking:** The flat, planar bases stack on top of each other perpendicular to the helix axis. Hydrophobic interactions and van der Waals forces between stacked bases provide significant stability to the helix.

Helical Geometry


- **Diameter:** ~20 Å (2 nanometers).
- **Helical Pitch:** One full turn is ~34 Å (3.4 nm) long and contains **10 base pairs**.
- **Distance between bases:** ~3.4 Å (0.34 nm) along the axis.
- **Major and Minor Grooves:** The twisting creates two grooves of unequal size.
 - **Major Groove:** Wider, exposes edges of base pairs for protein recognition (e.g., transcription factors).
 - **Minor Groove:** Narrower, also used for specific protein binding.

Forms of DNA

- **B-DNA:** The most common, biologically prevalent form under physiological conditions. Features described above (10 bp/turn, right-handed).
- **A-DNA:** Shorter, wider, right-handed helix formed under dehydrating conditions (11 bp/turn). Found in some protein-DNA complexes.
- **Z-DNA:** A left-handed helix with a zigzag sugar-phosphate backbone. Forms in sequences with alternating purines/pyrimidines (e.g., GCGCGC) under high salt or negative supercoiling. May play a role in gene regulation.

Tertiary Structure: DNA Supercoiling

Long DNA molecules must be packaged to fit inside cells. This is achieved by **supercoiling**.





M
K
P
R
E
P
A
R
A
T
I
O
N
S

- **Relaxed DNA:** B-DNA has one twist per 10.4 base pairs.
- **Supercoiled DNA:** The DNA double helix is further twisted upon itself, like twisting a rubber band.
 - **Positive Supercoiling:** Overwinding of the helix.
 - **Negative Supercoiling:** Underwinding of the helix. **Most prevalent in nature**, facilitates processes requiring strand separation (e.g., replication, transcription).
- **Topoisomerases:** Enzymes that cut and reseal DNA strands to introduce or relieve supercoiling (e.g., DNA gyrase).

Quaternary Structure: DNA Packaging in Chromatin (Eukaryotes)

Nucleosomes: The Fundamental Unit

DNA wraps around protein complexes called **histones**.

- **Core Histone Octamer:** Two copies each of H2A, H2B, H3, and H4.
- ~147 base pairs of DNA wrap **1.65 times** around the octamer.
- **Linker DNA:** Connects nucleosomes (~20-60 bp), associated with **Histone H1**, which helps stabilize the structure.
- This forms the "**beads on a string**" structure (10 nm fiber).

Higher-Order Packaging

1. **30 nm Fiber:** Nucleosomes coil into a solenoid structure, stabilized by H1 histones.
2. **Chromatin Loops:** The 30 nm fiber forms loops (40,000-100,000 bp each) anchored to a protein **scaffold** (using proteins like cohesin, condensin).
3. **Metaphase Chromosome:** Further coiling and folding during cell division creates the highly condensed, visible chromosomes.

Chromatin States

- **Euchromatin:** Less condensed, transcriptionally **active**, gene-rich.
- **Heterochromatin:** Highly condensed, transcriptionally **inactive**.
 - **Constitutive:** Always condensed (e.g., centromeres, telomeres).
 - **Facultative:** Can condense or decondense (e.g., X-chromosome inactivation).

Prokaryotic DNA Packaging

- DNA is circular and supercoiled.
- Organized into **nucleoid-associated proteins (NAPs)** (e.g., HU, H-NS) that bend and bridge DNA, forming loops but no true nucleosomes.

Specialized DNA Structures

- **Palindromic Sequences:** Can form **hairpins** or **cruciforms** (in single-stranded or negatively supercoiled DNA).
- **Triplex DNA (H-DNA):** Three-stranded structure in polypurine/polypyrimidine tracts.
- **G-Quadruplex:** Four-stranded structure formed in guanine-rich sequences (e.g., telomeres).

Functional Implications of Structure

- **Complementarity & Replication:** Strands serve as templates for accurate copying.
- **Genetic Code:** The sequence of bases encodes information for protein synthesis.
- **Protein Recognition:** Specific base sequences and groove geometries allow proteins to bind and regulate genes.
- **Stability:** Hydrogen bonds provide specificity; base stacking and hydrophobic core provide overall stability. The absence of a 2' OH makes DNA more chemically stable than RNA for long-term storage.



Types of DNA: A Comparative Table

Category of Type	DNA Type	Key Characteristics	Primary Role/Notes
Based on Structural Conformation	B-DNA	The classic, right-handed double helix. ~10 base pairs per turn. Hydrated, biologically most common form.	Predominant form under physiological conditions. Standard model for genetic storage and function.
	A-DNA	Shorter, wider right-handed helix. ~11 base pairs per turn. Forms under dehydrating conditions.	Found in DNA-RNA hybrids and some protein-DNA complexes. More compact than B-DNA.
	Z-DNA	Left-handed double helix with a zigzag backbone. ~12 base pairs per turn. Forms in sequences with alternating purines/pyrimidines (e.g., GCGCGC).	Associated with gene regulation, especially near promoter regions. Induced by negative supercoiling.
Based on Functional Role	Coding DNA	Sequences of genes that are transcribed into mRNA and ultimately translated into amino acid sequences (proteins).	Makes up a small percentage of the genome (~1-2% in humans). Directly determines an organism's traits via proteins.
	Non-Coding DNA	Sequences that are not translated into proteins. Includes a variety of functional and non-functional elements.	Comprises the vast majority of eukaryotic genomes. Includes regulatory sequences, introns, and repetitive DNA.
	Regulatory DNA	A subset of non-coding DNA that controls gene expression (e.g., promoters, enhancers, silencers).	Acts as binding sites for transcription factors and other proteins to turn genes on/off.
Based on Sequence & Location in Genome	Unique (Single-Copy) DNA	Sequences that appear only once or a few times in the genome. Includes most protein-coding genes.	Forms the basis of genetic individuality and codes for most functional products.
	Repetitive DNA	Sequences repeated hundreds to millions of times throughout	Involved in chromosome structure, evolution, and some diseases.

		the genome. Two main classes:	
	• Tandem Repeats	Short sequences repeated head-to-tail in clusters (e.g., satellite, minisatellite, microsatellite DNA).	Satellite DNA: Found at centromeres & telomeres (structural). Microsatellites: Used in DNA fingerprinting.
	• Interspersed Repeats	Repeated sequences scattered across the genome, derived from transposable elements (e.g., SINEs like <i>Alu</i> , LINEs).	Makes up a large fraction of mammalian genomes (~45% in humans). Can influence gene expression and genome evolution.
Based on Cellular Location	Nuclear DNA (nDNA)	DNA enclosed within the cell nucleus. Organized into linear chromosomes and complexed with histones into chromatin.	Contains the vast majority of an organism's genetic material. Inherited from both parents (biparental inheritance).
	Mitochondrial DNA (mtDNA)	Small, circular DNA molecule found in mitochondria. Lacks histones and introns. Maternally inherited in most species.	Codes for tRNAs, rRNAs, and proteins essential for oxidative phosphorylation. Has a higher mutation rate than nDNA.
	Chloroplast DNA (cpDNA)	Circular DNA found in chloroplasts of plant cells and other photosynthetic eukaryotes.	Codes for proteins and RNAs involved in photosynthesis. Also maternally inherited in many plants.
Based on Physical State	Linear DNA	DNA molecules with two distinct ends. Characteristic of eukaryotic nuclear chromosomes.	Ends are protected by telomeres , which prevent degradation and fusion.
	Circular DNA	DNA molecules that form a closed loop with no free ends.	Found in prokaryotic chromosomes, plasmids, mitochondria, and chloroplasts. Supercoiling is common.
	Single-Stranded DNA (ssDNA)	DNA consisting of a single polynucleotide chain. Not a stable storage form.	Temporary state during DNA replication (on the lagging strand template) and recombination . Also the genetic material of some viruses (e.g., parvoviruses).
Special / Alternative Structures	Triplex DNA (H-DNA)	Triple-stranded structure that can form in sequences	May play a role in gene regulation (transcription repression) and recombination.

		with mirror symmetry (polypurine-polypyrimidine tracts).	
	G-Quadruplex DNA	Four-stranded structure formed in guanine-rich sequences (e.g., telomeres, gene promoters). Stabilized by Hoogsteen hydrogen bonding.	Implicated in the regulation of replication, transcription, and telomere maintenance. A target for potential cancer therapeutics.
	Catenated DNA	Two or more circular DNA molecules that are interlinked like rings in a chain.	A topological state that must be resolved by topoisomerases after DNA replication in circular genomes.
Extrachromosomal DNA	Plasmid DNA	Small, circular, extrachromosomal DNA molecules found in bacteria, archaea, and some eukaryotes (e.g., yeast).	Often carry genes conferring selective advantages (e.g., antibiotic resistance). Key tools in genetic engineering and biotechnology.
	Viral DNA	The genetic material of DNA viruses. Can be single or double-stranded, linear or circular.	Exists independently of the host chromosome during viral replication. Can integrate into the host genome (e.g., retroviruses via a DNA intermediate).

Ribonucleic Acid (RNA)

I. Fundamental Structure of RNA

A. Chemical Composition

RNA is a **ribonucleic acid** polymer with four key differences from DNA:

Sugar: Ribose (has a 2'-hydroxyl group, making it more chemically reactive than deoxyribose).

- Bases: Adenine (A), Guanine (G), Cytosine (C), and Uracil (U).** Uracil replaces thymine and pairs with adenine.
- Strandedness:** Primarily **single-stranded**, but folds into complex 3D shapes via intramolecular base pairing.
- Overall Stability:** Less chemically stable than DNA due to the 2'-OH group, which makes it more susceptible to hydrolysis. This suits its role as a transient information carrier.

B. Levels of Structural Organization

Level	Description	Significance
Primary Structure	The linear sequence of nucleotides linked by 3'→5' phosphodiester bonds . The sequence is written 5'→3'.	Contains the information (genetic code in mRNA, anticodon in tRNA, structural motifs in rRNA).
Secondary Structure	Local folding into double-stranded regions via intra-strand base pairing (A=U, G≡C). Common	Provides stability, creates recognition sites for proteins/other RNAs, and is the foundation for tertiary folding.

	motifs: Stem-loops (hairpins), internal loops, bulges, pseudoknots.	
Tertiary Structure	The overall three-dimensional shape formed by long-range interactions between secondary structure elements. Stabilized by: <ul style="list-style-type: none"> • Non-canonical base pairs (e.g., G=U wobble) • Stacking interactions between bases • Ionic interactions with Mg²⁺ and other metals • Interactions between the sugar-phosphate backbone 	Creates unique surfaces and clefts that determine the RNA's specific biological function, including catalytic sites in ribozymes.

II. Major Types of RNA and Their Detailed Structures

A. Messenger RNA (mRNA)

Function: Carries the genetic blueprint from DNA in the nucleus to the ribosome in the cytoplasm for protein synthesis.

Detailed Structure:

1. **5' Cap:** A modified guanine nucleotide (7-methylguanosine) added post-transcriptionally. It is linked via a **5'-5' triphosphate bridge**.
 - **Functions:** Protects from degradation, aids nuclear export, and is recognized by the ribosome for translation initiation.
2. **5' Untranslated Region (5' UTR):** Non-coding sequence that contains the **ribosome binding site** (Shine-Dalgarno sequence in prokaryotes; Kozak sequence in eukaryotes).
3. **Coding Sequence:** Composed of a series of **codons** (three-nucleotide units), each specifying an amino acid. Starts with an **AUG start codon** and ends with a **stop codon** (UAA, UAG, UGA).
4. **3' Untranslated Region (3' UTR):** Contains regulatory sequences that influence mRNA stability, localization, and translation efficiency.
5. **Poly-A Tail:** A stretch of 50-250 adenine nucleotides added to the 3' end.
 - **Functions:** Increases stability, aids in export, and is involved in translation termination and recycling.

Key Feature: Typically linear with minimal secondary structure in the coding region to allow smooth ribosome passage, though UTRs often contain regulatory structures.

B. Transfer RNA (tRNA)

Function: The "adapter" molecule that reads the mRNA codon and delivers the corresponding amino acid to the growing polypeptide chain.

Detailed Structure (The Cloverleaf Model - Secondary Structure):

1. **Acceptor Stem (3' CCA End):**
 - Formed by base pairing between the 5' and 3' ends.
 - The 3' end always terminates in the sequence **CCA**. The amino acid is covalently attached (esterified) to the 3'-OH of the terminal adenosine.
2. **D Arm:** Contains **dihydrouridine** (a modified base). Important for tRNA recognition by the correct **aminoacyl-tRNA synthetase**.
3. **Anticodon Arm:** Contains the **anticodon triplet** (e.g., 3'-UAC-5') that base-pairs with the complementary mRNA codon (e.g., 5'-AUG-3') during translation.
4. **Variable Loop:** Size varies among tRNAs; function not always clear.

- T ψ C Arm:** Contains the sequence T ψ C (ribothymidine, pseudouridine, cytidine). Critical for binding to the ribosome.

Tertiary Structure (L-shaped 3D Fold):


- The two "halves" of the cloverleaf fold at right angles.
- The **D arm and T ψ C arm** form one stem, stabilizing the core.
- The **acceptor stem and anticodon arm** form the other stem, positioned $\sim 70\text{\AA}$ apart.
- This L-shape places the **amino acid attachment site** (3' CCA) at one end and the **anticodon** at the other, perfectly adapted for its role on the ribosome.

C. Ribosomal RNA (rRNA)

Function: Forms the **core structural and catalytic framework** of the ribosome, the macromolecular machine that synthesizes proteins.

Ribosomal Subunit	rRNA Components (Prokaryotes - E. coli)	rRNA Components (Eukaryotes - Humans)	Key Structural & Functional Roles
Small Subunit	16S rRNA (1542 nucleotides)	18S rRNA (~ 1900 nt)	<ul style="list-style-type: none"> Binds mRNA Decodes codon-anticodon pairing Contains the decoding center.
Large Subunit	23S rRNA (2904 nt) 5S rRNA (120 nt)	28S rRNA (~ 5000 nt) 5.8S rRNA (156 nt) 5S rRNA (121 nt)	<ul style="list-style-type: none"> Catalyzes peptide bond formation (the peptidyl transferase center is a ribozyme composed of 23S/28S rRNA). Contains the exit tunnel for the nascent polypeptide.

- Extensive Secondary/Tertiary Structure:** rRNA is highly folded with numerous stem-loops, creating precise 3D scaffolds.
- Ribozymatic Activity:** The **peptidyl transferase** reaction (forming the peptide bond) is catalyzed by the rRNA itself, not the ribosomal proteins.
- Ribosome Assembly:** rRNAs act as a "backbone" around which ribosomal proteins assemble. The three-dimensional structure creates distinct functional sites: **A site** (aminoacyl-tRNA binding), **P site** (peptidyl-tRNA binding), **E site** (exit).



Other Crucial Functional RNAs

Type	Full Name	Size & Structure	Primary Function
snRNA	Small Nuclear RNA	100-200 nt; complex with proteins to form snRNPs ("snurps").	Key components of the spliceosome ; catalyze the removal of introns from pre-mRNA in eukaryotes.
snoRNA	Small Nucleolar RNA	60-300 nt; found in the nucleolus.	Guide the chemical modification (e.g., methylation, pseudouridylation) of other RNAs, primarily rRNAs and tRNAs.
miRNA	MicroRNA	~22 nt; form imperfect duplexes with target mRNA.	Gene regulation. Bind to complementary sequences in the 3'UTR of target mRNAs, leading to translational repression or mRNA degradation.
siRNA	Small Interfering RNA	~21-23 nt; form perfect duplexes.	Gene silencing. Derived from long double-stranded RNA, they guide the RISC complex to cleave complementary viral or transposon mRNA. Also used in RNAi technology.
lncRNA	Long Non-coding RNA	>200 nt; diverse structures.	Multifunctional regulators. Involved in X-chromosome inactivation, genomic imprinting, chromatin remodeling, and transcriptional regulation.
Catalytic RNAs (Ribozymes)	-	Varied (e.g., Group I/II introns, RNase P, Hammerhead ribozyme).	RNA molecules with enzymatic activity . They catalyze site-specific RNA cleavage, splicing, or peptide bond formation (rRNA).

Historical Discoveries of Major Biological Molecules

Molecule / Concept	Key Discoverer(s) & Contributors	Approx. Date	Significance of Discovery
DNA (as genetic material)	Friedrich Miescher	1869	Isolated a phosphorus-rich, acidic substance from white blood cell nuclei, which he named " nuclein " (later nucleic acid). Did not know its function.
	Oswald Avery, Colin MacLeod, Maclyn McCarty	1944	Avery-MacLeod-McCarty experiment provided the first definitive evidence that DNA (not protein) is the substance that carries genetic information, transforming the "transforming principle" in <i>Pneumococcus</i> .
	Alfred Hershey & Martha Chase	1952	Hershey-Chase experiment used bacteriophages to confirm that DNA, not protein, is the genetic material.
Structure of DNA	Rosalind Franklin & Maurice Wilkins	1950-1953	Franklin's Photo 51 , obtained via X-ray crystallography, revealed the helical,

			double-stranded pattern with bases stacked interiorly. Crucial, unacknowledged data.
	James Watson & Francis Crick	1953	Proposed the double helix model in their paper "Molecular Structure of Nucleic Acids," correctly explaining base pairing (A-T, G-C) and the mechanism for replication.
RNA	Multiple researchers (following Miescher)	Late 1800s/Early 1900s	Initially distinguished from DNA by its sugar (ribose) and presence in cytoplasm. Its diverse roles were elucidated much later in the 20th century.
Proteins	Jöns Jacob Berzelius	1838	Coined the term " protein " (from Greek <i>proteios</i> , "primary"). Recognized them as a distinct class of biological molecules essential to life.
	Frederick Sanger	1951-1955	Sequenced the amino acids of insulin , proving proteins have a specific, defined sequence. This is the first protein ever sequenced.
Enzymes	Anselme Payen & Jean-François Persoz	1833	Isolated diastase (amylase) from malt, the first enzyme preparation and discovery of a biological catalyst.
	Eduard Buchner	1897	Demonstrated that cell-free yeast extracts could ferment sugar, proving enzymatic activity was separate from living cells. Nobel Prize (1907).
	James B. Sumner	1926	First to crystallize an enzyme (urease) and prove its protein nature, settling a long debate. Nobel Prize (1946).
Carbohydrates	Ancient knowledge	-	Sugars and starches known since antiquity.
	Emil Fischer	1880s-1890s	Elucidated the structures of many simple sugars (glucose, fructose), proposed the " lock and key " model for enzyme action, and pioneered carbohydrate chemistry. Nobel Prize (1902).
Lipids & Membranes	William Prout	Early 1800s	Classified food components into saccharinous (carbs), albuminous (proteins), and oleaginous (fats) .
	E. Gorter & F. Grendel	1925	Extracted lipids from red blood cell membranes and concluded membranes are a lipid bilayer , the foundation of all modern membrane models.
ATP (Energy Currency)	Karl Lohmann	1929	Discovered Adenosine triphosphate (ATP) in muscle extracts.



	Fritz Albert Lipmann	1941	Formulated the " high-energy phosphate bond " concept, establishing ATP as the central carrier of chemical energy in cells. Nobel Prize (1953).
--	-----------------------------	------	--

M
K
P
R
E
P
A
R
A
T
I
O
N
S

Biological Molecules & Enzymes: One Liners

- **Biochemistry** is the branch of science that explores the chemical processes and substances occurring within living organisms.
- It serves as a bridge between biology and chemistry, explaining life at a molecular level.
- All biochemical processes occur within or are mediated by cells (**Cellular Basis**).
- The structural hierarchy in living systems is: Atoms → Small Molecules → Monomers → Polymers → Supramolecular Complexes → Organelles → Cells.
- The three-dimensional shape (**conformation**) of a biomolecule is directly linked to its biological function.
- **Denaturation** is the loss of a biomolecule's shape, which leads to loss of function.
- **Metabolism** is the totality of an organism's chemical reactions, transforming energy and matter.
- Biochemical systems are tightly regulated to maintain a stable internal environment, a principle called **homeostasis**.
- Genetic information flows from **DNA → RNA → Protein**, a concept known as **The Central Dogma**.
- **Catabolism** refers to breakdown pathways that release energy.
- **Anabolism** refers to build-up (biosynthesis) pathways that consume energy.
- Catabolic reactions are **exergonic**, releasing energy, some of which is captured as ATP.
- Anabolic reactions are **endergonic**, consuming energy, often driven by ATP hydrolysis.
- Catabolism is **oxidative**, often releasing electrons captured by carriers like NAD⁺.
- Anabolism is **reductive**, often consuming electrons from carriers like NADPH.
- Catabolism converts complex molecules (carbs, fats) into smaller, simpler molecules like CO₂.
- Anabolism builds complex macromolecules from simple precursors like amino acids and sugars.
- The primary goal of catabolism is to generate usable energy (ATP) and create precursor metabolites.
- The primary goal of anabolism is to synthesize cellular components and store energy.
- Examples of catabolic pathways include Glycolysis, Krebs Cycle, and β-Oxidation of fats.
- Examples of anabolic pathways include Protein synthesis, Glycogenesis, and DNA Replication.
- A **metabolic pathway** is a series of linked, enzyme-catalyzed reactions.
- **ATP (Adenosine Triphosphate)** is the universal "energy currency" of the cell.
- Energy from catabolism is used to phosphorylate **ADP** into **ATP**.
- Anabolic processes hydrolyze **ATP** back to **ADP + Pi**, releasing energy.
- **NAD⁺/NADH & FAD/FADH₂** are redox coenzymes primarily involved in **catabolic** reactions.
- **NADP⁺/NADPH** is a redox coenzyme primarily involved in **anabolic** reactions.
- **Allosteric regulation** occurs when a molecule binds at a site other than the active site, changing enzyme activity.
- **Covalent modification** involves the reversible addition/removal of chemical groups like phosphate.
- **Compartmentalization** separates opposing pathways into different organelles for regulation.
- **Metabolic disorders** are diseases resulting from defects in enzymes or pathways (e.g., PKU, Diabetes).
- The **First Law of Thermodynamics** states energy can be neither created nor destroyed, only converted.



- In biochemistry, chemical potential energy in nutrients is converted into work, heat, or storage.
- The **Second Law of Thermodynamics** states the total entropy (disorder) of the universe always increases in a spontaneous process.
- **Entropy (S)** is a measure of randomness or disorder.
- Living organisms are highly ordered (low entropy) systems.
- Organisms maintain order by constantly increasing the entropy of their surroundings (e.g., releasing heat and waste).
- **Gibbs Free Energy (ΔG)** predicts reaction spontaneity at constant temperature and pressure.
- The equation for Gibbs Free Energy is $\Delta G = \Delta H - T\Delta S$.
- A reaction with $\Delta G < 0$ is **exergonic** and spontaneous (e.g., ATP hydrolysis).
- A reaction with $\Delta G > 0$ is **endergonic** and non-spontaneous (e.g., protein synthesis).
- A reaction with $\Delta G = 0$ is at equilibrium.
- **ATP acts as an energy coupler**, driving endergonic reactions by coupling them to its exergonic hydrolysis.
- **Kinetics** (speed) is separate from **thermodynamics** (spontaneity); enzymes affect kinetics by lowering activation energy.
- **Redox reactions** release free energy which is harnessed in processes like the electron transport chain.
- Water constitutes **70-90%** of cell mass.
- A water molecule has a **bent geometry**, making it a **polar** molecule.
- The **electronegativity** difference between oxygen and hydrogen creates a **dipole moment**.
- Water molecules form up to **four hydrogen bonds** with neighboring molecules.
- Water's **hydrogen bonding network** is responsible for its high cohesiveness, surface tension, and unique thermal properties.
- Water is an **excellent solvent** for polar and charged (**hydrophilic**) substances.
- **Hydration shells** form when water molecules surround solutes.
- Water's **high specific heat capacity (4.184 J/g·°C)** buffers cells against rapid temperature changes.
- Water's **high heat of vaporization (~2260 J/g)** provides a powerful cooling mechanism via evaporation.
- Water is a direct **reactant** in **hydrolysis** reactions, which break bonds.
- Water is a **product** in **dehydration synthesis (condensation)** reactions, which build molecules.
- The **hydrophobic effect** is the clustering of nonpolar molecules to minimize disruption of water's H-bonding network.
- The hydrophobic effect is the primary driving force for **membrane formation, protein folding, and DNA base stacking**.
- **Cohesion** is the attraction between water molecules (via H-bonds).
- **Adhesion** is the attraction of water to other polar surfaces.
- Together, cohesion and adhesion enable **capillary action**.
- Water undergoes slight **autoionization**: $H_2O \rightleftharpoons H^+ + OH^-$.
- The concentration of H^+ defines the **pH** of a solution.
- Proton gradients across membranes, established in water, are crucial for **ATP synthesis** via chemiosmosis.
- Ice is **less dense** than liquid water because hydrogen bonds form a crystalline lattice, causing it to float.
- **Carbon** is the fundamental element of all known life.
- Carbon's **tetravalency** allows it to form four stable covalent bonds.
- **C-C and C-H bonds** are strong, providing stability to large organic molecules.
- Carbon can form **single, double, and triple bonds**, leading to versatile molecular shapes.



- **Catenation** is carbon's ability to form long chains and rings.
- Carbon's moderate **electronegativity** makes its bonds with biogenic elements relatively non-polar.
- The combination of tetravalency and catenation leads to immense **molecular diversity**.
- **Isomers** are molecules with the same formula but different structures.
- **Structural isomers** have different bonding arrangements (e.g., glucose vs. fructose).
- **Stereoisomers (Enantiomers)** are mirror-image molecules.
- Life almost exclusively uses **L-amino acids** and **D-sugars**.
- **Functional groups** are specific, reactive clusters of atoms that determine a molecule's chemical properties.
- The **hydroxyl (-OH)** group is characteristic of alcohols and sugars; it is polar and forms H-bonds.
- The **carbonyl (>C=O)** group is present in aldehydes and ketones (e.g., in sugars).
- The **carboxyl (-COOH)** group is characteristic of organic acids; it is acidic and donates H⁺.
- The **amino (-NH₂)** group is characteristic of amines; it is basic and accepts H⁺.
- The **phosphate (-OPO₃²⁻)** group is found in nucleotides and ATP; it is energetic and acidic.
- The **sulfhydryl (-SH)** group is found in thiols like cysteine; it can form disulfide bonds.
- In **carbohydrates**, carbon forms the sugar backbone (e.g., in glucose and cellulose).
- In **lipids**, carbon forms hydrocarbon chains (fatty acids) and steroid ring systems.
- In **proteins**, carbon forms the amino acid backbone, specifically the central **α-carbon**.
- In **nucleic acids**, carbon forms the pentose sugar and the nitrogenous bases in nucleotides.
- **Cellular respiration** is fundamentally the controlled, stepwise oxidation of carbon-based molecules to release energy.
- The **Carbon Cycle** involves the fixation of CO₂ into organic molecules by autotrophs and their metabolism by heterotrophs.
- **Carbohydrates** are polyhydroxy aldehydes or ketones, or substances that hydrolyze to yield them.
- Biochemically, they are defined by the empirical formula **(CH₂O)_n**, where n ≥ 3.
- **Photosynthesis (Calvin Cycle)** fixes CO₂ into 3-phosphoglycerate, eventually producing glucose and starch.
- The key enzyme in the Calvin Cycle is **Ribulose-1,5-bisphosphate carboxylase/oxygenase (RuBisCO)**.
- **Gluconeogenesis** is the 11-step pathway that synthesizes glucose from non-carbohydrate precursors.
- **Glycogenesis** is the synthesis of glycogen from glucose-1-phosphate.
- Dietary carbohydrates are hydrolyzed by enzymes like **α-amylase**, **sucrase**, **lactase**, and **maltase**.
- **Glycolysis** is the 10-step cytosolic pathway converting glucose to pyruvate.
- **D-sugars** are the stereoisomers predominantly used in biochemistry.
- **Epimers** are diastereomers differing at only one chiral center (e.g., glucose and galactose are C-4 epimers).
- **Anomerism** is a type of diastereomerism specific to the **anomeric carbon** upon ring formation.
- The **α-anomer** has the -OH group *trans* to the CH₂OH group (axial in glucose).
- The **β-anomer** has the -OH group *cis* to the CH₂OH group (equatorial in glucose).
- **Mutarotation** is the interconversion of α and β anomers in solution.
- The predominant ring structures are **pyranose** (6-membered) and **furanose** (5-membered).
- The **anomeric carbon** is the most reactive center in a sugar, involved in glycosidic bond formation and reducing ability.
- A **glycosidic bond** is a covalent bond formed between the anomeric carbon of one sugar and a hydroxyl group of another.
- **Reducing sugars** have a free anomeric carbon and can reduce agents like Cu²⁺ in Benedict's test.



- **Sucrose** is a non-reducing sugar because both of its anomeric carbons are involved in the glycosidic bond.
- High solubility and low molecular weight of monosaccharides create high **osmotic pressure**.
- Cells store glucose as large, insoluble polymers (glycogen, starch) to avoid osmotic issues.
- **Monosaccharides** are the simplest carbohydrate units that cannot be hydrolyzed into smaller carbs.
- Their empirical formula is typically $(\text{CH}_2\text{O})_n$, where $n = 3-9$.
- They are classified as **aldoses** (aldehyde group) or **ketoses** (ketone group).
- Based on carbon count: **Trioses** (3C), **Tetroses** (4C), **Pentoses** (5C), **Hexoses** (6C), **Heptoses** (7C), **Nonoses** (9C).
- Examples: **Glyceraldehyde** (aldotriose), **Dihydroxyacetone** (ketotriose), **Ribose** (aldopentose in RNA), **2-Deoxyribose** (in DNA), **Glucose**, **Galactose**, **Mannose** (aldohexoses), **Fructose** (ketohexose).
- **D-Glucose** is the central metabolic fuel and monomer for major polysaccharides.
- **D-Fructose** is a very sweet dietary sugar metabolized in the liver.
- **D-Galactose** is a component of lactose and glycoconjugates.
- **D-Mannose** is a component of N-linked glycans.
- **D-Ribose** is the backbone of RNA and nucleotide cofactors (ATP, NADH).
- Monosaccharides are highly soluble in water due to extensive hydrogen bonding via -OH groups.
- All monosaccharides (except dihydroxyacetone) are **optically active**.
- All monosaccharides are **reducing sugars**.
- They can form **esters** (e.g., phosphate esters) and **glycosides**.
- **Disaccharides** are formed by two monosaccharides joined by a **glycosidic bond** via dehydration.
- The general molecular formula for two hexoses is $\text{C}_{12}\text{H}_{22}\text{O}_{11}$.
- **Reducing disaccharides** have one free anomeric carbon (e.g., Maltose, Lactose, Cellobiose).
- **Non-reducing disaccharides** have both anomeric carbons involved in the bond (e.g., Sucrose).
- **Sucrose** is composed of **Glucose + Fructose** linked by an $\alpha(1\rightarrow2\beta)$ glycosidic bond; it is non-reducing.
- **Lactose** is composed of **Galactose + Glucose** linked by a $\beta(1\rightarrow4)$ bond; it is reducing.
- **Maltose** is composed of **Glucose + Glucose** linked by an $\alpha(1\rightarrow4)$ bond; it is reducing.
- **Cellobiose** is composed of **Glucose + Glucose** linked by a $\beta(1\rightarrow4)$ bond; it is reducing but not digestible by humans.
- **Trehalose** is composed of **Glucose + Glucose** linked by an $\alpha(1\rightarrow1\alpha)$ bond; it is non-reducing.
- **Polysaccharides (Glycans)** are polymers of monosaccharide units (typically >10).
- They serve as **energy storage** (e.g., starch, glycogen), **structural materials** (e.g., cellulose), or **cell recognition** components.
- **Homopolysaccharides** are composed of one type of monosaccharide (e.g., Starch, Glycogen, Cellulose, Chitin).
- **Heteropolysaccharides** are composed of two or more different monosaccharides (e.g., Hyaluronic acid).
- **Storage polysaccharides** are compact, often branched, and osmotically inert.
- **Structural polysaccharides** are typically fibrous and water-insoluble.
- **Linear polysaccharides** include Cellulose, Amylose, and Chitin.
- **Branched polysaccharides** include Amylopectin and Glycogen.
- **Amylose** is a linear polymer of α -D-glucose with $\alpha(1\rightarrow4)$ linkages.
- **Amylopectin** is a branched polymer of α -D-glucose with $\alpha(1\rightarrow4)$ linkages and $\alpha(1\rightarrow6)$ branches.
- **Glycogen** is the animal storage polysaccharide; it is highly branched with $\alpha(1\rightarrow4)$ and frequent $\alpha(1\rightarrow6)$ linkages.



- **Cellulose** is a structural polysaccharide in plants; it is a linear polymer of β -D-glucose with $\beta(1\rightarrow4)$ linkages.
- **Chitin** is a structural polysaccharide in arthropods and fungi; it is a polymer of N-Acetyl- β -D-Glucosamine (NAG).
- The **Iodine Test** forms colored complexes with helical polysaccharides: deep blue with **Amylose**, reddish-purple with **Amylopectin/Glycogen**, and no color with Cellulose/Chitin.
- Polysaccharides are generally **non-reducing** and **tasteless**.
- **Lipids** are a heterogeneous group of organic compounds that are **insoluble in water (hydrophobic)** but soluble in organic solvents.
- They are characterized by predominantly **nonpolar hydrocarbon chains**.
- They are composed mainly of **carbon, hydrogen, and oxygen** (some contain P, N, S).
- Functions include: **Energy storage** (9 kcal/g), **Structural components** of membranes, **Insulation and protection**, **Hormone precursors**, **Vitamin carriers**, **Signaling molecules**.
- Most biological lipids are **amphipathic** (have both hydrophobic and hydrophilic regions).
- **Saturated fats** have higher melting points and are solid at room temperature.
- **Unsaturated fats** have lower melting points and are liquid at room temperature.
- **Saponification** is the alkaline hydrolysis of triglycerides to produce soap and glycerol.
- The **saponification number** indicates the average chain length of fatty acids in a fat.
- **Hydrogenation** adds H_2 to double bonds in unsaturated fats, converting them to saturated fats.
- The **iodine number** measures the degree of unsaturation in a fat (grams of iodine absorbed by 100g fat).
- **Rancidity** is the spoilage of fats, either by **hydrolytic** (lipase action) or **oxidative** (auto-oxidation) processes.
- **Simple lipids** yield only fatty acids and alcohols upon hydrolysis (e.g., Triacylglycerols, Waxes).
- **Complex lipids** contain additional groups like phosphate, carbohydrates, or nitrogenous bases (e.g., Phospholipids, Glycolipids).
- **Derived lipids** are substances derived from simple or complex lipids (e.g., Fatty acids, Steroids, Terpenes).
- **Triacylglycerols** consist of a **glycerol** backbone esterified with three **fatty acids**.
- **Simple triglycerides** have three identical fatty acids (e.g., Tristearin).
- **Mixed triglycerides** have two or three different fatty acids.
- **Fats** are solid at room temperature, high in saturated fatty acids (e.g., animal fats).
- **Oils** are liquid at room temperature, high in unsaturated fatty acids (e.g., plant oils).
- **Waxes** are esters of **long-chain fatty acids** with **long-chain alcohols** (no glycerol).
- They are hard, pliable, water-repellent, and have high melting points (60-100°C).
- Biological functions include waterproofing (plant cuticle, animal fur), protection, and structural roles (beeswax).
- **Fatty acids** are long hydrocarbon chains with a terminal carboxyl group.
- **Saturated fatty acids** have no double bonds (e.g., Palmitic C16:0, Stearic C18:0).
- **Unsaturated fatty acids** have one or more double bonds.
- **Monounsaturated fatty acids** have one double bond (e.g., Oleic acid C18:1 *cis*).
- **Polyunsaturated fatty acids (PUFAs)** have more than one double bond (e.g., Linoleic C18:2, α -Linolenic C18:3).
- **Essential fatty acids** cannot be synthesized by the body and must be obtained from the diet (e.g., Linoleic, α -Linolenic).
- Melting point **increases** with **increasing chain length**.
- Melting point **decreases** with **increasing unsaturation**.
- *Cis* double bonds lower melting point more than *trans* double bonds.



- Fatty acids regulate **membrane fluidity**: saturated decrease fluidity, unsaturated increase fluidity.
- **β -oxidation** is the major catabolic pathway for fatty acids, yielding large amounts of ATP.
- **Phospholipids** contain a phosphate group and are major components of biological membranes.
- **Glycerophospholipids** have a glycerol backbone, two fatty acids, a phosphate, and an alcohol (e.g., choline, ethanolamine).
- **Phosphatidylcholine (Lecithin)** is a major membrane component and a lung surfactant.
- **Phosphatidylethanolamine (Cephalin)** is abundant in brain and nerve tissue.
- **Phosphatidylserine** is important in apoptosis and blood clotting.
- **Steroids** have a basic structure of four fused carbon rings (**cyclopentanoperhydrophenanthrene nucleus**).
- **Cholesterol** is a key animal sterol, a component of membranes and a precursor for other steroids.
- **Steroid hormones** include sex hormones (estrogens, androgens) and adrenocortical hormones (cortisol, aldosterone).
- **Bile acids** (e.g., cholic acid) emulsify dietary fats.
- **Vitamin D** (calcitriol) is derived from cholesterol/ergosterol and regulates calcium metabolism.
- **Terpenes** are built from **isoprene (C_5H_8)** units.
- **Monoterpenes (C_{10})** are essential oils (e.g., menthol).
- **Diterpenes (C_{20})** include phytol (in chlorophyll) and gibberellins.
- **Tetraterpenes (C_{40})** include carotenoids (e.g., β -carotene).
- **Polyterpenes** include natural rubber.
- **Proteins** are large, complex macromolecules that are polymers of **amino acids**.
- They contain **Carbon (C), Hydrogen (H), Oxygen (O), Nitrogen (N)**, and often **Sulfur (S)**.
- There are **20 standard amino acids** commonly found in proteins.
- Every amino acid (except proline) has a central **α -carbon** bonded to an **amino group ($-NH_2$)**, a **carboxyl group ($-COOH$)**, a **hydrogen atom**, and a variable **side chain (R-group)**.
- At physiological pH (~ 7.4), the amino group is protonated (NH_3^+) and the carboxyl group is deprotonated (COO^-), forming a **zwitterion**.
- All amino acids in proteins are in the **L-configuration**.
- **Peptide bonds** are formed by a condensation reaction between the carboxyl group of one amino acid and the amino group of another, releasing water.
- The peptide bond has **partial double-bond character** due to resonance, making it rigid and planar.
- **Primary structure** is the linear sequence of amino acids linked by peptide bonds.
- **Secondary structure** involves local folding patterns stabilized by hydrogen bonds between backbone atoms.
- The **α -helix** is a right-handed coiled structure stabilized by intra-chain H-bonds between carbonyl O of residue $*n*$ and amide H of residue $*n+4*$.
- The **β -pleated sheet** consists of β -strands stabilized by inter-chain H-bonds; it can be parallel or antiparallel.
- **Beta turns (β -turns)** are tight reverse turns that allow the polypeptide chain to change direction.
- **Tertiary structure** is the overall 3D conformation of a single polypeptide chain.
- Tertiary structure is stabilized by: **Covalent disulfide bonds, Electrostatic (ionic) interactions, Hydrogen bonds, Van der Waals forces**, and the **Hydrophobic effect**.
- The **hydrophobic effect** is the major driving force for protein folding, clustering nonpolar side chains in the interior.
- **Quaternary structure** involves the association of two or more folded polypeptide chains (subunits) into a functional complex.
- Quaternary structure is stabilized by the same non-covalent interactions as tertiary structure (e.g., hemoglobin).



- **Structural proteins** provide mechanical support (e.g., Collagen, Keratin, Actin, Tubulin).
- **Enzymatic proteins** are biological catalysts that speed up specific reactions (e.g., Amylase, ATP Synthase).
- **Transport proteins** carry specific atoms or molecules (e.g., Hemoglobin transports O_2 , Membrane channels).
- **Hormonal proteins** act as chemical messengers (e.g., Insulin, Glucagon, Growth Hormone).
- **Defensive proteins** include antibodies that neutralize pathogens.
- **Storage proteins** store nutrients (e.g., Ferritin stores iron).
- **Receptor proteins** receive chemical signals.
- **Motor proteins** enable movement (e.g., Myosin in muscle contraction).
- **Enzymes** are primarily **globular proteins** that act as catalysts, increasing reaction rates by lowering **activation energy**.
- The **active site** is the region where the substrate binds.
- **Cofactors** are non-protein helper components required for enzyme activity.
- **Inorganic cofactors** (e.g., Mg^{2+} , Zn^{2+}) are called **activators**.
- **Coenzymes** are organic cofactors, often derived from vitamins (e.g., NAD^+ , FAD).
- A **prosthetic group** is a cofactor covalently and permanently bound to the enzyme.
- An enzyme without its cofactor is an **apoenzyme**.
- The complete, active enzyme (apoenzyme + cofactor) is the **holoenzyme**.
- The **Lock and Key Model** (Emil Fischer) proposes a rigid, pre-shaped active site.
- The **Induced Fit Model** (Daniel Koshland) proposes the active site molds around the substrate upon binding.
- Enzyme activity is affected by **temperature, pH, substrate concentration, and enzyme concentration**.
- For most human enzymes, the **optimum temperature** is $\sim 37^\circ C$.
- Above the optimum temperature, enzymes **denature** (lose their 3D structure and activity).
- Each enzyme has an **optimum pH** (e.g., Pepsin $\sim pH 2$, Trypsin $\sim pH 7.8-8.7$).
- At low substrate concentration, reaction rate is proportional to $[S]$ (first-order kinetics).
- At high substrate concentration, the rate plateaus at **V_{max}** (zero-order kinetics) because all active sites are saturated.
- The **Michaelis Constant (K_m)** is the substrate concentration at which the reaction rate is half of V_{max} ; it indicates enzyme affinity (low K_m = high affinity).
- With excess substrate, reaction rate is directly proportional to **enzyme concentration**.
- **Competitive inhibitors** resemble the substrate and bind reversibly to the active site, increasing apparent K_m but not affecting V_{max} .
- **Non-competitive inhibitors** bind reversibly to an allosteric site, decreasing V_{max} but not affecting K_m .
- **Uncompetitive inhibitors** bind only to the Enzyme-Substrate (ES) complex, decreasing both V_{max} and apparent K_m .
- **Irreversible inhibitors** form covalent bonds with the enzyme, permanently inactivating it (e.g., Penicillin).
- **Feedback inhibition** is a regulatory mechanism where the end product of a pathway inhibits the first enzyme.
- **Allosteric regulation**: Effector molecules bind to allosteric sites, changing enzyme activity.
- **Covalent modification**: Reversible addition/removal of groups (e.g., phosphorylation by kinases).
- **Zymogen (proenzyme) activation**: Inactive precursors are activated by proteolytic cleavage (e.g., Pepsinogen \rightarrow Pepsin).
- **Compartmentalization**: Separating enzymes and substrates into specific organelles.



- **Genetic control:** Regulating enzyme synthesis by turning genes on/off.
- **EC 1: Oxidoreductases** catalyze oxidation-reduction reactions (e.g., Alcohol Dehydrogenase).
- **EC 2: Transferases** transfer functional groups (e.g., Hexokinase).
- **EC 3: Hydrolases** catalyze hydrolysis reactions (e.g., Trypsin, Amylase).
- **EC 4: Lyases** cleave bonds by means other than hydrolysis or oxidation, often forming double bonds (e.g., Pyruvate Decarboxylase).
- **EC 5: Isomerases** catalyze intramolecular rearrangements (e.g., Triosephosphate Isomerase).
- **EC 6: Ligases** join two molecules using energy from ATP hydrolysis (e.g., DNA Ligase).
- **Nucleic acids** are biopolymers of **nucleotides** that store, transmit, and express genetic information.
- The two main types are **DNA (Deoxyribonucleic Acid)** and **RNA (Ribonucleic Acid)**.
- DNA contains the sugar **deoxyribose** and bases **A, G, C, T**; it is typically double-stranded and more stable.
- RNA contains the sugar **ribose** and bases **A, G, C, U**; it is typically single-stranded and less stable.
- A **nucleotide** consists of a **pentose sugar**, a **phosphate group**, and a **nitrogenous base**.
- The base is attached to the sugar's **1' carbon**, and the phosphate is attached to the **5' carbon**.
- Nucleotides are linked by **phosphodiester bonds** between the 3' carbon of one sugar and the 5' phosphate of the next.
- **Purine bases** are double-ring structures: **Adenine (A)** and **Guanine (G)**.
- **Pyrimidine bases** are single-ring structures: **Cytosine (C)**, **Thymine (T)** in DNA, **Uracil (U)** in RNA.
- A **nucleoside** is a sugar + base (no phosphate).
- The **primary structure** of DNA is its linear nucleotide sequence.
- The **secondary structure** is the **double helix** (Watson & Crick, 1953).
- The two strands are **antiparallel** (one 5'→3', the other 3'→5').
- Strands are held together by **complementary base pairing** via **hydrogen bonds**: **A=T (2 H-bonds)**, **G≡C (3 H-bonds)**.
- This explains **Chargaff's rules**: **A=T** and **G=C**.
- **Base stacking** (hydrophobic interactions between stacked bases) provides significant stability.
- Helical geometry: diameter ~20 Å, one turn ~34 Å containing **10 base pairs**, distance between bases ~3.4 Å.
- Grooves: **Major groove** (wider, important for protein recognition) and **Minor groove**.
- **B-DNA** is the most common form under physiological conditions.
- **A-DNA** is shorter and wider, formed under dehydrating conditions.
- **Z-DNA** is a left-handed helix formed in sequences with alternating purines/pyrimidines.
- **Supercoiling** is the further twisting of the DNA helix upon itself; **negative supercoiling** (underwinding) is prevalent and facilitates processes like replication.
- **Topoisomerases** are enzymes that cut and reseal DNA strands to manage supercoiling.
- DNA wraps around **histone** proteins to form **nucleosomes**, the fundamental unit of chromatin.
- A nucleosome core consists of an **octamer** of histones (H2A, H2B, H3, H4 x2) with ~147 bp of DNA wrapped around it.
- **Histone H1** binds to linker DNA, stabilizing higher-order structure.
- Nucleosomes coil into a **30 nm fiber**, which forms loops attached to a protein scaffold, ultimately condensing into **chromosomes**.
- **Euchromatin** is less condensed and transcriptionally active.
- **Heterochromatin** is highly condensed and transcriptionally inactive.
- Prokaryotic DNA is circular and organized by **nucleoid-associated proteins (NAPs)**.
- **Nuclear DNA (nDNA)** is linear, organized into chromosomes, and inherited from both parents.



- **Mitochondrial DNA (mtDNA)** is circular, lacks histones, has a higher mutation rate, and is maternally inherited.
- **Chloroplast DNA (cpDNA)** is circular and found in photosynthetic eukaryotes.
- **Coding DNA** sequences are transcribed into mRNA and translated into proteins (~1-2% of human genome).
- **Non-coding DNA** includes regulatory sequences, introns, and repetitive DNA.
- **Repetitive DNA** includes **tandem repeats** (e.g., satellite DNA at centromeres) and **interspersed repeats** (e.g., transposable elements like *Alu*).
- **Unique (Single-Copy) DNA** includes most protein-coding genes.
- **Plasmid DNA** is small, circular, extrachromosomal DNA found in bacteria, often carrying antibiotic resistance genes.
- RNA is primarily **single-stranded** but folds into complex 3D shapes via intramolecular base pairing.
- RNA contains **ribose** with a **2'-OH group**, making it more reactive and less stable than DNA.
- **Primary structure:** linear sequence of nucleotides.
- **Secondary structure:** local folds (stem-loops, hairpins, pseudoknots) via base pairing (A=U, G=C).
- **Tertiary structure:** overall 3D shape stabilized by non-canonical base pairs, stacking, and metal ions.
- **Messenger RNA (mRNA)** carries genetic information from DNA to the ribosome.
- mRNA has a **5' Cap** (7-methylguanosine) for protection and initiation, a **5' UTR**, a **coding sequence** (codons), a **3' UTR**, and a **Poly-A Tail** for stability.
- **Transfer RNA (tRNA)** is the "adapter" molecule that brings amino acids to the ribosome.
- tRNA has a **cloverleaf secondary structure** with an **acceptor stem (3' CCA end)**, **D arm**, **anticodon arm**, and **T ψ C arm**.
- tRNA folds into an **L-shaped tertiary structure**.
- **Ribosomal RNA (rRNA)** forms the core structural and catalytic framework of the ribosome.
- In prokaryotes: small subunit has **16S rRNA**, large subunit has **23S and 5S rRNA**.
- In eukaryotes: small subunit has **18S rRNA**, large subunit has **28S, 5.8S, and 5S rRNA**.
- The **peptidyl transferase** activity (peptide bond formation) is catalyzed by **rRNA**, making it a **ribozyme**.
- **Small Nuclear RNA (snRNA)** components of the spliceosome catalyze pre-mRNA splicing.
- **Small Nucleolar RNA (snoRNA)** guide chemical modifications of other RNAs (rRNA, tRNA).
- **MicroRNA (miRNA)** and **Small Interfering RNA (siRNA)** are involved in gene regulation and silencing via RNA interference (RNAi).
- **Long Non-coding RNA (lncRNA)** are >200 nt RNAs involved in various regulatory processes (e.g., X-inactivation).
- **Ribozymes** are RNA molecules with enzymatic activity (e.g., Group I introns, RNase P).
- **Friedrich Miescher** (1869) isolated "nuclein" (DNA) from cell nuclei.
- **Oswald Avery, Colin MacLeod, Maclyn McCarty** (1944) proved DNA is the genetic material (Avery-MacLeod-McCarty experiment).
- **Alfred Hershey & Martha Chase** (1952) confirmed DNA as genetic material using bacteriophages (Hershey-Chase experiment).
- **Rosalind Franklin & Maurice Wilkins** provided X-ray crystallography data (Photo 51) crucial for determining DNA structure.
- **James Watson & Francis Crick** (1953) proposed the double helix model of DNA.
- **Jöns Jacob Berzelius** (1838) coined the term "protein".
- **Frederick Sanger** (1951-1955) sequenced insulin, the first protein sequenced.
- **Anselme Payen & Jean-François Persoz** (1833) isolated the first enzyme, diastase (amylase).



- **Eduard Buchner** (1897) demonstrated cell-free fermentation, proving enzymatic activity is separate from living cells.
- **James B. Sumner** (1926) first crystallized an enzyme (urease), proving its protein nature.
- **Emil Fischer** elucidated sugar structures and proposed the Lock and Key model for enzymes.
- **William Prout** classified food components into saccharinous, albuminous, and oleaginous (fats).
- **E. Gorter & F. Grendel** (1925) concluded membranes are lipid bilayers.
- **Karl Lohmann** (1929) discovered ATP.
- **Fritz Albert Lipmann** (1941) formulated the concept of ATP as the central energy currency.

M
K

P
R
E
P
A
R
A
T
I
O
N
S



4. Biological Molecules & Enzymes

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. What is the branch of biology that deals with the study of biological molecules and their reactions?

- A) Physiology
- B) Biochemistry
- C) Molecular biology
- D) Genetics

Answer: Biochemistry

2. Water is an excellent solvent for polar substances because of its:

- A) High specific heat
- B) Cohesion
- C) Polarity
- D) Hydrogen bonding

Answer: Polarity

3. The property of water that allows it to resist changes in temperature is due to its:

- A) High specific heat capacity
- B) High heat of vaporization
- C) Cohesion
- D) Adhesion

Answer: High specific heat capacity

4. When water molecules stick to each other, it is called:

- A) Cohesion
- B) Adhesion
- C) Surface tension
- D) Capillary action

Answer: Cohesion

5. When water molecules stick to other surfaces, it is called:

- A) Cohesion
- B) Adhesion
- C) Surface tension
- D) Capillary action

Answer: Adhesion

6. Ice floats on water because:

- A) Ice is denser than water
- B) Ice is less dense than water
- C) Ice has a higher specific heat
- D) Ice has a lower heat of vaporization

Answer: Ice is less dense than water

7. Hydrophobic exclusion refers to:

- A) The attraction between water and nonpolar molecules
- B) The repulsion between water and nonpolar molecules

C) The ionization of water

D) The formation of hydrogen bonds

Answer: The repulsion between water and nonpolar molecules

8. The high specific heat of water is due to:

- A) Hydrogen bonding
- B) Covalent bonding
- C) Ionic bonding
- D) van der Waals forces

Answer: Hydrogen bonding

9. Water is a polar molecule because:

- A) It has a linear shape
- B) The oxygen atom is more electronegative than hydrogen
- C) It has a tetrahedral shape
- D) It can form hydrogen bonds

Answer: The oxygen atom is more electronegative than hydrogen

10. The cohesion of water molecules is responsible for:

- A) High specific heat
- B) High heat of vaporization
- C) Surface tension
- D) Low density of ice

Answer: Surface tension

11. Adhesion of water molecules to cell walls helps in:

- A) Temperature regulation
- B) Capillary action
- C) Floating of ice
- D) Solvent properties

Answer: Capillary action

12. Hydrophobic exclusion is important for:

- A) Formation of lipid bilayers
- B) Solubility of salts in water
- C) High specific heat
- D) Ionization of water

Answer: Formation of lipid bilayers

13. Water's high surface tension is primarily a result of:

- A) Covalent bonding
- B) Ionic bonding
- C) Hydrogen bonding
- D) Van der Waals forces

Answer: Hydrogen bonding

14. Capillary action in plants is possible due to water's properties of:

- A) High specific heat and heat of vaporization
 B) Cohesion and adhesion
 C) Universal solvent nature and polarity
 D) Low density as ice

Answer: Cohesion and adhesion

15. Water's high heat of vaporization is a result of:

- A) Low boiling point
 B) Breaking of hydrogen bonds to change from liquid to vapor
 C) High density
 D) Low specific heat

Answer: Breaking of hydrogen bonds to change from liquid to vapor

16. Carbon's ability to form four stable covalent bonds is termed:

- A) Catenation
 B) Tetravalency
 C) Isomerism
 D) Electronegativity

Answer: Tetravalency

17. The immense structural diversity of organic molecules is primarily due to carbon's:

- A) High electronegativity
 B) Ability to form long chains and rings (catenation)
 C) Small atomic radius
 D) Formation of ionic bonds

Answer: Ability to form long chains and rings (catenation)

18. Molecules with the same molecular formula but different structural arrangements are called:

- A) Isotopes
 B) Polymers
 C) Isomers
 D) Enantiomers

Answer: Isomers

19. In biochemistry, which functional group is characteristic of an alcohol?

- A) -COOH
 B) -NH₂
 C) -OH
 D) -SH

Answer: -OH

20. The general formula for monosaccharides is:

- A) (CH₂O)_n

- B) (CHO)_n
 C) (CH₃O)_n
 D) (C₂H₅O)_n

Answer: (CH₂O)_n

21. Which of the following is a monosaccharide with the empirical formula C₆H₁₂O₆?

- A) Ribose
 B) Glucose
 C) Sucrose
 D) Maltose

Answer: Glucose

22. Monosaccharides with a carbonyl group at the end of the carbon chain are called:

- A) Ketoses
 B) Aldoses
 C) Trioses
 D) Pentoses

Answer: Aldoses

23. Which of the following is a ketose sugar?

- A) Glucose
 B) Fructose
 C) Galactose
 D) Ribose

Answer: Fructose

24. The sugar found in RNA is:

- A) Deoxyribose
 B) Ribose
 C) Glucose
 D) Fructose

Answer: Ribose

25. The sugar found in DNA is:

- A) Deoxyribose
 B) Ribose
 C) Glucose
 D) Fructose

Answer: Deoxyribose

26. Which of the following is a pentose sugar?

- A) Glucose
 B) Fructose
 C) Ribose
 D) Galactose

Answer: Ribose

27. The bond formed between two monosaccharides is called:

- A) Peptide bond
 B) Glycosidic bond
 C) Ester bond

D) Phosphodiester bond

Answer: Glycosidic bond

28. Sucrose is composed of:

- A) Glucose and glucose
- B) Glucose and fructose
- C) Glucose and galactose
- D) Fructose and galactose

Answer: Glucose and fructose

29. Lactose is composed of:

- A) Glucose and glucose
- B) Glucose and fructose
- C) Glucose and galactose
- D) Fructose and galactose

Answer: Glucose and galactose

30. Maltose is composed of:

- A) Glucose and glucose
- B) Glucose and fructose
- C) Glucose and galactose
- D) Fructose and galactose

Answer: Glucose and glucose

31. Which of the following is a storage polysaccharide in plants?

- A) Starch
- B) Glycogen
- C) Cellulose
- D) Chitin

Answer: Starch

32. Which of the following is a storage polysaccharide in animals?

- A) Starch
- B) Glycogen
- C) Cellulose
- D) Chitin

Answer: Glycogen

33. Which of the following is a structural polysaccharide in plants?

- A) Starch
- B) Glycogen
- C) Cellulose
- D) Chitin

Answer: Cellulose

34. Which of the following is a structural polysaccharide in arthropods and fungi?

- A) Starch
- B) Glycogen
- C) Cellulose
- D) Chitin

Answer: Chitin

35. Cellulose is made up of:

- A) α -glucose units
- B) β -glucose units
- C) α -fructose units
- D) β -fructose units

Answer: β -glucose units

36. Starch is made up of:

- A) α -glucose units
- B) β -glucose units
- C) α -fructose units
- D) β -fructose units

Answer: α -glucose units

37. Glycogen is similar to starch but has more:

- A) Branching
- B) Glucose units
- C) Hydrogen bonds
- D) Nitrogen atoms

Answer: Branching

38. Cellulose digestion in herbivores is aided by:

- A) Amylase
- B) Cellulase
- C) Lactase
- D) Maltase

Answer: Cellulase

39. Chitin is a polymer of:

- A) N-acetyl glucosamine
- B) N-acetyl galactosamine
- C) Glucose
- D) Galactose

Answer: N-acetyl glucosamine

40. The building blocks of proteins are:

- A) Nucleotides
- B) Amino acids
- C) Fatty acids
- D) Monosaccharides

Answer: Amino acids

41. How many amino acids are commonly found in proteins?

- A) 10
- B) 20
- C) 30
- D) 40

Answer: 20

42. Amino acids are linked together by:

- A) Glycosidic bonds
- B) Peptide bonds
- C) Ester bonds

M
K
P
R
E
P
A
R
A
T
I
O
N
S

D) Phosphodiester bonds

Answer: Peptide bonds

43. The bond between the carboxyl group of one amino acid and the amino group of another is called:

- A) Hydrogen bond
- B) Ionic bond
- C) Peptide bond
- D) Disulfide bond

Answer: Peptide bond

44. The sequence of amino acids in a protein is called its:

- A) Primary structure
- B) Secondary structure
- C) Tertiary structure
- D) Quaternary structure

Answer: Primary structure

45. α -helix and β -pleated sheet are examples of:

- A) Primary structure
- B) Secondary structure
- C) Tertiary structure
- D) Quaternary structure

Answer: Secondary structure

46. The overall three-dimensional shape of a protein is called:

- A) Primary structure
- B) Secondary structure
- C) Tertiary structure
- D) Quaternary structure

Answer: Tertiary structure

47. When a protein consists of more than one polypeptide chain, it has:

- A) Primary structure
- B) Secondary structure
- C) Tertiary structure
- D) Quaternary structure

Answer: Quaternary structure

48. Which of the following is a fibrous protein?

- A) Hemoglobin
- B) Collagen
- C) Enzyme
- D) Antibody

Answer: Collagen

49. Which of the following is a globular protein?

- A) Collagen
- B) Keratin

C) Hemoglobin

D) Elastin

Answer: Hemoglobin

50. Lipids are generally insoluble in water because they are:

- A) Hydrophilic
- B) Hydrophobic
- C) Polar
- D) Ionic

Answer: Hydrophobic

51. Which of the following is not a lipid?

- A) Triglyceride
- B) Phospholipid
- C) Steroid
- D) Protein

Answer: Protein

52. A triglyceride is composed of:

- A) Glycerol and two fatty acids
- B) Glycerol and three fatty acids
- C) Glycerol and one fatty acid
- D) Glycerol and phosphate

Answer: Glycerol and three fatty acids

53. The bond between glycerol and fatty acid in a triglyceride is called:

- A) Peptide bond
- B) Glycosidic bond
- C) Ester bond
- D) Phosphodiester bond

Answer: Ester bond

54. A fatty acid with no double bonds is called:

- A) Saturated
- B) Unsaturated
- C) Monounsaturated
- D) Polyunsaturated

Answer: Saturated

55. A fatty acid with one double bond is called:

- A) Saturated
- B) Monounsaturated
- C) Polyunsaturated
- D) None of the above

Answer: Monounsaturated

56. A fatty acid with more than one double bond is called:

- A) Saturated
- B) Monounsaturated
- C) Polyunsaturated

D) None of the above

Answer: Polyunsaturated

57. Phospholipids are composed of:

- A) Glycerol, two fatty acids, and a phosphate group
- B) Glycerol, three fatty acids
- C) Glycerol, two fatty acids, and a sugar
- D) Glycerol, one fatty acid, and a phosphate group

Answer: Glycerol, two fatty acids, and a phosphate group

58. In a phospholipid, the head is:

- A) Hydrophobic
- B) Hydrophilic
- C) Nonpolar
- D) Neutral

Answer: Hydrophilic

59. In a phospholipid, the tails are:

- A) Hydrophobic
- B) Hydrophilic
- C) Polar
- D) Charged

Answer: Hydrophobic

60. Steroids are composed of:

- A) Four fused carbon rings
- B) Three fused carbon rings
- C) Two fused carbon rings
- D) One carbon ring

Answer: Four fused carbon rings

61. Cholesterol is an example of a:

- A) Triglyceride
- B) Phospholipid
- C) Steroid
- D) Wax

Answer: Steroid

62. The building blocks of nucleic acids are:

- A) Nucleotides
- B) Amino acids
- C) Fatty acids
- D) Monosaccharides

Answer: Nucleotides

63. A nucleotide consists of:

- A) Sugar, phosphate, and base
- B) Sugar and base
- C) Phosphate and base
- D) Sugar and phosphate

Answer: Sugar, phosphate, and base

64. Which base is found in DNA but not in RNA?

- A) Adenine
- B) Guanine
- C) Cytosine
- D) Thymine

Answer: Thymine

65. Which base is found in RNA but not in DNA?

- A) Adenine
- B) Guanine
- C) Uracil
- D) Thymine

Answer: Uracil

66. In DNA, adenine pairs with:

- A) Thymine
- B) Cytosine
- C) Guanine
- D) Uracil

Answer: Thymine

67. In DNA, guanine pairs with:

- A) Thymine
- B) Cytosine
- C) Adenine
- D) Uracil

Answer: Cytosine

68. The bonds that hold the two strands of DNA together are:

- A) Covalent bonds
- B) Hydrogen bonds
- C) Ionic bonds
- D) Peptide bonds

Answer: Hydrogen bonds

69. According to Chargaff's rule, in DNA the amount of adenine equals the amount of:

- A) Guanine
- B) Cytosine
- C) Thymine
- D) Uracil

Answer: Thymine

70. The model of DNA double helix was proposed by:

- A) Darwin and Mendel
- B) Watson and Crick
- C) Franklin and Wilkins
- D) Chargaff and Franklin

Answer: Watson and Crick

71. Which of the following types of RNA carries genetic information from DNA to the ribosome?

- A) mRNA

- B) tRNA
- C) rRNA
- D) snRNA

Answer: mRNA

72. Which of the following types of RNA is a component of the ribosome?

- A) mRNA
- B) tRNA
- C) rRNA
- D) snRNA

Answer: rRNA

73. Which of the following types of RNA carries amino acids to the ribosome?

- A) mRNA
- B) tRNA
- C) rRNA
- D) snRNA

Answer: tRNA

74. The triplet code on mRNA is called a:

- A) Codon
- B) Anticodon
- C) Gene
- D) Nucleotide

Answer: Codon

75. The complementary triplet on tRNA is called a:

- A) Codon
- B) Anticodon
- C) Gene
- D) Nucleotide

Answer: Anticodon

76. ATP is an example of a:

- A) Nucleotide
- B) Amino acid
- C) Fatty acid
- D) Monosaccharide

Answer: Nucleotide

77. The bond between nucleotides in a nucleic acid is called:

- A) Peptide bond
- B) Glycosidic bond
- C) Ester bond
- D) Phosphodiester bond

Answer: Phosphodiester bond

78. Which of the following is a conjugated molecule?

- A) Glycolipid
- B) Glycoprotein
- C) Lipoprotein

- D) All of the above

Answer: All of the above

79. The pH of pure water at 25°C is:

- A) 0
- B) 7
- C) 14
- D) 1

Answer: 7

80. The dissociation constant of water (K_w) at 25°C is:

- A) 10^{-7}
- B) 10^{-14}
- C) 10^{-1}
- D) 10^{-10}

Answer: 10^{-14}

81. Which of the following is a triose?

- A) Glyceraldehyde
- B) Ribose
- C) Glucose
- D) Fructose

Answer: Glyceraldehyde

82. In an aqueous solution, glucose exists mainly as:

- A) Open chain
- B) Ring form
- C) Linear chain
- D) Branched chain

Answer: Ring form

83. α -glucose and β -glucose are:

- A) Structural isomers
- B) Stereoisomers
- C) Enantiomers
- D) Diastereomers

Answer: Stereoisomers

84. Sucrose is a non-reducing sugar because:

- A) It has no free anomeric carbon
- B) It has a free anomeric carbon
- C) It is a disaccharide
- D) It is sweet

Answer: It has no free anomeric carbon

85. Lactose intolerance is due to deficiency of the enzyme:

- A) Sucrase
- B) Lactase
- C) Maltase
- D) Amylase

Answer: Lactase

86. Amylose is a linear polymer of glucose linked by:

- A) α -1,4 linkages
- B) α -1,6 linkages
- C) β -1,4 linkages
- D) β -1,6 linkages

Answer: α -1,4 linkages

87. Amylopectin is a branched polymer of glucose with branches at:

- A) α -1,4 linkages
- B) α -1,6 linkages
- C) β -1,4 linkages
- D) β -1,6 linkages

Answer: α -1,6 linkages

88. Cellulose is digested by:

- A) Amylase
- B) Cellulase
- C) Lactase
- D) Sucrase

Answer: Cellulase

89. Which of the following is a reducing sugar?

- A) Sucrose
- B) Lactose
- C) Trehalose
- D) None of the above

Answer: Lactose

90. The test for reducing sugars is:

- A) Iodine test
- B) Benedict's test
- C) Biuret test
- D) Emulsion test

Answer: Benedict's test

91. The test for starch is:

- A) Iodine test
- B) Benedict's test
- C) Biuret test
- D) Emulsion test

Answer: Iodine test

92. The test for proteins is:

- A) Iodine test
- B) Benedict's test
- C) Biuret test
- D) Emulsion test

Answer: Biuret test

93. The test for lipids is:

- A) Iodine test
- B) Benedict's test
- C) Biuret test
- D) Emulsion test

Answer: Emulsion test

94. Amino acids are amphoteric because they have:

- A) Both amino and carboxyl groups
- B) Only amino group
- C) Only carboxyl group
- D) Side chain

Answer: Both amino and carboxyl groups

95. At physiological pH, amino acids exist as:

- A) Cations
- B) Anions
- C) Zwitterions
- D) Neutral molecules

Answer: Zwitterions

96. Which of the following amino acids has a sulfur atom in its side chain?

- A) Cysteine
- B) Methionine
- C) Both A and B
- D) Neither

Answer: Both A and B

97. Which of the following amino acids is acidic?

- A) Aspartic acid
- B) Glutamic acid
- C) Both A and B
- D) Neither

Answer: Both A and B

98. Which of the following amino acids is basic?

- A) Lysine
- B) Arginine
- C) Histidine
- D) All of the above

Answer: All of the above

99. The amino acid that can form disulfide bonds is:

- A) Cysteine
- B) Methionine
- C) Serine
- D) Threonine

Answer: Cysteine

100. The peptide bond has partial double bond character because of:

- A) Resonance
- B) Hydrogen bonding
- C) Ionic bonding
- D) van der Waals forces

Answer: Resonance

101. In an α -helix, hydrogen bonds form between the carbonyl group of one amino acid and the amino group of another amino acid that is:

- A) 3 residues away
- B) 4 residues away
- C) 5 residues away
- D) 6 residues away

Answer: 4 residues away

102. In a β -pleated sheet, the polypeptide chains are:

- A) Parallel or antiparallel
- B) Only parallel
- C) Only antiparallel
- D) Neither

Answer: Parallel or antiparallel

103. Which of the following forces stabilizes the tertiary structure of proteins?

- A) Hydrogen bonds
- B) Ionic bonds
- C) Hydrophobic interactions
- D) All of the above

Answer: All of the above

104. Hemoglobin is an example of a protein with:

- A) Primary structure
- B) Secondary structure
- C) Tertiary structure
- D) Quaternary structure

Answer: Quaternary structure

105. Myoglobin is an example of a protein with:

- A) Primary structure
- B) Secondary structure
- C) Tertiary structure
- D) Quaternary structure

Answer: Tertiary structure

106. Denaturation of proteins involves disruption of:

- A) Primary structure
- B) Secondary and tertiary structure
- C) Quaternary structure
- D) Both B and C

Answer: Both B and C

107. Which of the following can denature proteins?

- A) Heat
- B) pH change
- C) Organic solvents

D) All of the above

Answer: All of the above

108. Lipids are not polymers in the same sense as carbohydrates because they are not built from repeating monomers. However, they are formed by dehydration reactions between:

- A) Glycerol and fatty acids
- B) Fatty acids only
- C) Glycerol only
- D) None of the above

Answer: Glycerol and fatty acids

109. The melting point of fatty acids increases with:

- A) Increasing chain length
- B) Increasing unsaturation
- C) Decreasing chain length
- D) Branching

Answer: Increasing chain length

110. The melting point of fatty acids decreases with:

- A) Increasing unsaturation
- B) Increasing chain length
- C) Saturation
- D) None of the above

Answer: Increasing unsaturation

111. Essential fatty acids for humans include:

- A) Linoleic acid
- B) Linolenic acid
- C) Arachidonic acid
- D) All of the above

Answer: All of the above

112. Phospholipids are amphipathic because they have:

- A) Hydrophilic head and hydrophobic tails
- B) Hydrophobic head and hydrophilic tails
- C) Both hydrophilic
- D) Both hydrophobic

Answer: Hydrophilic head and hydrophobic tails

113. Cholesterol is a precursor for:

- A) Bile acids
- B) Steroid hormones
- C) Vitamin D
- D) All of the above

Answer: All of the above

114. Waxes are esters of long-chain fatty acids and:

- A) Glycerol

M
K
P
R
E
P
A
R
A
T
I
O
N
S

- B) Long-chain alcohols
- C) Phosphate
- D) Sugars

Answer: Long-chain alcohols

115. Prostaglandins are derived from:

- A) Arachidonic acid
- B) Palmitic acid
- C) Stearic acid
- D) Oleic acid

Answer: Arachidonic acid

116. Aspirin works by inhibiting the synthesis of:

- A) Steroids
- B) Prostaglandins
- C) Terpenes
- D) Phospholipids

Answer: Prostaglandins

117. Lipids are transported in the blood as:

- A) Lipoproteins
- B) Glycolipids
- C) Phospholipids
- D) Triglycerides

Answer: Lipoproteins

118. Which lipoprotein has the highest density?

- A) HDL
- B) LDL
- C) VLDL
- D) Chylomicrons

Answer: HDL

119. Which lipoprotein is often called "bad cholesterol"?

- A) HDL
- B) LDL
- C) VLDL
- D) Chylomicrons

Answer: LDL

120. The nitrogenous base in nucleotides is attached to the sugar at which carbon?

- A) 1'
- B) 2'
- C) 3'
- D) 4'

Answer: 1'

121. The phosphate group in nucleotides is attached to the sugar at which carbon?

- A) 3'
- B) 4'
- C) 5'

- D) 6'

Answer: 5'

122. A nucleotide without a phosphate is called a:

- A) Nucleoside
- B) Nucleotide
- C) Nucleosome
- D) Nucleoprotein

Answer: Nucleoside

123. ATP is a nucleoside triphosphate. It contains:

- A) Adenine, ribose, three phosphates
- B) Adenine, deoxyribose, three phosphates
- C) Adenine, ribose, two phosphates
- D) Adenine, deoxyribose, two phosphates

Answer: Adenine, ribose, three phosphates

124. In DNA, how many hydrogen bonds are between A and T?

- A) 2
- B) 3
- C) 4
- D) 1

Answer: 2

125. In DNA, how many hydrogen bonds are between G and C?

- A) 2
- B) 3
- C) 4
- D) 1

Answer: 3

126. The two strands of DNA are:

- A) Antiparallel
- B) Parallel
- C) Both
- D) Neither

Answer: Antiparallel

127. The distance between two consecutive base pairs in DNA is about:

- A) 0.34 nm
- B) 3.4 nm
- C) 2 nm
- D) 10 nm

Answer: 0.34 nm

128. Which of the following is a pyrimidine base found in both DNA and RNA?

- A) Cytosine
- B) Thymine
- C) Uracil

D) Adenine

Answer: Cytosine

129. Which of the following is a purine base found in both DNA and RNA?

- A) Adenine
- B) Guanine
- C) Both A and B
- D) Neither

Answer: Both A and B

130. The difference between deoxyribose and ribose is that deoxyribose has one less oxygen at which carbon?

- A) 2'
- B) 3'
- C) 4'
- D) 5'

Answer: 2'

131. Nucleosomes are composed of DNA wrapped around a core of:

- A) Histones
- B) RNA
- C) Lipids
- D) Carbohydrates

Answer: Histones

132. The ends of chromosomes are protected by structures called:

- A) Telomeres
- B) Centromeres
- C) Kinetochores
- D) Origins of replication

Answer: Telomeres

133. Glycolipids are composed of:

- A) Carbohydrate and lipid
- B) Protein and lipid
- C) Carbohydrate and protein
- D) Nucleic acid and lipid

Answer: Carbohydrate and lipid

134. Glycoproteins are composed of:

- A) Carbohydrate and protein
- B) Lipid and protein
- C) Carbohydrate and lipid

D) Nucleic acid and protein

Answer: Carbohydrate and protein

135. Lipoproteins are composed of:

- A) Lipid and protein
- B) Carbohydrate and protein
- C) Carbohydrate and lipid
- D) Nucleic acid and protein

Answer: Lipid and protein

136. The amount of heat required to convert 1 gram of water into vapor is called:

- A) Specific heat
- B) Heat of vaporization
- C) Latent heat
- D) Boiling point

Answer: Heat of vaporization

137. When water ionizes, it produces:

- A) H^+ and O^{2-}
- B) H_2 and O_2
- C) H^+ and OH^-
- D) H_3O^+ and OH^-

Answer: H^+ and OH^-

138. How many hydrogen bonds can a single water molecule form?

- A) 1
- B) 2
- C) 3
- D) 4

Answer: 4

139. When water freezes, its volume:

- A) Increases
- B) Decreases
- C) Remains the same
- D) First increases then decreases

Answer: Increases

140. In an α -helix, there are how many amino acids per turn?

- A) 3.6
- B) 4.0
- C) 5.0
- D) 6.0

Answer: 3.6



Chapter: 5

Bioenergetics

Bioenergetics is defined as the study of energy transduction in living organisms, encompassing the capture of light energy and its storage as chemical bond energy, as well as the utilization of energetic compounds through processes such as aerobic respiration.

Photosynthesis

Photosynthesis is the fundamental **biochemical process** by which **photoautotrophs**—primarily plants, algae, and certain bacteria (like cyanobacteria)—capture and convert **light energy** from the sun into **stable chemical energy** stored in the bonds of organic molecules (e.g., glucose). This process transforms inorganic carbon (from atmospheric carbon dioxide) and water into energy-rich carbohydrates, releasing oxygen as a byproduct.

B. Overview of the Process

At its core, photosynthesis is a two-stage process occurring within specialized organelles called **chloroplasts**:

1. **Light-Dependent Reactions:** Occur in the **thylakoid membranes**. Light energy is captured by pigments like chlorophyll and used to split water molecules (**photolysis**), releasing oxygen. This energy is converted into short-term energy carriers (**ATP and NADPH**).
2. **Light-Independent Reactions (Calvin Cycle):** Occur in the **stroma** of the chloroplast. The chemical energy from ATP and NADPH is used to fix atmospheric carbon dioxide (**CO₂**) into organic sugars, like glucose. This stage does not require light directly but depends on the products of the light reactions.

2. The General Equation

The overall chemical equation summarizes the net input and output of the complex process:



- **Reactants:** Carbon Dioxide (**CO₂**) from the air and Water (**H₂O**) from the soil.
- **Products:** Glucose (**C₆H₁₂O₆**), a simple sugar used for energy and structure, and **Molecular Oxygen (O₂)**, released into the atmosphere.

The Chloroplast: Structure and Function

The **chloroplast** is a double-membrane-bound organelle.

Structure	Description & Function
Outer & Inner Membrane	Envelope the organelle. The inner membrane is selectively permeable.
Intermembrane Space	Narrow space between the two membranes.
Stroma	Viscous fluid filling the chloroplast. Site of the light-independent reactions (Calvin Cycle) . Contains enzymes, DNA, ribosomes, and starch grains.
Thylakoids	A system of interconnected, flattened, membranous sacs.
Grana (sing. granum)	Stacks of thylakoids (like a stack of coins).
Lumen	The interior space of a thylakoid.
Chlorophyll & Accessory Pigments	Embedded in the thylakoid membranes . Arranged in photosystems (I & II) to form light-harvesting complexes.
Function Summary:	Thylakoids = Site of light-dependent reactions (produce ATP, NADPH, O ₂). Stroma = Site of light-independent reactions (use ATP & NADPH to fix CO ₂ into sugar).

The Two Stages of Photosynthesis

A. Light-Dependent Reactions

- **Location:** Thylakoid Membranes.

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091



Bioenergetics: One Liner

M
K
P
R
E
P
A
R
A
T
I
O
N
S

5. Bioenergetics

- **Bioenergetics** is the study of energy flow and transformation through living systems.
- **Photosynthesis** converts **light energy** into **chemical energy** stored in organic molecules.
- **Cellular respiration** breaks down organic molecules to release usable energy (**ATP**).
- The **primary electron donor** in photosynthesis is **water**, which is oxidized to **O₂**.
- The **final electron acceptor** in aerobic respiration is **oxygen**, which is reduced to **water**.
- **ATP (Adenosine Triphosphate)** is the universal **energy currency** of the cell.
- **ATP synthesis** is driven by a **proton motive force** across a membrane in both processes.
- **Chloroplasts** are the organelles where **photosynthesis** occurs in plants and algae.
- **Mitochondria** are the organelles where **aerobic respiration** occurs in eukaryotic cells.
- The general equation for photosynthesis is $6\text{CO}_2 + 6\text{H}_2\text{O} + \text{Light} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2$.
- The general equation for aerobic respiration is $\text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2 \rightarrow 6\text{CO}_2 + 6\text{H}_2\text{O} + \text{ATP}$.
- Photosynthesis occurs in two main stages: the **light-dependent reactions** and the **Calvin cycle**.
- The **light-dependent reactions** occur in the **thylakoid membranes** of chloroplasts.
- The **Calvin cycle** occurs in the **stroma** of chloroplasts.
- Cellular respiration comprises **glycolysis**, the **Krebs cycle**, and the **electron transport chain**.
- **Glycolysis** occurs in the **cytosol** and is anaerobic.
- The **Krebs cycle** and **electron transport chain** occur in the **mitochondrial matrix** and **inner membrane**, respectively.
- **Photosystems I and II** are protein-pigment complexes in the thylakoid membrane that capture light.
- **P680** is the reaction center chlorophyll *a* dimer of **Photosystem II**.
- **P700** is the reaction center chlorophyll *a* dimer of **Photosystem I**.
- **Photolysis** of water occurs at the **oxygen-evolving complex** of Photosystem II.
- **Non-cyclic photophosphorylation** involves both photosystems and produces **ATP, NADPH, and O₂**.
- **Cyclic photophosphorylation** involves only Photosystem I and produces **ATP** only.
- **NADP⁺** is reduced to **NADPH** during the light reactions, acting as a **reducing power** for the Calvin cycle.
- The **Calvin cycle** uses **ATP and NADPH** to fix **CO₂** into carbohydrate.
- **Rubisco (Ribulose-1,5-bisphosphate carboxylase/oxygenase)** is the key enzyme for **CO₂ fixation**.
- **Rubisco** can also catalyze **photorespiration**, a wasteful process.
- **C3 plants** fix CO₂ directly via the Calvin cycle and are susceptible to photorespiration.
- **C4 plants** use the **Hatch-Slack pathway** to concentrate CO₂ in bundle sheath cells, minimizing photorespiration.
- **CAM plants** open stomata at night to fix CO₂ into organic acids, storing it for daytime use.
- **Glycolysis** breaks one glucose into two molecules of **pyruvate**, producing a net gain of **2 ATP and 2 NADH**.
- The **energy investment phase** of glycolysis consumes 2 ATP.
- The **energy payoff phase** of glycolysis produces 4 ATP and 2 NADH.
- In the absence of oxygen, **fermentation** occurs to regenerate **NAD⁺** from **NADH**.
- **Lactic acid fermentation** reduces pyruvate to lactate (e.g., in muscle cells, some bacteria).
- **Alcoholic fermentation** reduces pyruvate to ethanol and **CO₂** (e.g., in yeast).
- Before the Krebs cycle, **pyruvate oxidation** converts pyruvate to **acetyl-CoA**, producing **NADH and CO₂**.

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. The primary source of energy for most living organisms is:

- A. Glucose
- B. ATP
- C. Sunlight
- D. Heat

Correct Answer: Sunlight

2. The process by which plants convert light energy into chemical energy is called:

- A. Respiration
- B. Photosynthesis
- C. Transpiration
- D. Fermentation

Correct Answer: Photosynthesis

3. The overall chemical equation for photosynthesis is:

- A. $C_6H_{12}O_6 + 6O_2 \rightarrow 6CO_2 + 6H_2O + \text{energy}$
- B. $6CO_2 + 6H_2O \rightarrow C_6H_{12}O_6 + 6O_2$
- C. $C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$
- D. $6CO_2 + 12H_2O \rightarrow C_6H_{12}O_6 + 6O_2 + 6H_2O$

Correct Answer: $6CO_2 + 12H_2O \rightarrow C_6H_{12}O_6 + 6O_2 + 6H_2O$

4. The organelles where photosynthesis takes place are:

- A. Mitochondria
- B. Ribosomes
- C. Chloroplasts
- D. Leucoplasts

Correct Answer: Chloroplasts

5. The main photosynthetic pigment in green plants is:

- A. Chlorophyll a
- B. Chlorophyll b
- C. Xanthophyll
- D. Carotene

Correct Answer: Chlorophyll a

6. Which part of the chloroplast contains the photosynthetic pigments?

- A. Stroma
- B. Granum
- C. Thylakoid membrane
- D. Inner membrane

Correct Answer: Thylakoid membrane

7. The light-independent reactions of photosynthesis occur in the:

- A. Thylakoid lumen
- B. Stroma
- C. Grana

D. Intermembrane space

Correct Answer: Stroma

8. During the light reactions of photosynthesis, the initial electron donor is:

- A. NADP+
- B. Water
- C. Oxygen
- D. Carbon dioxide

Correct Answer: Water

9. The final electron acceptor in the light reactions of photosynthesis is:

- A. Oxygen
- B. Water
- C. NADP+
- D. Cytochrome

Correct Answer: NADP+

10. The products of the light-dependent reactions used in the Calvin cycle are:

- A. ATP and NADPH
- B. ATP and NADH
- C. ADP and NADP+
- D. Glucose and O₂

Correct Answer: ATP and NADPH

11. In photosynthesis, oxygen is released as a by-product from the splitting of:

- A. Carbon dioxide
- B. Glucose
- C. Water
- D. ATP

Correct Answer: Water

12. The enzyme responsible for fixing CO₂ in the Calvin cycle is:

- A. RuBP carboxylase-oxygenase (Rubisco)
- B. PEP carboxylase
- C. ATP synthase
- D. Phosphofructokinase

Correct Answer: RuBP carboxylase-oxygenase (Rubisco)

13. The five-carbon compound that accepts CO₂ in the Calvin cycle is:

- A. Ribulose-1,5-bisphosphate (RuBP)
- B. 3-Phosphoglycerate (3-PGA)
- C. Glyceraldehyde-3-phosphate (G3P)
- D. Phosphoenolpyruvate (PEP)

Correct Answer: Ribulose-1,5-bisphosphate (RuBP)

14. The first stable product of CO₂ fixation in C₃ plants is:

Chapter 6

DNA & Chromosomes

- **Nucleic acids** are linear, unbranched polymers of nucleotides that serve as the primary information-carrying molecules in all living organisms and viruses.
- They constitute the **chemical basis of heredity** and direct cellular metabolism.
- **Historical Perspective:** Initially, proteins were favored as genetic material due to their chemical complexity. The series of key experiments established DNA as the universal genetic material.

Landmark Experiments Proving DNA as Genetic Material

Experiment (Year)	Scientists	Key Organism/System	Method & Findings	Conclusion
Transformation (1928)	Frederick Griffith	<i>Streptococcus pneumoniae</i> strains (S-virulent, R-avirulent)	Heat-killed S + live R → mice died; live S recovered.	A " transforming principle " transferred genetic traits.
Identification of Transforming Principle (1944)	Oswald Avery, Colin MacLeod, Maclyn McCarty	<i>S. pneumoniae</i>	Purified components; only DNA fraction caused transformation; DNase destroyed activity.	DNA is the transforming substance and hereditary material in bacteria.
Hershey-Chase (1952)	Alfred Hershey, Martha Chase	Bacteriophage T2 & <i>E. coli</i>	Radioactive labeling: ³² P (DNA) entered bacteria; ³⁵ S (protein) remained outside.	DNA, not protein, is the genetic material that enters host cells.
Chargaff's Rules (1949)	Erwin Chargaff	Multiple species	Chemical analysis of DNA base composition.	A=T, G=C; (A+G)=(T+C); base ratios are species-specific.
X-ray Diffraction (1950-53)	Rosalind Franklin, Maurice Wilkins	DNA fibers	Produced "Photo 51": helical structure, 2 nm diameter, 3.4 nm repeat, 0.34 nm between bases.	Provided critical data for double helix model.
Double Helix Model (1953)	James Watson, Francis Crick	N/A	Combined Chargaff's rules and Franklin's X-ray data to build a molecular model.	Proposed the antiparallel double helix with specific A-T and G-C pairing .

Central Dogma of Molecular Biology

Original Concept (Crick, 1958): DNA → RNA → Protein

Revised Concept: Includes exceptions:

- **Reverse transcription** (RNA → DNA) by retroviruses (e.g., HIV)
- **RNA replication** in RNA viruses

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091



Chromosomes And DNA: One-Liner

M
K
P
R
E
P
A
R
A
T
I
O
N
S

- **Chromosomes** become visible under a light microscope only during **mitosis and meiosis**.
- A chromosome is composed of **DNA, histone proteins, non-histone proteins, and a small amount of RNA**.
- The **centromere** is the site of **kinetochore** assembly for microtubule attachment during cell division.
- The **nucleolar organizer region (NOR)** contains genes for **ribosomal RNA (rRNA)** synthesis.
- **Telomeres** are specialized structures at chromosome ends, consisting of **highly conserved, tandemly repeated, non-coding DNA sequences** (e.g., TTAGGG in vertebrates).
- Telomeres function to **prevent end-to-end fusion, degradation, and solve the end-replication problem**.
- **Satellite DNA** is located near the centromere and consists of highly repetitive, non-coding sequences.
- Based on centromere position: **Metacentric** (V-shaped), **Submetacentric** (L-shaped), **Acrocentric** (J-shaped), **Telocentric** (I-shaped, rare in humans).
- Based on function: **Autosomes** (22 pairs in humans, govern somatic traits) and **Sex Chromosomes (Allosomes)** (XX female, XY male).
- The **Y chromosome** contains the **SRY (Sex-determining Region Y)** gene, the primary testis-determining factor.
- **Dosage compensation** in females involves the random inactivation of one X chromosome, forming a **Barr body**, regulated by the **XIST lncRNA**.
- Based on centromere number: **Monocentric** (standard), **Dicentric** (unstable), **Acentric** (lost during division), **Holocentric** (entire length acts as centromere, e.g., *C. elegans*).
- The fundamental unit of chromatin is the **nucleosome**, consisting of ~146 bp of DNA wrapped **1.65 times** around a **histone octamer** (two each of H2A, H2B, H3, H4).
- **Histone H1** binds to linker DNA, facilitating the folding of nucleosomes into a **30-nm chromatin fiber**.
- **Euchromatin** is less condensed, transcriptionally active, and replicates early in S-phase.
- **Heterochromatin** is highly condensed and transcriptionally silent, subdivided into **constitutive** (permanent) and **facultative** (reversible, e.g., Barr body).
- Higher-order packaging involves forming **radial loop domains** attached to a **nuclear scaffold**, culminating in the **metaphase chromosome**.
- **Griffith's Experiment (1928)** demonstrated **bacterial transformation**, indicating a "**transforming principle**."
- **Avery, MacLeod, and McCarty (1944)** conclusively identified the transforming principle as **DNA**.
- **Hershey-Chase Experiment (1952)** used radioactive labeling (³²P for DNA, ³⁵S for protein) in bacteriophage T2 to prove **DNA is the genetic material** that enters the host cell.
- **Chargaff's Rules** state that in double-stranded DNA, the amount of **Adenine equals Thymine and Guanine equals Cytosine** (A=T, G=C), and the **total purines equal total pyrimidines** (A+G = T+C).
- The **Watson-Crick model** describes DNA as a **right-handed, antiparallel double helix**.
- The two strands are held together by **complementary base pairing**: **A=T** (2 hydrogen bonds) and **G≡C** (3 hydrogen bonds).
- The **sugar-phosphate backbone** is on the outside, with **nitrogenous bases stacked inward**.
- The double helix has a constant diameter of **2 nm** due to purine-pyrimidine pairing.
- The helix makes a full turn every **10 base pairs**, with a **helical pitch of 3.4 nm**.
- The structure features a **major groove** and a **minor groove**, which are binding sites for proteins.

6. DNA & Chromosomes

- **DNA Barcoding:** Uses a standard gene region (e.g., mitochondrial COI) for species identification.
- **CRISPR-Cas9:** A genome editing tool for targeted gene knockout/knock-in.

Practice MCQs

1. Which nitrogenous base is found in RNA but not in DNA?

- A) Adenine
- B) Guanine
- C) Thymine
- D) Uracil

Answer: Uracil

2. The Meselson-Stahl experiment demonstrated that DNA replication is:

- A) Conservative
- B) Dispersive
- C) Semiconservative
- D) Non-conservative

Answer: Semiconservative

3. Which enzyme is responsible for synthesizing RNA primers during DNA replication?

- A) DNA polymerase I
- B) DNA polymerase III
- C) Primase
- D) Ligase

Answer: Primase

4. In the Watson-Crick model of DNA, adenine pairs with:

- A) Guanine
- B) Cytosine
- C) Thymine
- D) Uracil

Answer: Thymine

5. Which type of RNA carries amino acids to the ribosome during translation?

- A) mRNA
- B) tRNA
- C) rRNA
- D) snRNA

Answer: tRNA

6. The condition characterized by trisomy 21 is:

- A) Turner syndrome
- B) Klinefelter syndrome
- C) Down syndrome
- D) Cri-du-chat syndrome

Answer: Down syndrome

7. Which of the following is a purine base?

- A) Cytosine

B) Thymine

C) Uracil

D) Adenine

Answer: Adenine

8. The Hershey-Chase experiment used which isotopes to label DNA and protein?

- A) ^{14}C and ^3H
- B) ^{32}P and ^{35}S
- C) ^{15}N and ^{14}N
- D) ^{18}O and ^2H

Answer: ^{32}P and ^{35}S

9. Which enzyme relieves supercoiling ahead of the replication fork?

- A) Helicase
- B) Topoisomerase
- C) Primase
- D) Ligase

Answer: Topoisomerase

10. The genetic code is said to be degenerate because:

- A) One codon codes for multiple amino acids
- B) One amino acid can be coded by multiple codons
- C) It is the same in all organisms
- D) It has start and stop signals

Answer: One amino acid can be coded by multiple codons

11. Which histone protein is not part of the nucleosome core octamer?

- A) H1
- B) H2A
- C) H3
- D) H4

Answer: H1

12. Transcription in eukaryotes is carried out by which RNA polymerase for mRNA?

- A) RNA polymerase I
- B) RNA polymerase II
- C) RNA polymerase III
- D) RNA polymerase IV

Answer: RNA polymerase II

13. Which of the following mutations changes a codon to a stop codon?

- A) Missense
- B) Nonsense

Chapter 7

Variation and Genetics

Introduction

- **Genetics** is the scientific study of **heredity** (transmission of traits from parents to offspring) and **variation** (differences among individuals).
- **Inheritance**, the process encompassing both heredity and variation, is crucial for evolution and speciation.
- Since **genes** control heredity and variation, genetics is fundamentally the study of genes.
- **Molecular Basis:** A gene is a specific DNA sequence that codes for a polypeptide via **transcription** (DNA to mRNA in nucleus) and **translation** (mRNA to protein at ribosome).

Fundamental Genetic Concepts

- **Gene** – Basic unit of heredity; a segment of DNA coding for a polypeptide/trait. (*Example: The gene for flower color in peas.*)
- **Allele** – Alternative form of a gene at the same locus. (*Example: The alleles for purple (P) or white (p) flowers.*)
- **Locus** – Specific position of a gene on a chromosome.
- **Genotype** – Genetic makeup of an individual. (*Example: PP, Pp, or pp.*)
- **Phenotype** – Observable expression of a trait. (*Example: Purple or white flowers.*)
- **Homozygous** – Having two identical alleles for a gene. (*Example: PP or pp.*)
- **Heterozygous** – Having two different alleles for a gene. (*Example: Pp.*)
- **Hemizygous** – Having only one allele for a gene (e.g., X-linked genes in males).
- **Wild type** – Most common phenotype in natural populations.
- **Mutant phenotype** – Trait alternative to wild type.
- **Gene Pool** – All alleles present in a breeding population at a given time.

Mendelian Principles & Crosses

- **Law of Segregation (Principle of Segregation)** – Alleles separate during gamete formation. (*Mendel's pea plant experiments.*)
- **Law of Independent Assortment** – Genes for different traits assort independently during gamete formation.
- **P generation** – Parental generation.
- **F₁ generation** – First filial generation.
- **F₂ generation** – Second filial generation.
- **True-breeding (Pure breeding)** – Organisms that produce identical offspring when self-fertilized.
- **Monohybrid cross** – Cross involving one trait. (*Example: Crossing pure-breeding tall and dwarf pea plants.*)
- **Dihybrid cross** – Cross involving two traits. (*Example: Crossing plants differing in seed shape and color.*)
- **Testcross** – Cross between an individual with unknown genotype and a homozygous recessive individual.

Extensions to Mendelian Genetics

- **Complete Dominance** – One allele completely masks the other. (*Example: Mendel's pea traits.*)
- **Incomplete dominance** – Heterozygote shows an intermediate phenotype. (*Example: Pink flowers from red and white snapdragons.*)
- **Codominance** – Both alleles are fully expressed in the heterozygote. (*Example: AB blood type; speckled chicken feathers.*)



Variation & Genetics: One Liner

- **Genetics** is the scientific study of **heredity** (transmission of characteristics) and **variation** (differences among individuals).
- **Gene**: The **basic unit of heredity**; a specific DNA sequence that codes for a functional product (usually a protein).
- **Locus**: The specific **physical location of a gene** on a chromosome.
- **Allele**: **Alternative forms of the same gene** that occupy corresponding loci on homologous chromosomes.
- **Gene Pool**: The **complete set of all alleles** present in all individuals of a breeding population at a given time.
- **Phenotype**: The **observable characteristics** of an organism, resulting from genotype and environment.
- **Genotype**: The **genetic constitution** of an organism for a particular trait; the specific combination of alleles.
- **Homozygous**: A condition where **both alleles at a given locus are identical** (e.g., TT or tt).
- **Heterozygous**: A condition where **the two alleles at a given locus are different** (e.g., Tt).
- **Dominant Allele**: An allele that **expresses its phenotype even in a heterozygous state**.
- **Recessive Allele**: An allele whose **phenotypic effect is masked by a dominant allele** and is only expressed when homozygous.
- **True-breeding (Pure-breeding)**: Individuals that, upon self-fertilization, produce offspring identical to themselves for a given trait over generations.
- **Wild Type**: The normal gene or phenotype found in a natural population.
- **Mutant**: An organism or gene with a changed DNA sequence.
- **Carrier**: A heterozygous individual carrying a recessive allele for a genetic disorder without expressing it.
- **Test Cross**: A cross between an individual with a dominant phenotype (unknown genotype) and a homozygous recessive individual to determine the unknown genotype.
- **Gregor Johann Mendel (1822-1884)**, an Augustinian monk, is recognized as the **founder of classical genetics**.
- He conducted **hybridization experiments on garden pea (*Pisum sativum*)** for over eleven years.
- His success was due to: using **true-breeding lines**, studying **seven pairs of contrasting traits**, performing **controlled crosses**, and applying **quantitative/statistical analysis**.
- The **P generation** is the parental true-breeding generation.
- The **F₁ (First Filial) generation** is the first generation of offspring from the P cross.
- The **F₂ (Second Filial) generation** is produced by self-fertilization or interbreeding of F₁ individuals.
- Mendel's work was **rediscovered independently in 1900** by Correns, De Vries, and Tschermak.
- The **Law of Segregation** states that the **two alleles for a heritable trait segregate (separate) during gamete formation** and end up in different gametes.
- Each gamete carries **only one allele** for each trait.
- **Monohybrid Cross** studies the inheritance of a **single trait**.
- Example: Cross between true-breeding round (RR) and wrinkled (rr) pea seeds.
- F₁ Generation: All offspring are **round (Rr)**, demonstrating **complete dominance**.
- F₂ Generation (from selfing F₁): Phenotypic ratio of **3 Round : 1 Wrinkled**; Genotypic ratio of **1 RR : 2 Rr : 1 rr**.
- The physical basis for segregation is the **separation of homologous chromosomes during Anaphase I of Meiosis**.
- The **Law of Independent Assortment** states that **genes for different traits assort independently of one another during gamete formation**.

Practice MCQs

- 1. What is the basic unit of heredity that codes for a functional product like a protein?**
 A) Allele
 B) Locus
 C) Gene
 D) Chromosome
Answer: Gene
- 2. The specific physical location of a gene on a chromosome is called its:**
 A) Allele
 B) Genome
 C) Locus
 D) Phenotype
Answer: Locus
- 3. Alternative forms of the same gene that occupy corresponding loci on homologous chromosomes are known as:**
 A) Genotypes
 B) Phenotypes
 C) Alleles
 D) Linkage groups
Answer: Alleles
- 4. The complete set of all alleles present in all individuals of a breeding population at a given time is the:**
 A) Genome
 B) Karyotype
 C) Gene pool
 D) Genotype frequency
Answer: Gene pool
- 5. The genetic constitution of an organism for a particular trait is its:**
 A) Phenotype
 B) Allele
 C) Genotype
 D) Karyotype
Answer: Genotype
- 6. The observable characteristics resulting from genotype and environment define the:**
 A) Genotype
 B) Allele
 C) Phenotype
 D) Locus
Answer: Phenotype
- 7. An individual with two identical alleles at a given locus is said to be:**
 A) Heterozygous
 B) Hemizygous
 C) Homozygous
 D) Homogametic
Answer: Homozygous
- 8. An allele that expresses its phenotypic effect even in a heterozygous state is termed:**
 A) Recessive
 B) Codominant
 C) Dominant
 D) Incompletely dominant
Answer: Dominant
- 9. Who is recognized as the founder of classical genetics?**
 A) Charles Darwin
 B) Thomas Hunt Morgan
 C) Gregor Mendel
 D) Alfred Sturtevant
Answer: Gregor Mendel
- 10. Mendel's success was due in part to his use of which experimental plant?**
 A) Drosophila
 B) Garden pea
 C) Maize
 D) Snapdragon
Answer: Garden pea
- 11. The law stating that two alleles for a trait segregate during gamete formation is the Law of:**
 A) Independent Assortment
 B) Dominance
 C) Segregation
 D) Purity of Gametes
Answer: Segregation
- 12. In a monohybrid cross of two heterozygous individuals, the expected phenotypic ratio in the F₂ generation is:**
 A) 1:2:1
 B) 3:1
 C) 9:3:3:1
 D) 1:1
Answer: 3:1
- 13. A cross between an individual with a dominant phenotype (unknown**



Chapter 8

Biotechnology

Recombinant DNA Technology (rDNA Technology)

- **Timeframe:** Emerged in the mid-1970s, building on discoveries like restriction enzymes (1970) and DNA ligase.
- **Core Concept:** Artificially combining DNA sequences from **different species** that would not normally interbreed.
- **Mechanism:** Uses **restriction enzymes** and **DNA ligase** to insert a gene of interest into a **vector** (e.g., plasmid), which is then introduced into a host cell (e.g., *E. coli*).
- **Primary Goal:** To **amplify (clone)** a specific DNA segment for detailed study, or to express a protein of interest.
- **Impact:**
 - Revolutionized molecular biology by enabling:
 - **Gene structure analysis** (introns, exons, promoters, enhancers).
 - Study of **gene regulation** and **development**.
 - Tracing **evolutionary relationships** through conserved genes.
- **Key Derivative Fields:**
 1. **Genetic Engineering:** The deliberate modification of an organism's genetic material to introduce **new traits**.
 2. **Biotechnology:** The **industrial and commercial application** of biological systems (cells, enzymes, organisms) to create products or services.
 - **Applications:** Medicine (therapeutics, diagnostics), agriculture (GM crops), industry (enzymes), forensics, environmental remediation.
- **Illustrative Example – GloFish®:**
 - First genetically engineered **pet**.
 - Genes for **green fluorescent protein (GFP)** from jellyfish (*Aequorea victoria*) or **red fluorescent protein** from sea corals are inserted into zebrafish embryos.
 - Used in research as **biological markers** and for aesthetic purposes in aquaria.

DNA Cloning: Core Techniques & Tools

A. Restriction Enzymes – "Molecular Scissors"

- **Origin:** Bacterial defense systems against viral (bacteriophage) DNA.
- **Function:** **Endonucleases** that recognize and cut DNA at specific **palindromic sequences** (4-8 bp long), known as **restriction sites**.
- **Examples & Naming:**
 - **EcoRI:** From *Escherichia coli* strain RY13. Cuts at 5'-GAATTC-3'.
 - **HindIII:** From *Haemophilus influenzae* strain Rd. Cuts at 5'-AAGCTT-3'.
- **Types of Cuts:**
 1. **Sticky Ends:** Staggered cut leaving short, single-stranded **overhangs**. They are complementary and can easily anneal with fragments cut by the same enzyme. *Most useful for cloning.*
 2. **Blunt Ends:** Cut straight across both strands. Requires more efficient ligation.
- **Essential Partner Enzyme:** **DNA Ligase** (often from T4 bacteriophage) forms **phosphodiester bonds** to seal nicks in the sugar-phosphate backbone, creating a stable recombinant molecule.

B. Vectors for Cloning

- **Definition:** A DNA molecule that acts as a **vehicle** to carry foreign genetic material into a host cell.
- **Ideal Features:**
 1. **Origin of Replication (ORI):** Allows independent replication within the host.



Biotechnology: One-Liners

- **Biotechnology** is the use of living organisms or their processes to make useful products.
- **Modern biotechnology** combines molecular biology, genetics, and bioinformatics with traditional techniques like selective breeding.
- **Genetic Engineering** is the direct manipulation of an organism's genome to modify its characteristics.
- The foundation of modern biotechnology is **Recombinant DNA (rDNA) Technology**.
- **Recombinant DNA** is a single DNA molecule formed by combining DNA from two different sources.
- The first recombinant DNA molecule was created by **Paul Berg in 1972** by inserting viral DNA into bacterial DNA.
- **Herbert Boyer and Stanley Cohen (1973)** demonstrated that genes from a toad could be expressed and inherited in bacteria.
- The ability to isolate and manipulate DNA accelerated biotechnological discovery and novel applications.
- The **21st century** is often called the "Century of Biology" due to the rapid growth of genetic engineering.
- **Restriction Endonucleases** are bacterial enzymes that cut DNA at specific nucleotide sequences.
- They act as a bacterial defense system by cutting (**restricting**) invading viral DNA.
- The first restriction enzyme was isolated by **Hamilton Smith** from *Haemophilus influenzae*.
- **Werner Arber, Hamilton Smith, and Daniel Nathans** shared the 1978 Nobel Prize for the discovery of restriction enzymes.
- **Type II restriction enzymes** are most useful in biotechnology as they cut at precise locations within their recognition sites.
- The **recognition site** for most Type II enzymes is a **palindromic sequence** (reads the same on both strands).
- Cutting DNA with restriction enzymes can produce **sticky ends** (single-stranded overhangs) or **blunt ends**.
- **Sticky ends** are more useful in genetic engineering as they facilitate the joining of DNA from different sources.
- Bacteria protect their own DNA from their restriction enzymes by **methylation** (adding CH₃ groups to) the nucleotides at their restriction sites.
- Restriction enzymes are named after the **genus, species, and strain** of the source bacterium (e.g., *EcoRI* from *Escherichia coli*).
- **DNA Ligase** is an enzyme that catalyzes the formation of a **phosphodiester bond** between adjacent nucleotides.
- It permanently joins compatible ends of DNA fragments, sealing recombinant DNA molecules.
- It performs the same function as the DNA ligase that joins Okazaki fragments during cellular DNA replication.
- **Gel Electrophoresis** is a technique that separates DNA fragments based on their **size**.
- It uses an electric field to drive negatively charged DNA through a gel matrix (agarose or polyacrylamide).
- **Smaller DNA fragments** migrate faster and farther through the gel than larger fragments.
- DNA in the gel is visualized using a **fluorescent dye** that binds to DNA (e.g., ethidium bromide).
- A **vector** is a DNA molecule used to carry foreign DNA into a host cell.
- The most common vector is a **plasmid**, a small, circular, extra-chromosomal DNA molecule found in bacteria.
- An ideal vector must have an **Origin of Replication (Ori)** for independent replication in the host.

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. What is the primary function of restriction enzymes in recombinant DNA technology?

- A) Joining DNA fragments
- B) Cutting DNA at specific sequences
- C) Synthesizing DNA from RNA
- D) Amplifying DNA sequences

Answer: Cutting DNA at specific sequences

2. Which enzyme is responsible for sealing nicks between DNA fragments during recombinant DNA formation?

- A) Restriction endonuclease
- B) DNA polymerase
- C) Reverse transcriptase
- D) DNA ligase

Answer: DNA ligase

3. What are the single-stranded complementary ends produced by some restriction enzymes called?

- A) Blunt ends
- B) Sticky ends
- C) Palindrome ends
- D) Ligase ends

Answer: Sticky ends

4. Which of the following is commonly used as a vector in genetic engineering?

- A) Ribosome
- B) Plasmid
- C) Mitochondrion
- D) Lysosome

Answer: Plasmid

5. What does PCR stand for in molecular biology?

- A) Protein Chain Reaction
- B) Polymerase Chain Reaction
- C) Polymer Copying Reaction
- D) Phosphate Chain Reaction

Answer: Polymerase Chain Reaction

6. Which heat-stable enzyme is essential for PCR amplification?

- A) DNA polymerase I
- B) Reverse transcriptase
- C) Taq polymerase
- D) RNA polymerase

Answer: Taq polymerase

7. In PCR, what is the purpose of the annealing step?

- A) To denature DNA strands
- B) To synthesize new DNA strands

- C) To allow primers to bind to template DNA
- D) To seal DNA fragments

Answer: To allow primers to bind to template DNA

8. What is the term for an organism that has had a foreign gene inserted into its genome?

- A) Mutant
- B) Clone
- C) Transgenic organism
- D) Hybrid

Answer: Transgenic organism

9. Which disease has been treated using ex vivo gene therapy by correcting bone marrow stem cells?

- A) Cystic fibrosis
- B) Hypercholesterolemia
- C) Severe Combined Immunodeficiency (SCID)
- D) Diabetes

Answer: Severe Combined Immunodeficiency (SCID)

10. What is the missing enzyme in patients suffering from Severe Combined Immunodeficiency (SCID)?

- A) DNA polymerase
- B) Adenosine deaminase
- C) Reverse transcriptase
- D) Restriction enzyme

Answer: Adenosine deaminase

11. In cystic fibrosis gene therapy, how is the correct gene typically delivered to lung cells?

- A) Through oral tablets
- B) Via aerosol liposomes
- C) By direct injection into blood
- D) Using bacterial vectors

Answer: Via aerosol liposomes

12. Which technique is used to separate DNA fragments based on their size?

- A) Centrifugation
- B) Chromatography
- C) Gel electrophoresis
- D) PCR

Answer: Gel electrophoresis

13. What is the name of the international project that successfully sequenced the entire human genome?

- A) Human Proteome Project
- B) Human Genome Project
- C) ENCODE Project



Chapter 9

Evolution

- **Evolution:** Descent with modification; change in allele frequencies in populations over time.
- **Organic Evolution:** Biological evolution through genetic change and natural selection.
- **Microevolution:** Change in allele frequencies within a population over generations.
- **Macroevolution:** Large-scale evolutionary changes (speciation, extinction) over geological time.
- **Common Descent:** All organisms share a common ancestor.

Special Creation vs. Evolution

Aspect	Special Creation	Evolution
Origin of Species	Independently created	Descended from common ancestors
Change Over Time	Fixed, immutable	Continuously changing
Mechanism	Divine intervention	Natural processes (selection, drift, etc.)
Evidence Base	Religious texts	Multiple scientific disciplines
Scientific Status	Non-testable, non-scientific	Well-supported scientific theory

Origin of Life & Evolution of Cellular Life

I. Origin of Life (Abiogenesis)

A. Prebiotic Conditions on Early Earth (~4.6 - 3.9 Ga)

- **Hadean Eon:** Hot, volcanic, frequent asteroid impacts, no free oxygen.
- **Atmosphere:** Reducing (H₂, CH₄, NH₃, H₂O, CO₂, N₂). No O₂ layer, high UV radiation.
- **Key Requirements for Life:**
 1. Source of organic molecules (monomers).
 2. Mechanism to polymerize monomers.
 3. Self-replication (information storage).
 4. Compartmentalization (protocell membranes).

B. Key Experiments & Hypotheses for Organic Molecule Formation

1. **Miller-Urey Experiment (1953):** Simulated early Earth atmosphere with electrical sparks produced amino acids and other organics.
2. **Extraterrestrial Origins (Panspermia/Meteorites):** Murchison meteorite contains amino acids and nucleobases.
3. **Hydrothermal Vent Hypothesis (A Primary Focus): Submarine alkaline hydrothermal vents (e.g., Lost City-type) are a leading theory.**
 - **Why Vents?** Provide a compelling environment for life's origin.
 - **Energy Gradient:** Natural proton gradient (alkaline vent fluid vs. acidic ocean) mimics modern cellular chemiosmosis (ATP production).
 - **Mineral Catalysts:** Porous chimneys of iron-sulfide (FeS) and mackinawite act as inorganic catalysts and compartment walls.
 - **Conditions:** Stable, protected from surface UV radiation and impacts.
 - **Organic Synthesis:** H₂ and CO₂/CO in vent fluids can react via Fischer-Tropsch-type reactions on catalytic mineral surfaces to form organic molecules.
 - **The "Protometabolism First" Model:** Networks of chemical reactions within vent pores could evolve complexity before the emergence of genetic code or membranes.

C. From Molecules to Cells

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

M
K
P
R
E
P
A
R
A
T
I
O
N
S

9. Evolution

Evolution: One Liner

- **Evolution** is defined as **descent with modification** or a change in **allele frequencies** in a population over time.
- **Organic Evolution** refers to biological evolution through **genetic change** and **natural selection**.
- **Microevolution** is a change in **allele frequencies within a population** over generations.
- **Macroevolution** refers to large-scale evolutionary changes like **speciation** and **extinction** over geological time.
- **Common Descent** is the principle that all organisms share a **common ancestor**.
- Lamarck proposed the first comprehensive evolutionary theory based on **use and disuse** of organs and the **inheritance of acquired characteristics**.
- He believed that organs used extensively become **larger and stronger**, while disused organs **deteriorate and disappear**.
- A classic example is the **giraffe's long neck**, which Lamarck explained as a result of generations stretching to reach leaves.
- His theory is considered **transformational**, suggesting individuals change during their lifetime and pass these changes to offspring.
- **Weismann's Germplasm Theory (1892)** disproved Lamarckism by distinguishing **heritable germ cells** from **non-heritable somatic cells**.
- **Experimental evidence** against Lamarckism includes Weismann's tail-cutting experiment in mice over 80 generations, which produced no tailless offspring.
- Darwin's theory is based on observations from his **HMS Beagle voyage (1831-1836)**, particularly in the **Galápagos Islands**.
- Key influences on Darwin included **Charles Lyell's uniformitarianism** (deep geological time) and **Thomas Malthus's essay** on population growth and struggle for existence.
- Darwin's theory of **natural selection** rests on five key observations: **overproduction** of offspring, **population stability**, **struggle for existence**, **heritable variation**, and **differential survival/reproduction**.
- **Natural selection** is a **non-random process** where individuals with favorable heritable variations are more likely to survive and reproduce.
- Darwin's concept of **descent with modification** states all organisms are related through a **branching evolutionary tree** from a common ancestor.
- **Alfred Russel Wallace** independently conceived natural selection, prompting Darwin to publish *On the Origin of Species*.
- Criticisms of Darwin's original theory included its inability to explain the **source of variations**, the **origin of complex adaptations** ("arrival of the fittest"), **vestigial organs**, and the **mechanism of inheritance**.
- Proposed that new species arise suddenly via **large, discontinuous mutations**
- Integrated **Darwinian natural selection** with **Mendelian genetics** and **population biology**.
- Redefined evolution as a **change in allele frequencies** in a population's **gene pool**.
- Key contributors included **R.A. Fisher, J.B.S. Haldane, Sewall Wright, Theodosius Dobzhansky, Ernst Mayr, George G. Simpson, and Julian Huxley**.
- It recognizes **mutation** and **recombination** as sources of variation, **natural selection** as the primary adaptive mechanism, and the importance of **genetic drift** and **gene flow**.
- It shifted the unit of evolution from the **individual** (Darwin) to the **population (gene pool)**.
- **Directional Selection**: Favors one extreme phenotype (e.g., peppered moth melanism, antibiotic resistance).
- **Stabilizing Selection**: Favors intermediate phenotypes (e.g., human birth weight).
- **Disruptive Selection**: Favors both extremes over intermediates (e.g., black-bellied seed cracker finch beak size).

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. What is the primary mechanism of evolution according to Darwin's theory?

- A) Inheritance of acquired characteristics
- B) Use and disuse of organs
- C) Natural selection
- D) Genetic drift

Answer: Natural selection

2. Which scientist first proposed a comprehensive theory of evolution based on the inheritance of acquired characteristics?

- A) Charles Darwin
- B) Alfred Russel Wallace
- C) Jean-Baptiste Lamarck
- D) Gregor Mendel

Answer: Jean-Baptiste Lamarck

3. Structures that are similar in structure but different in function, indicating common ancestry, are called:

- A) Analogous structures
- B) Vestigial structures
- C) Homologous structures
- D) Convergent structures

Answer: Homologous structures

4. Which of the following is a condition required for Hardy-Weinberg equilibrium?

- A) Non-random mating
- B) Small population size
- C) No gene flow
- D) Presence of natural selection

Answer: No gene flow

5. The wing of a bird and the wing of an insect are examples of:

- A) Homologous structures
- B) Vestigial structures
- C) Analogous structures
- D) Divergent evolution

Answer: Analogous structures

6. What does the Hardy-Weinberg equation $p^2 + 2pq + q^2 = 1$ represent?

- A) Phenotype frequencies
- B) Allele frequencies
- C) Genotype frequencies
- D) Mutation rates

Answer: Genotype frequencies

7. Which of the following provides the strongest evidence for common ancestry among all aerobic organisms?

- A) Presence of hemoglobin
- B) Presence of chlorophyll
- C) Presence of cytochrome c

D) Presence of cellulose

Answer: Presence of cytochrome c

8. According to Lamarck, the long neck of the giraffe evolved due to:

- A) Natural selection for longer necks
- B) Genetic drift in a small population
- C) Inheritance of characteristics acquired through stretching
- D) Mutation in the neck vertebrae gene

Answer: Inheritance of characteristics acquired through stretching

9. What is the ultimate source of new genetic variation in a population?

- A) Genetic drift
- B) Gene flow
- C) Mutation
- D) Natural selection

Answer: Mutation

10. The random change in allele frequencies in a small population is known as:

- A) Gene flow
- B) Natural selection
- C) Genetic drift
- D) Mutation pressure

Answer: Genetic drift

11. The study of geographical distribution of species is known as:

- A) Paleontology
- B) Biogeography
- C) Comparative anatomy
- D) Molecular biology

Answer: Biogeography

12. Which of the following is a vestigial structure in humans?

- A) Femur
- B) Appendix
- C) Humerus
- D) Tibia

Answer: Appendix

13. Neo-Darwinism is best described as the synthesis of:

- A) Lamarckism and mutation theory
- B) Darwinism and Lamarckism
- C) Mendel's genetics and Darwin's evolution
- D) Biogeography and paleontology

Answer: Mendel's genetics and Darwin's evolution

14. What type of selection favors individuals at both extremes of a phenotypic range?

- A) Directional selection



Chapter 10

Ecology & Ecosystem

- **Ecology:** Scientific study of interactions between organisms and their biotic and abiotic environment. Coined by Ernst Haeckel from Greek *oikos* (household) + *logy* (study).
- **Ecosystem:** Dynamic complex of biotic communities and their abiotic environment interacting as a functional unit through energy flows and biogeochemical cycles. Coined by Arthur Tansley (1935) to emphasize interconnectedness.
- **Environment:** All abiotic (non-living: climate, soil, water) and biotic (living: plants, animals, microbes) factors influencing an organism.
- **Biosphere:** Thin, life-supporting layer of Earth where all ecosystems exist.

M
K
P
R
E
P
A
R
A
T
I
O
N
S

Levels of Ecological Organization

1. **Organism:** Individual living entity.
2. **Population:** Group of interbreeding individuals of the same species in a specific area.
3. **Community:** Assemblage of different populations living and interacting in a defined area.
4. **Ecosystem:** Community + physical environment, interacting through nutrient cycling and energy flow.
5. **Biome:** Large geographical region with distinct climate and characteristic community.
6. **Biosphere:** All ecosystems collectively.

Key Ecological Concepts

- **Habitat:** Physical space where an organism lives.
- **Ecological Niche:** Multidimensional concept describing the functional role of a species (resources used, conditions tolerated).
 - *Fundamental Niche:* Full range theoretically usable.
 - *Realized Niche:* Actual range occupied due to interspecific interactions.
- **Metapopulation:** Set of local populations linked by immigration/emigration. The **Glanville fritillary butterfly** in Finland exists as scattered local populations in dry meadows, connected by occasional migration.
- **Symbiosis:** Close, long-term biological interaction between two different species (parasitic, mutualistic, or commensal).
- **Mutualism:** Clownfish and sea anemones.
- **Parasitism:** Tapeworms in mammals.
- **Commensalism:** Barnacles on whales.

ECOSYSTEM STRUCTURE

A. Abiotic Components

- **Physical Factors:**
 - Solar radiation (1–2% converted via photosynthesis).
 - Temperature (affects metabolic rates via Q_{10} relationships).
 - Water availability (creates productivity gradients).
 - Soil texture (water holding capacity, nutrient retention).
- **Chemical Factors:**
 - Nutrient availability (Liebig's Law of the Minimum).
 - Redox potential (influences nutrient speciation).
 - pH, salinity, oxygen availability.

Food Chain

A **food chain** is a **linear sequence** showing how energy and nutrients move from one organism to another in an ecosystem. It follows a single path.

Example of a simple food chain:

Grass → Grasshopper → Frog → Snake → Hawk

Key characteristics:

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091



Ecology & Ecosystems: One Liner

- **Ecology** is the scientific study of interactions between organisms and their **biotic** (living) and **abiotic** (non-living) environment.
- The term **ecology** was coined by the German zoologist **Ernst Haeckel**.
- **Environment** encompasses all abiotic (climate, soil, water) and biotic (other organisms) factors influencing an organism.
- An **ecosystem** is a dynamic complex of biotic communities and their abiotic environment interacting as a functional unit through energy flow and nutrient cycling.
- The term **ecosystem** was coined by British ecologist **Arthur Tansley (1935)**.
- The **Biosphere** is the thin, life-supporting layer of Earth (from ocean depths to the atmosphere) where all ecosystems exist.
- Ecological organization follows a hierarchy: **Organism** → **Population** → **Community** → **Ecosystem** → **Biome** → **Biosphere**.
- A **species** is a group of organisms that can interbreed freely in nature and produce fertile offspring.
- A **population** is a group of interbreeding individuals of the same species living in a specific geographical area at the same time.
- A **community** is an assemblage of different populations living and interacting in a defined area.
- A **biome** is a large geographical region with a distinct climate and characteristic community (e.g., rainforest, desert).
- **Habitat** is the physical space or location where an organism lives (its "address").
- **Ecological Niche** is the multidimensional functional role of a species, including resources used and conditions tolerated (its "profession").
- The **Fundamental Niche** is the full range of conditions and resources a species could theoretically use.
- The **Realized Niche** is the actual range a species occupies, often restricted by competition or other interactions.
- A **Metapopulation** is a set of geographically distinct local populations (demes) linked by immigration and emigration.
- **Symbiosis** refers to any close, long-term biological interaction between two different species (parasitism, mutualism, commensalism).
- **Abiotic components** are the non-living physical and chemical factors of an ecosystem.
- Key **physical factors** include **solar radiation, temperature, water availability, and soil texture**.
- Only about **1-2%** of incoming solar radiation is converted to chemical energy via photosynthesis.
- **Temperature** affects metabolic rates, often following a **Q₁₀ relationship** (2-3 fold increase per 10°C rise).
- Key **chemical factors** include **nutrient availability, pH, salinity, and redox potential**.
- **Liebig's Law of the Minimum** states that productivity is limited by the nutrient scarcest relative to an organism's needs.
- **Biotic components** are the living organisms, classified by their nutritional role or **trophic level**.
- **Producers (Autotrophs)** occupy the **first trophic level (T₁)**; they synthesize organic food from inorganic materials using sunlight (photosynthesis) or chemical energy (chemosynthesis).
- Examples of producers include green plants, algae, cyanobacteria, and phytoplankton.
- **Consumers (Heterotrophs)** cannot synthesize their own food and depend on producers or other consumers.
- **Primary Consumers (Herbivores - T₂)** feed directly on producers (e.g., insects, deer, zooplankton).

Practice MCQs

1. Who coined the term "ecology"?

- A) Arthur Tansley
- B) Ernst Haeckel
- C) Charles Darwin
- D) Joseph Grinnell

Answer: Ernst Haeckel

2. The term "ecosystem" was coined by:

- A) Ernst Haeckel
- B) Robert Paine
- C) Arthur Tansley
- D) Eugene Odum

Answer: Arthur Tansley

3. All the ecosystems on Earth collectively form the:

- A) Community
- B) Biome
- C) Biosphere
- D) Hydrosphere

Answer: Biosphere

4. A group of interbreeding individuals of the same species in a specific area is a:

- A) Community
- B) Population
- C) Guild
- D) Ecosystem

Answer: Population

5. The physical space where an organism lives is its:

- A) Niche
- B) Territory
- C) Habitat
- D) Biome

Answer: Habitat

6. The full range of conditions and resources a species could theoretically use defines its:

- A) Realized Niche
- B) Fundamental Niche
- C) Trophic Niche
- D) Spatial Niche

Answer: Fundamental Niche

7. A set of local populations linked by immigration and emigration is a:

- A) Community
- B) Metapopulation
- C) Species Complex
- D) Deme

Answer: Metapopulation

8. A close, long-term biological interaction between two different species is called:

- A) Competition
- B) Symbiosis
- C) Predation
- D) Commensalism

Answer: Symbiosis

9. Which of the following is an abiotic component of an ecosystem?

- A) Producer
- B) Herbivore
- C) Soil Texture
- D) Decomposer

Answer: Soil Texture

10. Liebig's Law of the Minimum states that productivity is limited by the:

- A) Most abundant resource
- B) Resource scarcest relative to needs
- C) Temperature
- D) Light availability

Answer: Resource scarcest relative to needs

11. Organisms that synthesize organic food from inorganic materials using sunlight are:

- A) Heterotrophs
- B) Decomposers
- C) Autotrophs
- D) Omnivores

Answer: Autotrophs

12. Which trophic level includes herbivores?

- A) T₁
- B) T₂
- C) T₃
- D) T₄

Answer: T₂

13. Frogs and foxes are examples of:

- A) Primary Consumers
- B) Secondary Consumers
- C) Tertiary Consumers
- D) Decomposers

Answer: Secondary Consumers

14. Organisms that break down dead organic matter, such as fungi and bacteria, are:

- A) Producers
- B) Consumers
- C) Decomposers
- D) Autotrophs

Answer: Decomposers

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Chapter 11

Viruses

- **Acellular entities** are infectious agents that **lack a cellular structure** (no cytoplasm, organelles, or metabolism). They include **viruses, viroids and prions**.
- They are **obligate intracellular parasites**, meaning they **can only replicate inside a living host cell** by hijacking its metabolic machinery.
- They are major pathogens causing significant global economic losses and health impacts in plants, animals, and bacteria.

History and Discovery of Viruses

- **Term Origin:** The word **virus** is derived from Latin for "poison" or "venom."
- **1884 (Charles Chamberland):** Found that the causative agent of rabies could pass through a porcelain filter that trapped bacteria, suggesting a filterable agent smaller than bacteria.
- **1892 (Dmitri Ivanowsky):** Demonstrated that the sap from tobacco plants with **Tobacco Mosaic Disease (TMD)** remained infectious after bacterial filtration, proposing it was caused by a filterable agent or a "bacterial toxin."
- **1898 (Martinus Beijerinck):** Conducted critical experiments showing the infectious agent in filtered sap could **replicate** only within a living host. He coined the term "**contagium vivum fluidum**" (living contagious fluid) and is credited with conceptualizing the virus.
- **1935 (Wendell Stanley):** Crystallized the **Tobacco Mosaic Virus (TMV)**, proving viruses could exist as non-living, infectious particles.
- The invention of the **electron microscope** later allowed direct visualization of viruses. The study of viruses is called **virology**.

Status of Viruses: Living or Non-Living?

Viruses exist in a gray area, exhibiting characteristics of both living and non-living entities.

Living Characteristics	Non-Living Characteristics
1. Possess genetic material (DNA or RNA) capable of mutation .	1. Acellular ; lack cellular organization, metabolic enzymes, and energy-generating systems (cannot make ATP).
2. Capable of reproduction (using host machinery).	2. Can be crystallized and stored like chemicals.
3. Show specificity and variation (exist in different strains).	3. Outside a host, they are inert , non-living particles.
4. Act as obligate intracellular parasites , causing disease.	4. Do not grow, respire, or carry out independent metabolic activities.
5. Susceptible to destruction by agents like heat and UV radiation.	5. Assemble spontaneously from pre-made components; do not divide.

- **Conclusion:** Most biologists consider viruses **non-living** because they cannot carry out life processes independently. They lead a "**borrowed life**."

Structure of Viruses

A complete, infectious viral particle is called a **virion**. It has two primary components: the **core** and the **coat**.

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

11. Viruses

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Viruses: One Liner

- **Acellular entities** lack cellular organization (no cytoplasm, organelles, or cell membrane).
- The major acellular infectious agents are **viruses, viroids, and prions**.
- Viruses are **obligate intracellular parasites**, meaning they can only replicate inside a living host cell.
- The study of viruses is called **virology**.
- The word **virus** is derived from the Latin word for "poison" or "venom".
- In 1884, **Charles Chamberland** found that the causative agent of rabies could pass through a bacteria-trapping porcelain filter.
- In 1892, **Dmitri Iwanowsky** demonstrated that the cause of **Tobacco Mosaic Disease** was a filterable agent, not a bacterium.
- In 1935, **Wendell M. Stanley** crystallized the **Tobacco Mosaic Virus (TMV)**, proving viruses could exist as non-living particles.
- The invention of the **electron microscope** allowed for the direct visualization of viruses.
- **Living characteristics of viruses** include possession of genetic material (DNA or RNA), ability to undergo mutation, reproduction using host machinery, existence in different strains, and susceptibility to destruction by agents like UV rays.
- **Non-living characteristics of viruses** include lack of cellular structure, no independent metabolism or energy generation, ability to be crystallized, and existing as inert particles outside a host.
- A complete, infectious viral particle is called a **virion**.
- The **core** contains the viral genome (DNA or RNA) and essential viral enzymes (e.g., reverse transcriptase).
- The **coat** consists of a protein shell called a **capsid**, made of repeating subunits called **capsomeres**.
- Some viruses have an additional **envelope**, a lipid bilayer derived from the host cell membrane, studded with viral **glycoprotein spikes**.
- Viral **symmetry** types are **icosahedral** (spherical, 20 triangular faces), **helical** (rod-shaped), and **complex** (a combination, e.g., bacteriophage).
- Based on **host organism**: **Bacteriophages** (infect bacteria), **Plant viruses**, and **Animal viruses**.
- The **Baltimore Classification System** categorizes viruses into 7 groups based on genome type (DNA/RNA, ss/ds) and replication strategy.
- **Group I**: Double-stranded DNA (dsDNA) viruses (e.g., Herpesvirus).
- **Group II**: Single-stranded DNA (ssDNA) viruses (e.g., Parvovirus).
- **Group III**: Double-stranded RNA (dsRNA) viruses (e.g., Rotavirus).
- **Group IV**: Positive-sense single-stranded RNA [(+)ssRNA] viruses (genome acts as mRNA) (e.g., Poliovirus, SARS-CoV-2).
- **Group V**: Negative-sense single-stranded RNA [(-)ssRNA] viruses (e.g., Influenza virus, Rabies virus).
- **Group VI**: Retroviruses (ssRNA reverse transcribing) (e.g., HIV).
- **Group VII**: Pararetroviruses (dsDNA reverse transcribing) (e.g., Hepatitis B Virus).
- Viruses are **obligate intracellular parasites** because they lack ribosomes and metabolic enzymes.
- **Host range** is determined by specific interactions between viral surface proteins (spikes) and receptor molecules on the host cell membrane.
- **Tropism** refers to a virus's specificity for certain tissues or cell types within a host (e.g., HIV infects CD4+ T-cells, poliovirus infects motor neurons).
- **Attachment/Adsorption**: Viral surface proteins bind to specific host cell receptors.

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. What is the primary structural component that surrounds the viral genome?

- A) Envelope
- B) Capsid
- C) Capsomere
- D) Nucleoid

Answer: Capsid

2. Who first crystallized a virus, proving it could exist as a non-living particle?

- A) Dmitri Ivanowsky
- B) Martinus Beijerinck
- C) Wendell Stanley
- D) Adolf Mayer

Answer: Wendell Stanley

3. Which characteristic is NOT shared by all viruses?

- A) Genetic material
- B) Protein coat
- C) Independent metabolism
- D) Obligate intracellular parasitism

Answer: Independent metabolism

4. The protein subunits that assemble to form a viral capsid are called:

- A) Protomers
- B) Capsomeres
- C) Envelopes
- D) Spikes

Answer: Capsomeres

5. A virus that infects bacteria is specifically termed a:

- A) Retrovirus
- B) Bacteriophage
- C) Viroid
- D) Prion

Answer: Bacteriophage

6. Which viral structure is derived from the host cell's membrane?

- A) Capsid
- B) Core
- C) Envelope
- D) Capsomere

Answer: Envelope

7. Tobacco mosaic virus (TMV) exhibits which type of capsid symmetry?

- A) Icosahedral
- B) Helical
- C) Complex

D) Spherical

Answer: Helical

8. The Baltimore classification system for viruses is primarily based on:

- A) Host range and disease symptoms
- B) Capsid symmetry and presence of an envelope
- C) Genome type and method of mRNA synthesis
- D) Size and morphological complexity

Answer: Genome type and method of mRNA synthesis

9. A virus with a single-stranded RNA genome that can be directly translated by host ribosomes is classified as:

- A) Negative-sense RNA virus
- B) Double-stranded RNA virus
- C) Positive-sense RNA virus
- D) Retrovirus

Answer: Positive-sense RNA virus

10. Which of the following is a DNA virus?

- A) Influenza virus
- B) HIV
- C) Adenovirus
- D) Poliovirus

Answer: Adenovirus

11. The host range of a virus is primarily determined by:

- A) The type of viral nucleic acid
- B) The presence of an envelope
- C) Specific interactions between viral surface proteins and host cell receptors
- D) The speed of viral replication

Answer: Specific interactions between viral surface proteins and host cell receptors

12. What is the first step in the viral replicative cycle?

- A) Penetration
- B) Attachment
- C) Uncoating
- D) Biosynthesis

Answer: Attachment

13. In the lytic cycle of a bacteriophage, what happens immediately after the viral DNA enters the host cell?

- A) Assembly of new virions
- B) Integration into host chromosome
- C) Degradation of host DNA



Chapter 12

Bacteria and Archaea

Prokaryotes are unicellular organisms that constitute the domains **Bacteria** and **Archaea**. They are the most abundant and widespread life forms on Earth, found in virtually every habitat. Key defining features include:

- **Lack of a membrane-bound nucleus.** Genetic material is located in a region called the **nucleoid**.
- **Absence of membrane-bound organelles** (e.g., mitochondria, chloroplasts).
- Generally smaller (0.5–5.0 μm) and structurally simpler than eukaryotic cells.
- Possess a rigid **cell wall** (with exceptions like *Mycoplasma*).
- Exhibit extraordinary **metabolic diversity**, enabling survival in extreme environments.

Despite their simplicity, they are highly evolved, having existed for over 3.5 billion years. While a minority are pathogens, most play crucial ecological roles in nutrient cycling, decomposition, and symbiotic relationships.

Historical Discovery and Foundations

- **Antonie van Leeuwenhoek (1673):** First to observe and describe bacteria ("animalcules") using a simple microscope.
- **Louis Pasteur (19th Century):**
 - Disproved spontaneous generation.
 - Developed **germ theory of disease**.
 - Created vaccines (anthrax, rabies).
 - Developed the **pasteurization** process.
- **Robert Koch (19th Century):** Established the definitive link between specific microbes and specific diseases using **Koch's Postulates**.
 1. The pathogen must be present in all diseased individuals and absent in healthy ones.
 2. The pathogen must be isolated and grown in pure culture.
 3. The cultured pathogen must cause the same disease when inoculated into a healthy host.
 4. The pathogen must be re-isolated from the experimentally infected host.

Taxonomy and Phylogenetic Position

- **Historical Classification:** All prokaryotes were initially placed in the kingdom **Monera**.
- **The Three-Domain System (Carl Woese, 1977):** Based on comparisons of **small-subunit ribosomal RNA (rRNA)** gene sequences, life is divided into three domains:
 1. **Bacteria (Eubacteria):** "True bacteria."
 2. **Archaea (Archaeobacteria):** A distinct lineage sharing some features with eukaryotes.
 3. **Eukarya:** All eukaryotic organisms.
- This system revealed that **Archaea are more closely related to Eukarya than to Bacteria**. The last common ancestor of Bacteria and Archaea was likely a **hyperthermophile**.

Structure of Bacteria

Size and Shape

- **Size:** Typically 0.5–5.0 μm. Exceptions include *Mycoplasma* (~0.1 μm) and the giant *Epulopiscium fishelsoni* (up to 600 μm).
- **Shapes:** Used for primary classification.
 - **Cocci (spherical):** Arrangements include diplococci (pairs), streptococci (chains), staphylococci (clusters), tetrads (4), and sarcinae (8).
 - **Bacilli (rod-shaped):** Can be single, diplobacilli (pairs), or streptobacilli (chains). Vibrios are comma-shaped rods.
 - **Spiral:** Includes **spirilla** (rigid, helical) and **spirochetes** (flexible, tightly coiled).
- **Pleomorphic:** Some bacteria (e.g., *Mycoplasma*) lack a constant shape.



Bacteria & Archaea: One Liner

- **Prokaryotes** are unicellular organisms that lack a membrane-bound nucleus and membrane-bound organelles.
- The **five-kingdom system** (Robert Whittaker, 1969) placed prokaryotes in the kingdom **Monera**.
- The **three-domain system** (Carl Woese, 1977) classifies life into **Bacteria, Archaea,** and **Eukarya**.
- **Phylogenetics** is the study of evolutionary relatedness among groups of organisms.
- **Bacteria** and **Archaea** are the two distinct **domains** of prokaryotic life.
- The most recent common ancestor of bacteria and archaea is estimated to have lived **2.5–3.2 billion years ago**.
- Prokaryotes were the **first life forms** on Earth, appearing approximately **4 billion years ago**.
- **Archaea** were originally called **archaebacteria** and are known for inhabiting extreme environments.
- **Methanogens** are archaea that produce **methane** as a metabolic by-product in anaerobic conditions.
- **Halophiles** are salt-loving archaea that thrive in highly saline environments.
- **Thermoacidophiles** are archaea that live in extremely **hot and acidic** environments, like hot springs.
- Archaeal cell walls **lack peptidoglycan**.
- Archaeal membrane lipids have **branched hydrocarbon chains** linked by **ether** bonds, unlike bacterial ester-linked unbranched chains.
- Archaeal **RNA polymerase** is complex and similar to eukaryotic RNA polymerase.
- Some archaeal genes contain **introns**, which are rare in bacteria.
- *Thermoplasma acidophilum* and *Thermus aquaticus* are examples of **thermoacidophiles**.
- *Halobacterium halobium* is an example of an extreme **halophile**.
- The study of bacteria is called **bacteriology**, a branch of microbiology.
- **Anton van Leeuwenhoek** first observed bacteria in 1674, calling them "animalcules".
- **Christian Gottfried Ehrenberg** introduced the name "bacterium" in 1828.
- Bacteria inhabit the **widest range of habitats** on Earth, from soil to human intestines to extreme environments.
- **Flagella** provide motility, allowing bacteria to move toward favorable conditions (**taxis**).
- **Pili** (or fimbriae) are hair-like surface structures that aid in **adhesion** to surfaces and other cells.
- **Endospores** are highly resistant, dormant structures formed by some bacteria to survive harsh conditions.
- **Actinomycetes** are a group of bacteria that form branching filaments, common in soil for decomposing organic matter.
- **Aquificales** are extremely **thermophilic** and represent the oldest branch of bacteria (e.g., *Aquifex*).
- **Green Non-Sulfur Bacteria** are mostly photosynthetic and some are thermophilic (e.g., *Chloroflexus*).
- **Green Sulfur Bacteria** are green phototrophic bacteria that do **not produce oxygen** during photosynthesis (e.g., *Chlorobium*).
- **Proteobacteria (Purple Bacteria)** are a large, diverse group of **Gram-negative** bacteria (e.g., *Escherichia, Salmonella*).
- **Gram-positive Bacteria** include endospore-formers, lactic acid bacteria, and pathogens (e.g., *Staphylococcus, Bacillus*).
- **Cyanobacteria** (blue-green algae) are **oxygen-producing** photosynthetic bacteria (e.g., *Nostoc, Anabaena*).
- **Bacteroides** are **Gram-negative, rod-shaped** bacteria (e.g., *Bacteroides*).

- **Antibiotic Sensitivity:** Bacteria are often inhibited by streptomycin/chloramphenicol; Archaea are not.
- **Initiator Amino Acid:** Bacteria use **formyl-methionine**; Archaea use **methionine** (like eukaryotes).

Practice MCQs

1. Which scientist first observed bacteria using a homemade microscope?

- A) Louis Pasteur
- B) Robert Koch
- C) Antonie van Leeuwenhoek
- D) Christian Gram

Answer: Antonie van Leeuwenhoek

2. The domain Archaea differs from Bacteria in having:

- A) Peptidoglycan in cell wall
- B) Ester-linked membrane lipids
- C) Ether-linked membrane lipids
- D) Circular chromosome

Answer: Ether-linked membrane lipids

3. Gram-positive bacteria stain purple due to:

- A) Thin peptidoglycan layer
- B) Outer lipopolysaccharide membrane
- C) Thick peptidoglycan layer retaining crystal violet
- D) Presence of periplasmic space

Answer: Thick peptidoglycan layer retaining crystal violet

4. Which of the following is a photosynthetic prokaryote?

- A) Escherichia coli
- B) Streptococcus pneumoniae
- C) Nostoc
- D) Staphylococcus aureus

Answer: Nostoc

5. Koch's postulates are used to:

- A) Classify bacteria based on shape
- B) Prove causal relationship between microbe and disease
- C) Identify antibiotic resistance
- D) Stain bacterial capsules

Answer: Prove causal relationship between microbe and disease

6. Bacteria that require oxygen for growth are called:

- A) Obligate anaerobes
- B) Facultative anaerobes
- C) Obligate aerobes
- D) Microaerophiles

Answer: Obligate aerobes

7. The rigid spiral-shaped bacteria are known as:

- A) Vibrio
- B) Spirillum
- C) Spirochete
- D) Coccus

Answer: Spirillum

8. Which structure is involved in bacterial conjugation?

- A) Flagella
- B) Pili
- C) Capsule
- D) Endospore

Answer: Pili

9. The process of taking up free DNA from the environment is:

- A) Transduction
- B) Conjugation
- C) Transformation
- D) Binary fission

Answer: Transformation

10. Which of the following bacteria produce endospores?

- A) Escherichia coli
- B) Bacillus anthracis
- C) Neisseria gonorrhoeae
- D) Vibrio cholerae

Answer: Bacillus anthracis

11. Plasmids are best described as:

- A) Essential for bacterial growth
- B) Circular, double-stranded extrachromosomal DNA
- C) Part of the bacterial chromosome
- D) Involved in protein synthesis

Answer: Circular, double-stranded extrachromosomal DNA

12. Cyanobacteria perform oxygenic photosynthesis due to presence of:

- A) Bacteriochlorophyll
- B) Chlorophyll a and photosystem II
- C) Chloroplasts
- D) Chromatophores

Answer: Chlorophyll a and photosystem II

M
K
P
R
E
P
A
R
A
T
I
O
N
S

12. Bacteria and Archaea

Chapter 13

Protists

Protists are a diverse, informal assemblage of **primarily aquatic eukaryotic organisms** that are **not classified as true fungi, plants, or animals**. The term is derived from the Greek for "the very first," reflecting their status as **among the first eukaryotes to evolve**.

They represent a **paraphyletic group**, meaning the group includes some but not all descendants of a common ancestor. In modern taxonomy, the formal **Kingdom Protista** has been largely abandoned because molecular phylogenetics shows various protist lineages are more closely related to plants, animals, or fungi than to each other. The term "protist" is now used informally for **eukaryotes that are not plants, animals, or fungi**.

Reasons for Creation of Kingdom Protista

The Kingdom Protista was established to resolve classification challenges posed by certain eukaryotes.

- **i) Extreme Diversity:** Protists exhibit vast variation in body forms (unicellular, colonial, multicellular), reproduction (asexual, sexual), nutrition (autotrophic, heterotrophic, mixotrophic), and habitats, making them hard to characterize under traditional plant or animal kingdoms.
- **ii) Definition by Exclusion:** The kingdom is largely defined by what its members are **not**—they lack the specialized traits that place organisms definitively in Plantae, Animalia, Fungi, or Monera.
- **iii) Taxonomic Difficulty:** Early taxonomists struggled to classify eukaryotes like *Euglena* (which has both plant-like chloroplasts and animal-like motility) into existing kingdoms.
- **iv) Evolutionary Origin:** Protista is considered the ancestral eukaryotic group from which the other three eukaryotic kingdoms (Plantae, Animalia, Fungi) independently evolved.
- **v) Polyphyly:** Modern molecular data indicates protists do not share a single common ancestor; the kingdom is a **polyphyletic** assemblage of multiple lineages.

General Characteristics of Protists

Protists exhibit extraordinary **structural, functional, and ecological diversity**.

Cellular Organization

- All are **eukaryotic**, possessing a **membrane-bound nucleus** and other membrane-bound organelles (mitochondria, Golgi apparatus, etc.).
- They have a **well-developed cytoskeleton** enabling complex shapes, motility, and phagocytosis.

Body Plan Diversity

- **Unicellular:** Most common form (e.g., *Amoeba*, *Paramecium*). Can be extremely complex, performing all life functions within a single cell.
- **Colonial:** Loosely organized groups of cells, often embedded in a matrix (e.g., *Volvox*).
- **Coenocytic:** A multinucleate mass of cytoplasm without separate cell walls (e.g., plasmodial slime molds, some water molds like Oomycetes).
- **Simple Multicellular:** Have differentiated cells but lack complex tissues (e.g., brown algae, red algae). The body is a **thallus** (not differentiated into roots, stems, leaves).

Locomotion

Motile at some life stage via:

- **Pseudopodia:** Cytoplasmic extensions used for amoeboid movement and feeding (e.g., amoebas).
- **Flagella:** Long, whip-like structures with a **9+2 microtubule arrangement**.
- **Cilia:** Short, hair-like structures, similar to flagella but more numerous (e.g., ciliates).
- **Gliding:** Movement on secreted slime (e.g., diatoms, apicomplexans).

Nutritional Modes

Exhibit all major types:

- **Photoautotrophs:** Photosynthetic with chloroplasts (e.g., algae).

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091



Kingdom Protista: One Liner

- **Kingdom Protista** is a diverse assemblage of primarily **aquatic eukaryotic organisms** that do not fit into the kingdoms Fungi, Plantae, or Animalia.
- This kingdom is defined largely by **exclusion**.
- It is considered a **polyphyletic group**, meaning its members do not share a single common ancestor.
- Protists are among the **first eukaryotes** to evolve.
- The traditional kingdom **Protista** has been abandoned in modern systematics, though the term "protist" remains useful informally.
- Protists show immense variation in **body form, reproduction, nutrition, and lifestyle**, making them hard to classify.
- The kingdom is defined negatively; its members possess characteristics that exclude them from the other four eukaryotic kingdoms.
- Scientists historically found it difficult to place certain eukaryotes like *Euglena* into the plant or animal kingdom.
- The other eukaryotic kingdoms (**Plantae, Animalia, and Fungi**) **originated from various protistan ancestors**.
- This kingdom serves as the foundational group from which plants, animals, and fungi arose.
- Most protists are **aquatic**, living in marine and freshwater habitats.
- All protists are **eukaryotes**, with cells containing a nucleus and membrane-bound organelles.
- Their body organization can be **unicellular, colonial, or simple multicellular**.
- They **do not develop from a blastula or embryo** like animals or plants.
- They exhibit diverse modes of nutrition: **autotrophic (photosynthetic), heterotrophic (ingestive or absorptive), and mixotrophic**.
- They are motile at some life stage, using **pseudopodia, cilia, flagella, or gliding**.
- They reproduce both **asexually** (binary fission, budding, multiple fission) and **sexually**.
- Their life cycles can be **haplontic, diplontic, or haplodiplontic (alternation of generations)**.
- They exhibit an extreme size range, from microscopic to over 60 meters in length (e.g., kelps).
- In **1861, John Hogg** proposed the **kingdom Protoctista** for microscopic organisms.
- In **1866, Ernst Haeckel** proposed the **kingdom Protista**, including bacteria and microorganisms like *Euglena*.
- In **1938, Herbert Copeland** elevated prokaryotes to **kingdom Monera**, separating them from protists.
- In **1969, Robert Whittaker** formally included unicellular eukaryotes in **Kingdom Protista** as part of his **Five-Kingdom Classification**.
- In **1982, Margulis and Schwartz** modified the five-kingdom system, retaining Protista/Protoctista and including colonial and simple multicellular eukaryotes.
- Protists exhibit remarkable diversity in **size, habitat, structure, locomotion, reproduction, nutrition, and ecological interactions**.
- Due to their vast diversity and polyphyletic origins, Margulis and Schwartz listed **27 phyla** within Kingdom Protista.
- **Endosymbiosis** is a key process in the evolution of eukaryotic cells.
- **Mitochondria** are derived from an **alpha proteobacterium** engulfed by an ancestral archaeal (or archaea-related) host cell.
- **Plastids** are derived from a **cyanobacterium** engulfed by a heterotrophic eukaryote via **primary endosymbiosis**.
- **Secondary endosymbiosis** occurred when a heterotrophic eukaryote engulfed a photosynthetic eukaryote (a red or green alga), leading to plastids surrounded by three or four membranes.

M
K
P
R
E
P
A
R
A
T
I
O
N
S

13. Protists

Practice MCQs

1. What is the estimated time frame for the appearance of the first eukaryotic fossils?

- A) 3.5 billion years ago
- B) 2.5 billion years ago
- C) 1.5 billion years ago
- D) 500 million years ago

Answer: 1.5 billion years ago

2. Which structure is believed to have evolved from infoldings of the plasma membrane in prokaryotic ancestors?

- A) Mitochondria
- B) Chloroplasts
- C) Nucleus and endoplasmic reticulum
- D) Golgi apparatus

Answer: Nucleus and endoplasmic reticulum

3. According to the endosymbiotic theory, what is the proposed origin of mitochondria?

- A) Engulfed cyanobacterium
- B) Engulfed aerobic alpha-proteobacterium
- C) Infoldings of the plasma membrane
- D) Vesicles from the Golgi apparatus

Answer: Engulfed aerobic alpha-proteobacterium

4. Which evidence supports the endosymbiotic origin of chloroplasts?

- A) Presence of 70S ribosomes and circular DNA
- B) Presence of a double membrane and histone proteins
- C) Ability to undergo mitosis independently
- D) Synthesis of cellulose cell walls

Answer: Presence of 70S ribosomes and circular DNA

5. Which of the following is a characteristic feature of all protists?

- A) They are all unicellular
- B) They are all photosynthetic
- C) They are all eukaryotic
- D) They all have cell walls

Answer: They are all eukaryotic

6. What is the primary locomotory structure in amoebas?

- A) Flagella
- B) Cilia
- C) Pseudopodia
- D) Undulating membrane

Answer: Pseudopodia

7. Which term describes protists that can switch between autotrophic and heterotrophic

nutrition?

- A) Saprophytes
- B) Mixotrophs
- C) Phototrophs
- D) Osmotrophs

Answer: Mixotrophs

8. What is the process called when a protist undergoes multiple nuclear divisions before cytokinesis?

- A) Binary fission
- B) Budding
- C) Schizogony
- D) Conjugation

Answer: Schizogony

9. Which excavate group is characterized by having two nuclei and lacks functional mitochondria?

- A) Parabasalids
- B) Euglenozoans
- C) Diplomonads
- D) Kinetoplastids

Answer: Diplomonads

10. What human disease is caused by the parabasalid *Trichomonas vaginalis*?

- A) Malaria
- B) Trichomoniasis
- C) Giardiasis
- D) Sleeping sickness

Answer: Trichomoniasis

11. What unique structure is found in the flagella of euglenozoans?

- A) Mastigonemes
- B) Crystalline rod
- C) Axoneme
- D) Basal body

Answer: Crystalline rod

12. Which kinetoplastid causes African sleeping sickness?

- A) *Trypanosoma cruzi*
- B) *Trypanosoma brucei*
- C) *Leishmania donovani*
- D) *Giardia intestinalis*

Answer: *Trypanosoma brucei*

13. What characteristic feature is found on the flagella of stramenopiles?

- A) Alveoli
- B) Mastigonemes (hairy projections)
- C) Crystalline rods

M
K

P
R
E
P
A
R
A
T
I
O
N
S

13. Protists



Chapter 14

Fungi

Mycology is the scientific study of fungi. This kingdom comprises a vast diversity of eukaryotic, heterotrophic organisms, including mushrooms, molds, yeasts, and lichen partners. Fungi are primarily terrestrial and play indispensable ecological roles as decomposers, mutualists, and pathogens. While macroscopic structures like mushrooms are familiar, they represent only the reproductive parts; the main vegetative body is typically a subterranean, extensive network called a **mycelium**.

Key Estimates & Significance:

- **Described species:** ~100,000 – 145,000.
- **Estimated total species:** 1.5 – 5 million (suggesting massive undiscovered diversity).
- **Evolutionary origin:** Appeared ~1 billion years ago; major diversification occurred ~500 million years ago alongside land plants.
- **Evolutionary relationship:** Molecular and structural evidence places fungi in the **Opisthokont** clade, making them the **closest relatives of animals** (shared ancestor ~460 MYA), not plants.
- **Ecological dominance:** A single clone of the fungus *Armillaria ostoyae* (the "Humongous Fungus") in Oregon covers over **2,400 acres**, weighs ~35,000 tons, and is estimated to be **2,400 years old**, making it one of Earth's largest and oldest individual organisms.

GENERAL CHARACTERISTICS OF FUNGI

1. Nutritional Mode

Fungi are **chemoheterotrophs** with a unique **absorptive** strategy:

- **Secrete powerful exoenzymes** (e.g., cellulases, lignin peroxidases, proteases) into their surroundings.
- **Externally digest** complex organic matter (living or dead) into simpler, soluble molecules.
- **Absorb nutrients** directly across their cell membranes via active transport.

2. Cellular & Structural Features

- **Cell Wall:** Composed of **chitin** (β -1,4-linked N-acetylglucosamine), providing rigidity and resistance to degradation. This differs fundamentally from plant cellulose.
- **Storage Polysaccharide: Glycogen**, identical to animal glycogen, stored in cytoplasmic granules.
- **Mitochondria:** Have **flattened, plate-like cristae** (tubular in some early-diverging groups).
- **Nuclear Division (Mitosis):** Often "**closed mitosis**"—the nuclear envelope persists, and the mitotic spindle forms inside the nucleus. Centrioles are absent in most groups; spindle poles are organized by **spindle pole bodies**.
- **Habitat Tolerance:** Prefer slightly acidic pH (~5.6) but tolerate a wide range (pH 2–9). Many are **osmotolerant**, thriving in high-sugar or high-salt environments (e.g., jam, salted meats).

3. Body Plan (Thallus Organization)

Fungi exhibit two fundamental growth forms:

Form	Description	Example
Unicellular Yeasts	Spherical or oval cells. Reproduce asexually by budding or fission . Common in moist, nutrient-rich fluids.	<i>Saccharomyces cerevisiae</i> (Baker's yeast)

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Kingdom Fungi: One Liner

- **Mycology** is the scientific study of fungi.
- Fungi are a kingdom of **eukaryotic, heterotrophic** organisms.
- Their primary mode of nutrition is **absorptive heterotrophy**.
- They secrete digestive **exoenzymes** to break down complex matter externally and absorb the nutrients.
- The optimal pH for most fungal growth is about **5.6**, but they can tolerate a wide range from **pH 2 to 9**.
- Fungi are largely **osmotolerant**, able to grow in high-sugar or high-salt environments.
- Fungi are classified within the **Opisthokont** clade, making them the **closest relatives of animals**, not plants.
- This close relationship makes treating fungal infections challenging, as drugs targeting fungi can also harm the host.
- They first appeared approximately **1 billion years ago**, with major diversification around **500 million years ago**.
- Mycologists have described about **100,000 to 145,000** fungal species.
- The total estimated number of fungal species is between **1.5 and 5 million**.
- A single clone of the fungus *Armillaria ostoyae* in Oregon covers over **2,400 acres**, weighs ~35,000 tons, and is estimated to be **2,400 years old**.
- The fungal cell wall is primarily composed of **chitin**, a polymer of **N-acetylglucosamine**.
- **Chitin** is also a key component of insect and arthropod exoskeletons.
- Fungi store carbohydrates as **glycogen**, similar to animals, not as starch.
- Fungal mitochondria typically have **flattened, plate-like cristae**.
- Fungal mitosis is often "**closed mitosis**", where the nuclear envelope persists.
- The vegetative body of a fungus is called a **thallus**.
- The basic structural unit of multicellular fungi is a **hypha** (pl. hyphae).
- A mass of interconnected hyphae is called a **mycelium**.
- The mycelial structure provides an **extremely high surface-area-to-volume ratio** for efficient absorption.
- Hyphae grow at their tips, a process known as **apical growth**.
- **Coenocytic (aseptate) hyphae** lack cross-walls and contain multiple nuclei in a continuous cytoplasm.
- **Septate hyphae** are divided by cross-walls called **septa**.
- Septa often contain **pores** that allow cytoplasm and organelles to flow between cells.
- **Yeasts** are unicellular fungi with a round or oval shape.
- Yeasts reproduce asexually by **budding** or **fission**.
- **Dimorphic fungi** can switch between yeast and hyphal forms depending on environmental conditions.
- **Rhizoids** are root-like anchoring hyphae (e.g., in *Rhizopus*).
- **Haustoria** are specialized hyphal tips of parasitic fungi that penetrate host cell walls to absorb nutrients.
- **Appressoria** are swollen, adhesive hyphal tips that generate pressure to physically penetrate host surfaces.
- **Arbuscules** are highly branched, tree-like structures formed inside plant root cells by **arbuscular mycorrhizal fungi** for nutrient exchange.
- **Trapping structures** like constricting rings or adhesive nets are formed by predatory fungi to capture nematodes.
- **Sclerotia** are compact, hardened masses of hyphae that serve as dormant, resistant structures.

Practice MCQs

M
K

P
R
E
P
A
R
A
T
I
O
N
S

1. What is the primary structural component of fungal cell walls?

- A) Cellulose
- B) Chitin
- C) Peptidoglycan
- D) Lignin

Answer: Chitin

2. Fungi are more closely related to which of the following groups?

- A) Plants
- B) Animals
- C) Bacteria
- D) Archaea

Answer: Animals

3. Which of the following describes the nutritional mode of fungi?

- A) Autotrophic
- B) Absorptive heterotrophs
- C) Ingestive heterotrophs
- D) Photosynthetic

Answer: Absorptive heterotrophs

4. The vegetative body of a multicellular fungus is called a:

- A) Thallus
- B) Mycelium
- C) Hypha
- D) Spore

Answer: Mycelium

5. Individual fungal filaments are known as:

- A) Septa
- B) Hyphae
- C) Mycelia
- D) Rhizoids

Answer: Hyphae

6. Which fungal phylum is characterized by coenocytic hyphae and zygospores?

- A) Ascomycota
- B) Basidiomycota
- C) Zygomycota
- D) Chytridiomycota

Answer: Zygomycota

7. In which phylum are sac-like asci that contain ascospores found?

- A) Zygomycota
- B) Glomeromycota
- C) Ascomycota
- D) Basidiomycota

Answer: Ascomycota

8. Club fungi, which produce basidiospores on basidia, belong to:

- A) Ascomycota
- B) Basidiomycota
- C) Zygomycota
- D) Deuteromycota

Answer: Basidiomycota

9. Which fungal phylum forms arbuscular mycorrhizae?

- A) Glomeromycota
- B) Zygomycota
- C) Ascomycota
- D) Chytridiomycota

Answer: Glomeromycota

10. Fungi that lack a known sexual stage were historically grouped as:

- A) Zygomycetes
- B) Deuteromycetes
- C) Basidiomycetes
- D) Ascomycetes

Answer: Deuteromycetes

11. What is the primary storage carbohydrate in fungi?

- A) Starch
- B) Glycogen
- C) Cellulose
- D) Chitin

Answer: Glycogen

12. The fusion of cytoplasm from two parent mycelia is called:

- A) Karyogamy
- B) Plasmogamy
- C) Meiosis
- D) Mitosis

Answer: Plasmogamy

13. The fusion of nuclei in fungal sexual reproduction is termed:

- A) Plasmogamy
- B) Karyogamy
- C) Heterokaryosis
- D) Dikaryosis

Answer: Karyogamy

14. A fungal mycelium containing two genetically distinct haploid nuclei per cell is:

- A) Heterokaryotic
- B) Dikaryotic
- C) Diploid



Chapter 15

Kingdom Plantae

Kingdom Plantae (Embryophytes) represents multicellular, photosynthetic eukaryotes that have successfully colonized terrestrial environments. Modern classification follows a **phylogenetic system**, reflecting evolutionary relationships rather than mere morphological similarity. Plants evolved from **freshwater charophyte green algae** approximately 500 million years ago.

M
K
P
R
E
P
A
R
A
T
I
O
N
S

General Diagnostic Features Of Land Plants

- **Eukaryotic & Multicellular:** Composed of complex cells with membrane-bound organelles and true nuclei.
- **Autotrophic Nutrition:** Perform **photosynthesis** using chlorophylls *a* and *b* within chloroplasts. Storage product is **starch**.
- **Cell Wall:** Structural support provided by walls composed primarily of **cellulose**.
- **Cuticle:** A protective, **lipophilic, waxy layer** (containing cutin and waxes) covering aerial epidermal surfaces to minimize water loss.
- **Stomata (Sing. Stoma): Regulatable pores** flanked by guard cells that allow for gas exchange (CO₂ intake for photosynthesis, O₂ release) while controlling **transpirational water loss**.
- **Gametangia:** Multicellular organs that produce and protect gametes.
 - **Antheridium:** Produces numerous **flagellated sperm cells**.
 - **Archegonium:** A flask-shaped structure producing a single, non-motile **egg cell**.
- **Embryo Retention (Embryophytic Condition):** The defining plant trait. The **zygote develops into a multicellular diploid embryo** retained within and nourished by the maternal gametophyte tissue.
- **Alternation of Generations:** A life cycle with **multicellular haploid (gametophyte)** and **multicellular diploid (sporophyte)** phases.
 - **Gametophyte (n):** Develops from a spore. Produces **gametes via mitosis**.
 - **Sporophyte (2n):** Develops from the zygote. Produces **haploid spores via meiosis in sporangia**.
 - **Evolutionary Trend:** A progressive **reduction of the gametophyte** and **elaboration of the sporophyte**, correlating with increasing terrestriality.
- **Apical Meristems:** Localized regions of **perpetual embryonic cells** at root and shoot tips, enabling **primary growth** (extension) and complex tissue formation.
- **Secondary Metabolites:** Synthesis of diverse compounds (e.g., alkaloids, terpenes, phenolics) for defense against herbivores, pathogens, and UV radiation.

Classification of Kingdom Plantae

I. Non-Vascular Plants (Bryophytes)

- **Division: Bryophyta**
 - **Class: Hepaticae** (Liverworts) - e.g., *Marchantia*
 - **Class: Anthocerotae** (Hornworts) - e.g., *Anthoceros*
 - **Class: Musci** (Mosses) - e.g., *Funaria*

II. Vascular Plants (Tracheophyta)

A. Seedless Vascular Plants (Cryptogams)

- **Division: Pteridophyta** (Ferns & Allies)
 - **Class: Psilopsida** (Whisk ferns) - e.g., *Psilotum*
 - **Class: Lycopsidea** (Club mosses) - e.g., *Lycopodium, Selaginella*
 - **Class: Sphenopsida** (Horsetails) - e.g., *Equisetum*
 - **Class: Pteropsida** (True Ferns) - e.g., *Pteris, Adiantum*

B. Seed-Bearing Plants (Phanerogams/Spermatophytes)

- **Division: Gymnospermae** (Naked Seeds)



Kingdom Plantae: One Liner

- Modern plant classification follows a **phylogenetic system** based on evolutionary relationships.
- Kingdom Plantae (**Embryophytes**) are multicellular, photosynthetic eukaryotes that have colonized land.
- Plants evolved from **freshwater charophyte green algae**, specifically within the **Zygnematophyceae** (or Charophyceae), around 500 million years ago.
- Shared ancestry with charophytes is evidenced by biochemical similarities (chlorophyll *a* & *b*, cellulose, starch), cell division patterns (phragmoplast), and genetic homology.
- The transition to land presented challenges like **desiccation, UV radiation, gravity, and reproduction without water**.
- Primary divisions are **non-vascular plants (Bryophytes)** and **vascular plants (Tracheophytes)**.
- Tracheophytes comprise **Lycophytes** and **Euphyllophytes** (ferns, horsetails, and seed plants).
- Plants are **eukaryotic, multicellular, and autotrophic**, performing **photosynthesis** with chlorophylls *a* and *b*.
- The primary storage product is **starch**.
- The cell wall is composed mainly of **cellulose**.
- A protective, waxy **cuticle** (containing cutin) covers aerial surfaces to minimize water loss.
- **Stomata** are regulatable pores flanked by guard cells that allow gas exchange (CO₂ in, O₂ out) while controlling transpiration.
- Gametes are produced within multicellular organs called **gametangia** (**antheridia** produce sperm; **archegonia** produce eggs).
- They exhibit the **embryophytic condition**: the zygote develops into a multicellular embryo retained within and nourished by the maternal gametophyte.
- Their life cycle involves a **heteromorphic alternation of generations** between a multicellular haploid **gametophyte** and a multicellular diploid **sporophyte**.
- The evolutionary trend is a **reduction of the gametophyte and dominance of the sporophyte** with increasing terrestriality.
- **Apical meristems** at root and shoot tips enable primary growth.
- They produce **secondary metabolites** (e.g., alkaloids, terpenes) for defense.
- Bryophytes are a **paraphyletic group** representing the earliest land plant lineages.
- They **lack true vascular tissue (xylem/phloem) and true roots, stems, and leaves** (though they have analogous structures).
- They are **poikilohydric** (water content varies with the environment) and are **habitat-restricted** to moist areas.
- They require water for **sperm motility**.
- The **gametophyte generation is dominant, persistent, and photosynthetic**.
- The sporophyte is **dependent** on the gametophyte and is typically **homosporous**.
- Bryophytes are divided into three phyla: **Marchantiophyta (Liverworts)**, **Bryophyta (True Mosses)**, and **Anthocerotophyta (Hornworts)**.
- **Liverworts (Marchantiophyta)**: Gametophyte can be **thalloid** (e.g., *Marchantia*) or **leafy** (e.g., *Porella*). They reproduce asexually via **gemmae cups**. The sporophyte is ephemeral. They generally **lack true stomata** (some have pores) and possess **oil bodies** in cells.
- **Mosses (Bryophyta)**: Gametophyte has erect, "leafy" shoots with **multicellular rhizoids**. Sporophyte is differentiated into **foot, seta, and capsule**. **Stomata are present on the sporophyte capsule**. The spore germinates into a filamentous **protonema**. Examples: *Funaria*, *Polytrichum*, *Sphagnum*.
- **Hornworts (Anthocerotophyta)**: Gametophyte is simple and thalloid. Cells contain a **single large, plate-like chloroplast**. The sporophyte is an **elongated, photosynthetic capsule** that

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. Which modern system of classification reflects evolutionary relationships among plants?

- A) Artificial system
- B) Natural system
- C) Phylogenetic system
- D) Linnaean system

Answer: Phylogenetic system

2. Plants are believed to have evolved from which group of algae?

- A) Rhodophyceae
- B) Phaeophyceae
- C) Charophyceae (Zygnematophyceae)
- D) Chlorophyceae

Answer: Charophyceae (Zygnematophyceae)

3. What is the primary storage product in plants?

- A) Glycogen
- B) Cellulose
- C) Starch
- D) Lipids

Answer: Starch

4. Which structure minimizes water loss in land plants by covering aerial epidermal surfaces?

- A) Stomata
- B) Cuticle
- C) Lenticels
- D) Hydathodes

Answer: Cuticle

5. Regulatable pores flanked by guard cells that facilitate gas exchange are called?

- A) Hydathodes
- B) Lenticels
- C) Stomata
- D) Trichomes

Answer: Stomata

6. The multicellular organ that produces and protects the egg in plants is the?

- A) Antheridium
- B) Archegonium
- C) Sporangium
- D) Gametangium

Answer: Archegonium

7. The defining trait of embryophytes is?

- A) Presence of vascular tissue
- B) Retention of the multicellular embryo
- C) Dominant sporophyte generation

D) Production of seeds

Answer: Retention of the multicellular embryo

8. In the alternation of generations, the haploid phase that produces gametes is the?

- A) Sporophyte
- B) Zygote
- C) Gametophyte
- D) Embryo

Answer: Gametophyte

9. The evolutionary trend in land plants shows a progressive reduction of the?

- A) Sporophyte
- B) Gametophyte
- C) Zygote
- D) Embryo

Answer: Gametophyte

10. Localized regions of perpetual embryonic cells at root and shoot tips are?

- A) Vascular cambium
- B) Cork cambium
- C) Apical meristems
- D) Intercalary meristems

Answer: Apical meristems

11. Bryophytes lack which of the following?

- A) Cuticle
- B) Stomata
- C) True vascular tissue
- D) Chlorophyll

Answer: True vascular tissue

12. The dominant generation in bryophytes is the?

- A) Sporophyte
- B) Gametophyte
- C) Zygote
- D) Embryo

Answer: Gametophyte

13. Which bryophyte phylum has a thalloid gametophyte and reproduces asexually via gemmae cups?

- A) Bryophyta (Mosses)
- B) Anthocerotophyta (Hornworts)
- C) Marchantiophyta (Liverworts)
- D) Lycopodiophyta

Answer: Marchantiophyta (Liverworts)

14. In mosses, the spore germinates into a filamentous stage called the?

- A) Prothallus



Chapter 16

Form and Functions in Plants

Plant Nutrition

Nutrition: The process by which organisms obtain and utilize nutrients for growth, metabolism, and maintenance. Plants are **autotrophs** (specifically photoautotrophs), synthesizing organic compounds from inorganic materials using light energy.

Essential Plant Nutrients

Sixteen elements are considered essential (Arnon & Stout, 1939). Essentiality criteria:

1. Required for normal life cycle completion.
2. Function is specific and cannot be replaced.
3. Directly involved in metabolism.

Classification by Quantity & Mobility:

- **Macronutrients** (> 0.05% dry weight): Required in large amounts.
- **Micronutrients/Trace Elements** (< 0.05% dry weight): Required in minute amounts.
- **Mobile Nutrients:** Symptoms appear first in older leaves (e.g., N, P, K, Mg).
- **Immobile Nutrients:** Symptoms appear first in young leaves/meristems (e.g., Ca, Fe, B).

NUTRIENT FUNCTIONS & DEFICIENCIES

Nutrient (Absorbed Form)	Category	Major Physiological & Biochemical Functions	Specific Deficiency Symptoms	Mobility
Nitrogen (NO ₃ ⁻ , NH ₄ ⁺)	Macro	- Component of amino acids, proteins, nucleic acids, chlorophyll, ATP, coenzymes. - Central to metabolism, growth, and photosynthesis.	General chlorosis (yellowing) starting in older leaves ; stunted growth; thin stems; delayed flowering & fruiting.	Mobile
Phosphorus (H ₂ PO ₄ ⁻ /HPO ₄ ²⁻)	Macro	- Key component of ATP, nucleic acids (DNA/RNA), phospholipids (membranes), coenzymes (NADP⁺). - Energy transfer, genetic information, membrane integrity.	Dark green/blue-green leaves with possible purple/red anthocyanin pigmentation (sugar accumulation); poor root development; delayed maturity.	Mobile
Potassium (K ⁺)	Macro	- Enzyme activation (over 50 enzymes, e.g., for protein synthesis, starch formation). - Osmotic regulation & turgor (stomatal opening). - Solute transport, charge balance.	Chlorosis & necrosis (dead tissue) at leaf margins and tips ; "scorched" appearance; weak stalks (lodging); poor resistance to stress.	Mobile
Calcium (Ca ²⁺)	Macro	- Structural: Component of middle lamella (calcium pectate). - Signaling: Second messenger in hormone & stress responses.	Death of apical meristems (root & shoot tips); blossom end rot in fruits (tomatoes, peppers); young leaves crinkled	Immobile

M
K
P
R
E
P
A
R
A
T
I
O
N
S

16. Forms & Functions in Plants



Forms & Functions in Plants: One Liner

- **Nutrition** is the process by which organisms obtain and utilize nutrients for growth, metabolism, and maintenance.
- Plants are **autotrophs**, specifically **photoautotrophs**, synthesizing organic compounds from inorganic materials using light energy.
- An **essential element** is required for a plant to complete its normal life cycle, has a specific function, and is directly involved in metabolism.
- **Hydroponics** is a technique of growing plants in a nutrient solution without soil to determine essential elements.
- In hydroponics, omitting one element at a time helps identify **essential nutrients** and study deficiency symptoms.
- There are **16 essential elements** for plant growth as defined by Arnon & Stout (1939).
- **Macronutrients** are required in amounts greater than **0.05%** of a plant's dry weight.
- **Micronutrients** or **trace elements** are required in amounts less than **0.05%** of a plant's dry weight.
- **Mobile nutrients** show deficiency symptoms first in **older leaves** (e.g., N, P, K, Mg).
- **Immobile nutrients** show deficiency symptoms first in **young leaves or meristems** (e.g., Ca, Fe, B).
- **Nitrogen** is a macronutrient absorbed as NO_3^- or NH_4^+ and is a component of amino acids, proteins, nucleic acids, chlorophyll, and ATP.
- Nitrogen deficiency causes **general chlorosis** (yellowing) starting in older leaves, stunted growth, and delayed flowering.
- **Phosphorus** is a macronutrient absorbed as $\text{H}_2\text{PO}_4^-/\text{HPO}_4^{2-}$ and is a key component of ATP, nucleic acids (DNA/RNA), and phospholipids.
- Phosphorus deficiency results in **dark green or blue-green leaves** with possible **purple/red anthocyanin pigmentation** and poor root development.
- **Potassium** is a macronutrient absorbed as K^+ and functions in **enzyme activation**, osmotic regulation, turgor (stomatal opening), and solute transport.
- Potassium deficiency causes **chlorosis and necrosis at leaf margins and tips**, a scorched appearance, and weak stalks.
- **Calcium** is a macronutrient absorbed as Ca^{2+} and is a structural component of the **middle lamella** (calcium pectate) and acts as a second messenger.
- Calcium deficiency leads to **death of apical meristems**, **blossom end rot** in fruits, and distorted young leaves.
- **Magnesium** is a macronutrient absorbed as Mg^{2+} and is the **central atom in the chlorophyll molecule** and an enzyme activator.
- Magnesium deficiency causes **interveinal chlorosis** (veins remain green) in older leaves.
- **Sulfur** is a macronutrient absorbed as SO_4^{2-} and is a component of amino acids (cysteine, methionine), coenzymes (CoA), and ferredoxin.
- Sulfur deficiency causes **general chlorosis** appearing first in younger leaves, resembling nitrogen deficiency.
- **Iron** is a micronutrient absorbed as $\text{Fe}^{2+}/\text{Fe}^{3+}$ and is essential for **chlorophyll synthesis** (as a catalyst) and redox reactions in cytochromes.
- Iron deficiency causes **severe interveinal chlorosis in young leaves**.
- **Manganese** is a micronutrient absorbed as Mn^{2+} and activates decarboxylase/dehydrogenase enzymes and is involved in the **water-splitting complex of PS II**.
- Manganese deficiency causes **interveinal chlorosis with small necrotic spots**.
- **Zinc** is a micronutrient absorbed as Zn^{2+} and is an enzyme component/activator (e.g., alcohol dehydrogenase) and essential for **tryptophan synthesis** (auxin precursor).

M
K

P
R
E
P
A
R
A
T
I
O
N
S

16. Forms & Functions in Plants

Practice MCQs

1. Which element is the central atom in the chlorophyll molecule?

- A) Iron
- B) Magnesium
- C) Calcium
- D) Potassium

Answer: Magnesium

2. Chlorosis in older leaves is a typical deficiency symptom of which nutrient?

- A) Nitrogen
- B) Calcium
- C) Iron
- D) Boron

Answer: Nitrogen

3. Which micronutrient is essential for nitrate reductase activity?

- A) Zinc
- B) Molybdenum
- C) Copper
- D) Manganese

Answer: Molybdenum

4. "Whiptail" disease in cauliflower is caused by the deficiency of:

- A) Boron
- B) Molybdenum
- C) Zinc
- D) Copper

Answer: Molybdenum

5. Carnivorous plants primarily supplement their intake of which element?

- A) Carbon
- B) Nitrogen
- C) Phosphorus
- D) Potassium

Answer: Nitrogen

6. Which plant has a modified leaf that forms a snap trap?

- A) Nepenthes
- B) Drosera
- C) Dionaea muscipula
- D) Utricularia

Answer: Dionaea muscipula

7. The primary site for gas exchange in leaves is the:

- A) Palisade mesophyll
- B) Spongy mesophyll
- C) Epidermis

D) Vascular bundle

Answer: Spongy mesophyll

8. In dorsiventral leaves, the palisade mesophyll is located towards the:

- A) Lower epidermis
- B) Upper epidermis
- C) Both sides equally
- D) Leaf margin

Answer: Upper epidermis

9. Stomatal opening is triggered directly by the absorption of:

- A) Red light
- B) Far-red light
- C) Blue light
- D) Green light

Answer: Blue light

10. The hormone that induces stomatal closure during water stress is:

- A) Auxin
- B) Gibberellin
- C) Abscisic Acid
- D) Cytokinin

Answer: Abscisic Acid

11. At the compensation point, the net exchange of which gas is zero?

- A) Oxygen
- B) Carbon dioxide
- C) Nitrogen
- D) Water vapor

Answer: Carbon dioxide

12. Which tissue is responsible for the upward conduction of water and minerals?

- A) Phloem
- B) Xylem
- C) Cambium
- D) Pericycle

Answer: Xylem

13. Dead, elongated cells with tapered ends and lignified walls in xylem are called:

- A) Vessel elements
- B) Sieve tubes
- C) Tracheids
- D) Companion cells

Answer: Tracheids

14. The only living cells in the xylem are:

- A) Tracheids
- B) Vessel elements
- C) Xylem fibres

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Chapter 17

Kingdom Animalia

Kingdom Animalia comprises multicellular, eukaryotic, heterotrophic organisms that lack cell walls. They are **ingestive feeders**, deriving nutrients by consuming other organisms. Animals typically develop from a **blastula** during embryonic development and have a dominant diploid stage. This kingdom is distinct from Protozoa, which are placed in Kingdom Protocista.

Characteristics:

- **Multicellular Eukaryotes:** Composed of eukaryotic cells without rigid cell walls. Structural support is provided by an extracellular matrix containing proteins like **collagen**.
- **Heterotrophic Nutrition:** Obligate heterotrophs that ingest and internally digest food.
- **Specialized Tissues:** Possess true tissues (except in sponges). The evolution of **nervous** and **muscle tissue** is a key innovation.
- **Blastula Formation:** A hollow ball of cells formed after zygote cleavage.
- **Sexual Reproduction:** Most reproduce sexually with haploid gametes (sperm and egg). Fertilization produces a diploid zygote.
- **Motility:** Most are motile at some life stage, aided by muscle tissues.
- **Regulative Development:** Cell fate is determined relatively late, allowing for high developmental plasticity.

Habitat & Adaptations:

- **Marine (Original):** Buoyancy, stable temperature. Adaptations include sessile attachment, burrowing, or planktonic forms.
- **Freshwater:** Challenges include osmoregulation (hypoosmotic environment) and variable conditions.
- **Terrestrial:** Major challenges are desiccation, gravity, and temperature extremes. Key adaptations include impermeable body coverings, internal respiratory surfaces, internal fertilization, amniotic eggs/vivipary, and supportive skeletons.

Animal Body Plans & Classification Criteria

A **body plan** is an integrated set of morphological and developmental traits. Key aspects are used to classify animals and infer evolutionary relationships.

1. Levels of Organization & Tissue Complexity

- **Cellular Level (Parazoa):** Cells are loosely associated; no true tissues or organs. Example: **Phylum Porifera (sponges)**.
- **Tissue Level:** Cells organized into tissues. Example: **Phylum Cnidaria**.
- **Organ & Organ System Level:** Tissues form organs and complex systems. Example: All higher phyla (**Eumetazoa**).

2. Germ Layers (Embryonic Tissue Layers)

Formed during gastrulation.

Feature	Diploblastic	Triploblastic
Germ Layers	Two: Ectoderm & Endoderm	Three: Ectoderm, Mesoderm & Endoderm
Intermediate Layer	Non-cellular Mesoglea	Cellular Mesoderm
Complexity	Limited tissue complexity.	Allows development of complex organs and systems (muscular, circulatory, skeletal).
Examples	Cnidaria, Ctenophora	All Bilateria (Platyhelminthes to Chordata)

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Kingdom Animalia: One Liner

- **Kingdom Animalia** comprises **multicellular, eukaryotic, heterotrophic** organisms that **lack cell walls**.
- Animals are **ingestive feeders**, deriving nutrients by **consuming other organisms**.
- They develop from a **blastula** during embryonic development and have a **dominant diploid** stage.
- The kingdom is **monophyletic** and also called **Metazoa**.
- Animals, fungi, and **choanoflagellates** form the **Opisthokonta** clade.
- Their structural support comes from an **extracellular matrix** containing the protein **collagen**.
- Most possess **true tissues** (except sponges); the evolution of **nervous and muscle tissue** is a key innovation.
- They exhibit **regulative development** where **cell fate is determined relatively late**.
- The **original animal habitat** is **marine**, offering buoyancy and stable temperature.
- **Freshwater** habitats pose **osmoregulation** challenges due to a **hyposmotic environment**.
- **Terrestrial colonization** required adaptations like **impermeable body coverings, internal respiratory surfaces, internal fertilization, and amniotic eggs/vivipary**.
- **Levels of organization** range from **cellular (Parazoa)** to **tissue to organ system (Eumetazoa)**.
- **Diploblastic** animals have **two germ layers (ectoderm & endoderm)** separated by a **non-cellular mesoglea** (e.g., Cnidaria).
- **Triploblastic** animals have **three germ layers (ectoderm, mesoderm, endoderm)**, allowing complex organs (all Bilateria).
- **Radial symmetry** involves arrangement around a **central axis** and is associated with **sessile or floating life**.
- **Bilateral symmetry** allows for **directed movement** and **cephalization** (sensory organs at the anterior).
- **Biradial symmetry**, a variant of radial, is found in **Ctenophora**.
- An **acoelomate** (e.g., Platyhelminthes) lacks a body cavity; space is filled with **mesenchyme/parenchyma**.
- A **pseudocoelomate** (e.g., Nematoda) has a body cavity (**pseudocoelom**) **not fully lined by mesoderm**.
- A **coelomate** has a **true coelom** fully lined by **mesoderm-derived peritoneum** (e.g., Annelida, Chordata).
- The **coelom functions** as a **hydrostatic skeleton**, provides **cushioning**, and allows **independent organ movement**.
- In arthropods, the main body cavity is a **hemocoel** where **hemolymph circulates**.
- **Protostomes** ("mouth first") exhibit **spiral, determinate cleavage** and form the coelom via **schizocoely**.
- **Deuterostomes** ("mouth second") exhibit **radial, indeterminate cleavage** and form the coelom via **enterocoely**.
- **Segmentation (metamerism)** allows for **specialization (tagmatization)** and evolved **convergently** in Annelida, Arthropoda, and Chordata.
- **Hox genes** are master regulators of **segmentation and body plan**.
- Molecular evidence identifies **choanoflagellates** as the **closest living relatives** of animals.
- The **Ediacaran Period** (~635-541 mya) saw the first **macroscopic, soft-bodied animal fossils**.
- The **Cambrian Explosion** (~541-515 mya) was the rapid diversification of most **major animal phyla**.
- **Arthropods** were the **first animals to colonize land** (~490-440 mya).
- **Porifera (sponges)** are **basal metazoans** with **cellular-level organization** and **no true tissues**.
- Sponges have **choanocytes (collar cells)** that drive **filter-feeding** through a **water canal system**.

M
K
P
R
E
P
A
R
A
T
I
O
N
S

17. Kingdom Animalia

Practice MCQs

1. Which of the following is NOT a defining characteristic of Kingdom Animalia?

- A) Multicellularity
- B) Presence of cell walls
- C) Heterotrophic nutrition
- D) Blastula formation during development

Answer: Presence of cell walls

2. Animals are distinguished from protozoans by being:

- A) Unicellular
- B) Placed in Kingdom Protocista
- C) Multicellular and ingestive feeders
- D) Autotrophic

Answer: Multicellular and ingestive feeders

3. The structural protein found in the extracellular matrix of animals is:

- A) Keratin
- B) Chitin
- C) Cellulose
- D) Collagen

Answer: Collagen

4. The hollow ball of cells formed after zygote cleavage is called:

- A) Gastrula
- B) Blastula
- C) Morula
- D) Neurula

Answer: Blastula

5. Which of the following is an autapomorphy of animals?

- A) Photosynthesis
- B) Regulative development
- C) Presence of cell walls
- D) Haploid dominant life cycle

Answer: Regulative development

6. The original habitat of animals is considered to be:

- A) Freshwater
- B) Terrestrial
- C) Marine
- D) Aerial

Answer: Marine

7. A major challenge for freshwater animals is:

- A) Buoyancy
- B) Osmoregulation
- C) Stable temperature

D) High salinity

Answer: Osmoregulation

8. Which adaptation is NOT crucial for terrestrial life?

- A) Impermeable body covering
- B) External fertilization
- C) Amniotic egg
- D) Internal respiratory surfaces

Answer: External fertilization

9. Animals with loosely associated cells and no true tissues are at which level of organization?

- A) Tissue level
- B) Organ system level
- C) Cellular level (Parazoa)
- D) Organ level

Answer: Cellular level (Parazoa)

10. True tissues are first observed in which group?

- A) Porifera
- B) Eumetazoa
- C) Parazoa
- D) Protozoa

Answer: Eumetazoa

11. Diploblastic animals possess how many germ layers?

- A) One
- B) Two
- C) Three
- D) Four

Answer: Two

12. The non-cellular layer between ectoderm and endoderm in diploblastic animals is called:

- A) Mesoderm
- B) Mesoglea
- C) Mesenchyme
- D) Peritoneum

Answer: Mesoglea

13. Triploblastic condition allows for the development of:

- A) Only epithelial tissue
- B) Simple nerve nets
- C) Complex organs and systems
- D) Choanocytes

Answer: Complex organs and systems

14. Radial symmetry is typically associated with which type of lifestyle?

M
K
P
R
E
P
A
R
A
T
I
O
N
S

17. Kingdom Animalia



Chapter 18

Nutrition and Digestion

Nutrition is the sum of all processes involved in the procurement, intake, digestion, absorption, and utilization of substances necessary for growth, maintenance, repair, and metabolic functions. Nutrients are substances that supply the body with elements essential for metabolism.

Importance of Nutrition

Aspect	Physiological & Molecular Role
Growth (Hyperplasia & Hypertrophy)	Provides substrates (amino acids, fatty acids, minerals) for the synthesis of new tissues (e.g., bone matrix, muscle protein). Critical during developmental windows (prenatal, adolescent). Hormones (GH, IGF-1) are nutrient-sensitive.
Repair & Maintenance (Homeostasis)	Enables continuous tissue turnover (e.g., intestinal epithelium renewal every 3-5 days). Nutrients act as cofactors (Zn in DNA polymerase) and antioxidants (Vitamins C & E) to mitigate oxidative damage and support apoptosis/autophagy of damaged cells.
Energy (ATP Production)	Macronutrients undergo catabolism to yield ATP: <ul style="list-style-type: none"> • Carbohydrates: Primary fuel via glycolysis & oxidative phosphorylation. • Lipids: High-yield energy reserve via β-oxidation. • Proteins: Emergency fuel via gluconeogenesis (catabolic states).

Nutrition Vs Digestion

Feature	NUTRITION	DIGESTION
Definition	Holistic process of obtaining & utilizing nutrients.	Specific breakdown of food into absorbable units.
Scope	Extremely broad (behavior, physiology, ecology).	Narrow (focused on GI tract processes).
Primary Goal	Acquire matter & energy for life functions.	Convert food into absorbable form.
Key Processes	Ingestion, Digestion, Absorption, Transport, Assimilation, Catabolism, Egestion.	Ingestion, Mechanical/Chemical Breakdown, Propulsion.
Systems Involved	Digestive, Circulatory, Lymphatic, Endocrine, Excretory, Nervous.	Primarily Digestive System & exocrine glands.
End Point	Cellular metabolism (ATP, biosynthesis).	Lumen of small intestine (simple molecules ready for absorption).
Regulation	Systemic (e.g., insulin, leptin).	Largely local (e.g., gastrin, secretin, enteric nervous system).

Fundamental Nutritional Dichotomy:

- **Autotrophy:** Organisms synthesize their own complex organic molecules from simple inorganic substances.

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Nutrition And Digestion: One Liner

M
K

P
R
E
P
A
R
A
T
I
O
N
S

- Nutrition is defined as the **sum of processes for growth, maintenance, and metabolism.**
- The **fundamental nutritional division** is between **autotrophs and heterotrophs.**
- **Autotrophs synthesize their own organic compounds** from inorganic sources.
- **Photoautotrophs**, like plants, perform **photosynthesis using light energy.**
- **Chemoautotrophs**, like nitrifying bacteria, obtain energy from **oxidizing inorganic compounds.**
- **Heterotrophs cannot synthesize their own organic compounds** and rely on other organisms.
- **Holozoic nutrition** involves **ingestion, digestion, absorption, and egestion.**
- **Saprophytic nutrition** involves **absorbing nutrients from dead matter via extracellular enzymes.**
- **Parasitic nutrition** derives nutrients from a **living host, causing harm.**
- **Symbiotic mutualism** is a relationship where **both species benefit nutritionally.**
- **Insectivorous plants** are **autotrophs that supplement nutrients** by digesting animals.
- **Herbivores** eat plant material and often have **symbiotic microbes for cellulose digestion.**
- **Carnivores** eat other animals and possess adaptations like **sharp teeth and short digestive tracts.**
- **Omnivores** consume both plants and animals and have **generalized digestive systems.**
- **Filter feeders** obtain food by **straining particles from water.**
- **Fluid feeders** consume **nutrient-rich fluids** through specialized mouthparts.
- **Detritivores** feed on **decomposing organic matter.**
- **Intracellular digestion** occurs **inside cells within food vacuoles.**
- A **gastrovascular cavity** has a **single opening serving as both mouth and anus.**
- A **complete digestive tract (alimentary canal)** has **two openings and allows one-way flow.**
- **Mammals** have **heterodont dentition**, meaning **different types of teeth.**
- The **dental formula** reflects an **animal's diet.**
- **Saliva** contains **salivary amylase (ptyalin)**, which **begins starch digestion.**
- The **stomach** secretes **HCl**, which **denatures proteins and activates pepsinogen.**
- **Pepsin** is the active **proteolytic enzyme in the stomach.**
- The **mixture of partially digested food in the stomach** is called **chyme.**
- The **small intestine** is the **primary site for digestion and absorption.**
- The **duodenum** receives secretions from the **liver and pancreas.**
- **Bile**, produced by the **liver**, **emulsifies fats** but contains **no digestive enzymes.**
- **Pancreatic juice** contains **enzymes for digesting all major macromolecules.**
- **Brush border enzymes** in the small intestine complete the digestion of **disaccharides and peptides.**
- **Hormones like secretin and CCK** **regulate pancreatic and biliary secretions.**
- **Enterokinase** is the brush border enzyme that **activates trypsinogen to trypsin.**
- **Villi and microvilli** massively increase the **surface area for absorption** in the small intestine.
- **Monosaccharides and amino acids** are absorbed into the **blood capillaries of villi.**
- Absorbed **monosaccharides and amino acids** travel to the **liver via the hepatic portal vein.**
- **Lipids** are absorbed as **micelles** and are reformed into **chylomicrons in enterocytes.**
- **Chylomicrons** enter the **lymphatic system (lacteals)** before entering the bloodstream.
- The **large intestine (colon)** primarily **reabsorbs water and electrolytes.**
- The **gut microbiome** resides in the **large intestine** and aids in **fermentation and vitamin synthesis.**
- The **liver** performs key functions including **bile production, detoxification, and glycogen storage.**
- The **gallbladder** stores and **concentrates bile.**
- The **pancreas** has both **exocrine (digestive) and endocrine (hormonal) functions.**
- **Essential amino acids** are those that **cannot be synthesized by the body.**
- There are **9 essential amino acids** for adults.
- **Essential fatty acids** include **linoleic acid and alpha-linolenic acid.**
- **Water-soluble vitamins (B-complex and C)** act as **coenzymes** and are **not stored.**
- **Fat-soluble vitamins (A, D, E, K)** are **stored in the liver and fatty tissues.**

18. Nutrition and Digestion



Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. Which of the following is the correct sequence of the digestive tract from proximal to distal?

- A) Esophagus, Stomach, Duodenum, Jejunum, Ileum
- B) Stomach, Esophagus, Ileum, Jejunum, Duodenum
- C) Duodenum, Jejunum, Ileum, Stomach, Esophagus
- D) Jejunum, Ileum, Duodenum, Stomach, Esophagus

Answer: Esophagus, Stomach, Duodenum, Jejunum, Ileum

2. Which hormone is responsible for stimulating the release of bicarbonate-rich pancreatic juice?

- A) Gastrin
- B) Secretin
- C) Cholecystokinin
- D) Gastric Inhibitory Peptide

Answer: Secretin

3. Which of the following cells secrete pepsinogen in the stomach?

- A) Parietal Cells
- B) G Cells
- C) Chief Cells
- D) Mucous Neck Cells

Answer: Chief Cells

4. The majority of nutrient absorption occurs in which part of the small intestine?

- A) Duodenum
- B) Jejunum
- C) Ileum
- D) All parts equally

Answer: Jejunum

5. Which vitamin requires intrinsic factor for its absorption in the ileum?

- A) Vitamin C
- B) Vitamin B12
- C) Vitamin D
- D) Vitamin K

Answer: Vitamin B12

6. The process of breaking down large fat globules into smaller droplets is primarily the function of:

- A) Pancreatic Lipase
- B) Bile Salts
- C) Gastric Lipase

D) Colipase

Answer: Bile Salts

7. Which of the following is NOT a primary function of the liver?

- A) Gluconeogenesis
- B) Production of Insulin
- C) Synthesis of Plasma Proteins
- D) Detoxification of Ammonia

Answer: Production of Insulin

8. The muscular contractions that mix food with digestive juices in the small intestine are called:

- A) Peristalsis
- B) Segmentation
- C) Haustration
- D) Mass Movements

Answer: Segmentation

9. Which enzyme is responsible for the digestion of starch in the mouth and small intestine?

- A) Maltase
- B) Sucrase
- C) Amylase
- D) Lactase

Answer: Amylase

10. The hormone that inhibits gastric emptying and secretion when fats are present in the duodenum is:

- A) Gastrin
- B) Secretin
- C) Cholecystokinin
- D) Motilin

Answer: Cholecystokinin

11. The main function of the large intestine is:

- A) Digestion of Proteins
- B) Absorption of Amino Acids
- C) Absorption of Water and Electrolytes
- D) Production of Bile

Answer: Absorption of Water and Electrolytes

12. Which of the following is a zymogen (inactive enzyme precursor)?

- A) Pepsin
- B) Trypsin
- C) Trypsinogen
- D) Amylase

Answer: Trypsinogen



Chapter 19

Gaseous Exchange

Respiration is the integrated physiological process that encompasses the **exchange of gases (oxygen, O₂ and carbon dioxide, CO₂)** between an organism and its environment, and the subsequent **intracellular utilization of O₂ for aerobic metabolism** to produce ATP. This fundamental process consists of four interconnected stages, ensuring O₂ delivery to cells and CO₂ removal.

1. **Pulmonary Ventilation (Breathing):** The mechanical process of moving the respiratory medium (air or water) into and out of the respiratory organs.
2. **External Respiration:** The diffusion of O₂ from the environment into the blood and CO₂ from the blood into the environment across a specialized **respiratory surface**.
3. **Transport of Respiratory Gases:** The carriage of O₂ from respiratory organs to tissues and CO₂ from tissues to respiratory organs via the circulatory system (blood or hemolymph).
4. **Internal (Cellular) Respiration:** The diffusion of O₂ from blood into tissue cells and CO₂ from cells into blood, followed by **cellular respiration** within mitochondria, where O₂ acts as the final electron acceptor in the electron transport chain.

In single-celled organisms, gas exchange occurs directly across the cell membrane. Multicellular organisms require specialized respiratory systems due to a decreased **surface area-to-volume ratio** and the increased distance of internal cells from the environment.

Fundamental Physical Principles of Gas Exchange

Gas exchange occurs via **simple diffusion** down partial pressure gradients, governed by key physical laws.

Dalton's Law of Partial Pressures: In a mixture of gases, the total pressure exerted is equal to the sum of the pressures that each gas would exert if it occupied the volume alone. The pressure of an individual gas is its **partial pressure (P_x)**, which is the critical determinant of its diffusion gradient (e.g., at sea level, atmospheric P_{O₂} ≈ 159 mm Hg).

Fick's Law of Diffusion: Quantifies the rate of diffusion (R) of a gas across a membrane: $R = (D \times A \times \Delta P) / d$.

- R = Rate of diffusion.
- D = Diffusion constant (depends on gas properties, temperature, and membrane permeability).
- A = Surface area available for diffusion.
- ΔP = Partial pressure difference across the membrane.
- d = Distance (thickness) over which diffusion occurs.

Evolutionary adaptations of respiratory systems optimize these variables: maximizing A, minimizing d, and maintaining a steep ΔP.

Henry's Law: At a constant temperature, the amount of a gas dissolved in a liquid is directly proportional to its partial pressure above the liquid. **CO₂ is approximately 20-25 times more soluble in water/plasma than O₂**, which significantly influences its transport mechanics and buffering in blood.

Properties of Efficient Respiratory Surfaces

Efficient respiratory surfaces, where external respiration occurs, share common characteristics:

- **Large Surface Area:** Relative to body volume, to maximize the area for diffusion (increases A in Fick's Law).
- **Minimal Thickness:** Often just one cell layer thick, to minimize diffusion distance (decreases d).
- **Moisture:** Gases must dissolve in a fluid before they can diffuse across a membrane.
- **Permeability:** The membrane must be readily permeable to O₂ and CO₂.
- **Rich Vascularization:** A good blood supply (in most animals) maintains a steep partial pressure gradient by rapidly carrying away O₂ and delivering CO₂ (maintains ΔP).
- **Effective Ventilation Mechanism:** Ensures a constant supply of fresh, oxygen-rich medium (air/water) to the surface, renewing the gradient.



Gas Exchange & Respiratory System: One Liner

- **Respiration** is the integrated physiological process encompassing the exchange of gases (O_2 and CO_2) between an organism and its environment and the intracellular utilization of O_2 for aerobic metabolism.
- The four interconnected stages of respiration ensure O_2 delivery to cells and CO_2 removal.
- **Pulmonary ventilation (breathing)** is the mechanical process of moving the respiratory medium (air or water) into and out of the respiratory organs.
- **External respiration** is the diffusion of O_2 from the environment into the blood and CO_2 from the blood into the environment across a specialized respiratory surface.
- **Transport of respiratory gases** is the carriage of O_2 from respiratory organs to tissues and CO_2 from tissues to respiratory organs via the circulatory system.
- **Internal (cellular) respiration** involves the diffusion of O_2 from blood into tissue cells and CO_2 from cells into blood, followed by cellular respiration within mitochondria.
- In mitochondria, O_2 acts as the final electron acceptor in the electron transport chain.
- In single-celled organisms, gas exchange occurs directly across the cell membrane.
- Multicellular organisms require specialized respiratory systems due to a decreased **surface area-to-volume ratio** and the increased distance of internal cells from the environment.
- **CO_2 is approximately 20-25 times more soluble in water/plasma than O_2** , which significantly influences its transport mechanics and buffering in blood.
- Efficient respiratory surfaces share common characteristics.
- They have a **large surface area** relative to body volume to maximize the area for diffusion (increases **A** in Fick's Law).
- They have **minimal thickness**, often just one cell layer thick, to minimize diffusion distance (decreases **d**).
- They must be **moist** because gases must dissolve in a fluid before they can diffuse across a membrane.
- The membrane must be **readily permeable to O_2 and CO_2** .
- **Rich vascularization** (a good blood supply) maintains a steep partial pressure gradient by rapidly carrying away O_2 and delivering CO_2 (maintains ΔP).
- An **effective ventilation mechanism** ensures a constant supply of fresh, oxygen-rich medium (air/water) to the surface, renewing the gradient.
- **Body surface (cutaneous)** respiration involves simple diffusion across thin, moist, highly vascularized skin and is limited by **surface area-to-volume ratio**.
- Cutaneous respiration is found in **Porifera, Cnidaria, Platyhelminthes, Earthworms, and Amphibians** (as a supplementary method).
- The **tracheal system** is a network of chitin-lined tubes (**tracheae & tracheoles**) delivering air directly to cells, bypassing the circulatory system.
- The tracheal system is found in **Insects, some Myriapods, and some Arachnids**.
- Valvular **spiracles** in the tracheal system minimize water loss.
- **Gills (branchial)** are **evaginations** (outgrowths) of the body surface specialized for aquatic exchange.
- Gills are found in **Fish, Crustaceans, Mollusks, Amphibian larvae, and some Annelids**.
- **Countercurrent flow** in gill lamellae allows **80-90% O_2 extraction** from water.
- **Lungs (pulmonary)** are **invaginations** (internal pouches) of the body surface specialized for terrestrial life.
- Lungs are found in **terrestrial vertebrates, some Snails (pulmonary cavity), and Spiders (book lungs)**.
- **Surfactant** in mammalian lungs prevents alveolar collapse.
- Air has a **higher O_2 content** than water.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

19. Gaseous Exchange

Practice MCQs

1. What is the primary driving force for gas exchange across respiratory surfaces?

- A) Active transport
- B) Osmotic pressure
- C) Diffusion down partial pressure gradients
- D) Facilitated diffusion

Answer: Diffusion down partial pressure gradients

2. Which law states that the total pressure of a gas mixture is the sum of the partial pressures of each gas?

- A) Henry's Law
- B) Boyle's Law
- C) Dalton's Law
- D) Fick's Law

Answer: Dalton's Law

3. Fick's Law of Diffusion rate is directly proportional to all except:

- A) Surface area
- B) Diffusion distance
- C) Partial pressure difference
- D) Permeability constant

Answer: Diffusion distance

4. Which of the following is NOT a characteristic of an efficient respiratory surface?

- A) Dry surface
- B) Large surface area
- C) Thin epithelium
- D) Rich blood supply

Answer: Dry surface

5. In humans, the actual site of gas exchange is the:

- A) Trachea
- B) Bronchi
- C) Alveoli
- D) Bronchioles

Answer: Alveoli

6. Which structure prevents food from entering the larynx during swallowing?

- A) Glottis
- B) Epiglottis
- C) Uvula
- D) Pharynx

Answer: Epiglottis

7. The trachea is reinforced by C-shaped rings of:

- A) Bone

- B) Smooth muscle
- C) Hyaline cartilage
- D) Elastic cartilage

Answer: Hyaline cartilage

8. Which part of the respiratory system has no cartilage but contains smooth muscle?

- A) Trachea
- B) Primary bronchi
- C) Bronchioles
- D) Larynx

Answer: Bronchioles

9. Surfactant in the alveoli is secreted by:

- A) Type I pneumocytes
- B) Type II pneumocytes
- C) Alveolar macrophages
- D) Goblet cells

Answer: Type II pneumocytes

10. The total lung capacity in an average adult human is approximately:

- A) 3 liters
- B) 5 liters
- C) 7 liters
- D) 10 liters

Answer: 5 liters

11. Residual volume is the air:

- A) Inhaled during normal breathing
- B) Exhaled forcibly after normal expiration
- C) Remaining in lungs after maximal exhalation
- D) Inhaled forcibly after normal inspiration

Answer: Remaining in lungs after maximal exhalation

12. During quiet inspiration, the diaphragm:

- A) Relaxes and moves up
- B) Contracts and flattens
- C) Remains stationary
- D) Contracts and moves up

Answer: Contracts and flattens

13. Which muscles are primarily involved in forced expiration?

- A) External intercostals
- B) Diaphragm
- C) Internal intercostals and abdominals
- D) Scalene muscles

Answer: Internal intercostals and abdominals

14. The normal breathing rate (eupnea) in adults at rest is:

- A) 5–10 breaths/min
- B) 12–20 breaths/min



Chapter 20

Circulation and Transport

Biological transport refers to the movement of materials (nutrients, gases, wastes, hormones) within an organism, essential for maintaining **homeostasis** in multicellular life forms. In very small or simple organisms (e.g., **protozoans**, sponges, hydra), **diffusion** across the cell membrane or via a **gastrovascular cavity** suffices. As body size and complexity increase, the time for diffusion becomes prohibitively long (increasing with the square of the distance), necessitating specialized **circulatory systems**. These systems overcome diffusion limitations, ensuring rapid, directed delivery and removal of substances.

Types Of Circulatory Systems

Based on architectural plan, circulatory systems are classified into two primary types.

OPEN CIRCULATORY SYSTEM

- **Definition:** The circulatory fluid, called **hemolymph**, is not always enclosed within vessels. It is pumped by a heart into open spaces or sinuses called the **hemocoel**, where it directly bathes tissues.
- **Pathway:** Heart → Arteries → Open Hemocoel/Sinuses → Tissues → Returns to heart via openings (**ostia**).
- **Characteristics:**
 - Low-pressure system.
 - **Hemolymph** is a mixture of blood and interstitial fluid; no separation.
 - Exchange occurs directly between hemolymph and cells.
 - Less efficient for rapid, targeted transport; suitable for animals with lower metabolic rates.
 - **Examples:** Most **arthropods** (insects, crustaceans) and most **molluscs** (e.g., snails, clams). In insects, gas exchange is handled by a separate **tracheal system**; hemolymph mainly transports nutrients and hormones.
 - **Respiratory Pigments:** Some (e.g., crustaceans, molluscs) use **hemocyanin** (copper-based, blue pigment).

CLOSED CIRCULATORY SYSTEM

- **Definition:** The **blood** remains confined within a continuous, branching network of vessels (arteries → capillaries → veins) throughout its circuit.
- **Pathway:** Heart → Arteries → Arterioles → Capillaries (site of exchange) → Venules → Veins → Heart.
- **Characteristics:**
 - Relatively high-pressure system.
 - Blood and interstitial fluid are separate.
 - Highly efficient, rapid, and precisely regulated flow.
 - Supports higher metabolic rates, greater body size, and activity.
 - **Examples:** **Annelids** (earthworm), some advanced invertebrates (**cephalopod molluscs** like squid, octopus), and **all vertebrates**.

OPEN VS. CLOSED CIRCULATORY SYSTEMS

Feature	Open Circulatory System	Closed Circulatory System
Fluid Containment	Not always in vessels; flows in sinuses/hemocoel.	Always confined within a continuous network of vessels.
Circulating Fluid	Hemolymph (mixed blood & interstitial fluid).	Blood (distinct from interstitial fluid).

20. Circulation and Transport

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Circulation and Transport: One Liner

- **Biological transport** is the movement of materials within an organism to maintain homeostasis.
- In small organisms like protozoans, **diffusion** across the cell membrane is sufficient for transport.
- Larger, complex organisms require specialized **circulatory systems** to overcome diffusion limitations.
- The time for diffusion increases with the **square of the distance**, making it inefficient for large bodies.
- **Circulatory systems** ensure rapid, directed delivery and removal of substances.
- Based on architectural plan, circulatory systems are classified into **open** and **closed** systems.
- An **open circulatory system** is also called a **hemocoelic system**.
- In an open system, the circulatory fluid (**hemolymph**) is not always enclosed within vessels.
- Hemolymph is pumped by a heart into open spaces called the **hemocoel**.
- Exchange in an open system occurs directly between hemolymph and cells in the hemocoel.
- Open systems are **low-pressure** and less efficient for rapid transport.
- **Examples** of open circulatory systems include most **arthropods** (insects, crustaceans) and most **molluscs**.
- In insects, gas exchange is handled by a separate **tracheal system**.
- Some invertebrates with open systems use **hemocyanin**, a copper-based blue respiratory pigment.
- A **closed circulatory system** confines **blood** within a continuous network of vessels.
- The pathway in a closed system is: Heart → Arteries → Capillaries → Veins → Heart.
- Closed systems maintain **separation between blood and interstitial fluid**.
- Closed systems are **high-pressure**, efficient, and support higher metabolic rates.
- **Examples** of closed systems include annelids (earthworms), cephalopod molluscs (squid, octopus), and **all vertebrates**.
- **Closed systems** allow precise control of flow via vasoconstriction and vasodilation.
- **Fishes** have a **two-chambered heart** (one atrium, one ventricle) with **single circulation**.
- In single circulation, blood pressure **drops after passing through gill capillaries**.
- **Amphibians** have a **three-chambered heart** (two atria, one ventricle) with **incomplete double circulation**.
- Amphibians exhibit some separation of blood via **ridges in the ventricle**.
- The amphibian skin serves as an additional respiratory surface, forming a **pulmocutaneous circuit**.
- **Most reptiles** have a **three-chambered heart** with a **partially divided ventricle**.
- Reptilian hearts allow better **control of blood flow** to lungs versus body.
- **Crocodylians, birds, and mammals** have a **four-chambered heart** with **complete double circulation**.
- A **complete double circulation** ensures **no mixing** of oxygenated and deoxygenated blood.
- The four-chambered heart is essential for supporting **endothermy** (warm-bloodedness).
- The evolution of the four-chambered heart in birds and mammals is a classic example of **convergent evolution**.
- The human cardiovascular system is a **closed, double circulatory system**.
- The heart is located in the **mediastinum** of the thoracic cavity.
- The heart is enclosed by a double-walled sac called the **pericardium**.
- The **fibrous pericardium** is the tough outer layer for protection and anchorage.
- The **serous pericardium** is a thin inner layer forming the **pericardial cavity**.
- The pericardial cavity contains **pericardial fluid** (≈15–50 mL) for lubrication.
- The heart wall has three layers: **epicardium, myocardium, and endocardium**.
- The **epicardium** is the visceral layer of the serous pericardium.
- The **myocardium** is the thick, middle layer of **cardiac muscle tissue** (cardiomyocytes).

M
K
P
R
E
P
A
R
A
T
I
O
N
S

20. Circulation and Transport

Practice MCQs

1. Which of the following is the primary function of a circulatory system in multicellular organisms?

- A) To produce hormones
- B) To maintain homeostasis by transporting materials
- C) To provide structural support
- D) To generate body heat

Answer: To maintain homeostasis by transporting materials

2. In very small organisms like protozoans, which process is sufficient for internal transport?

- A) Active transport
- B) Osmosis
- C) Diffusion
- D) Bulk flow

Answer: Diffusion

3. What is the term for the fluid circulating in an open circulatory system?

- A) Blood
- B) Plasma
- C) Hemolymph
- D) Lymph

Answer: Hemolymph

4. In an open circulatory system, where does exchange between the circulatory fluid and cells directly occur?

- A) Capillaries
- B) Hemocoel
- C) Arteries
- D) Veins

Answer: Hemocoel

5. Which of the following animals typically possesses an open circulatory system?

- A) Earthworm
- B) Squid
- C) Insect
- D) Human

Answer: Insect

6. What is the main respiratory pigment found in the hemolymph of many arthropods and molluscs?

- A) Hemoglobin
- B) Hemocyanin
- C) Chlorocruorin
- D) Myoglobin

Answer: Hemocyanin

7. Which characteristic is true for a closed circulatory system?

- A) Low-pressure system
- B) Blood and interstitial fluid are mixed
- C) Exchange occurs in sinuses
- D) Blood remains within a continuous network of vessels

Answer: Blood remains within a continuous network of vessels

8. All vertebrates possess which type of circulatory system?

- A) Open
- B) Hemocoelic
- C) Closed
- D) Gastrovascular

Answer: Closed

9. The circulatory system of an earthworm is an example of:

- A) Open system
- B) Closed system
- C) Hemocoelic system
- D) Tracheal system

Answer: Closed system

10. In a closed circulatory system, which vessels are the primary sites of exchange?

- A) Arteries
- B) Arterioles
- C) Capillaries
- D) Veins

Answer: Capillaries

11. Fish have a heart with how many chambers?

- A) One
- B) Two
- C) Three
- D) Four

Answer: Two

12. What type of circulation is found in fish?

- A) Double circulation
- B) Single circulation
- C) Incomplete double circulation
- D) Pulmocutaneous circulation

Answer: Single circulation

13. Amphibians possess a heart with how many chambers?

- A) Two
- B) Three
- C) Four



Chapter 21

Homeostasis

Homeostasis is the maintenance of a **stable internal environment** (the *milieu intérieur*) within a narrow, optimal range despite fluctuations in the external environment. It is a **dynamic equilibrium** achieved through self-regulating mechanisms, essential for optimal enzyme function and cellular metabolism. The concept was pioneered by **Claude Bernard** and later named by **Walter B. Cannon**.

Core Principles & Significance

- **Dynamic Constancy:** It is not a static, fixed state but a condition maintained within a specific, optimal range through continuous adjustments.
- **Universal Phenomenon:** Observed in all living organisms, from unicellular entities to complex animals.
- **Evolutionary Adaptation:** Enables functional independence from the external environment, allowing colonization of diverse habitats.
- **Efficiency:** Biochemical reactions and physiological processes function with maximum efficiency within narrow homeostatic ranges.

Components of a Homeostatic Control System

Homeostatic regulation operates via a **feedback loop** with three integrated components:

- **Receptor (Sensor):** Specialized structures (e.g., nerve endings, specialized cells like thermoreceptors, osmoreceptors, chemoreceptors) that detect changes (**stimuli**) in a specific physiological variable and send input to the control center.
- **Control Center (Integrator):** Typically a region of the brain (often the **hypothalamus**) or an endocrine gland. It receives input, compares it to the **set point** (desired value), and determines the appropriate corrective response. It then sends output instructions to the effector.
- **Effector:** An organ (muscle or gland) that carries out the corrective response directed by the control center, thereby influencing the regulated variable and moving it back toward the set point.

Feedback Mechanisms

- **Negative Feedback:** The **most common** homeostatic mechanism. The effector's response **counteracts or negates** the original stimulus, reversing the change and shutting off the response loop. This stabilizes the system.
 - *Process:* Stimulus → Receptor → Control Center → Effector (Response reduces stimulus) → Homeostasis restored.
 - *Examples:*
 - **Thermoregulation:** Shivering in cold, sweating in heat.
 - **Blood Glucose Regulation:** Insulin lowers high blood glucose; glucagon raises low blood glucose.
 - **Baroreceptor Reflex:** Adjusts heart rate and vessel diameter to maintain blood pressure.
 - **Renin-Angiotensin-Aldosterone System (RAAS):** Raises low blood pressure/volume.
- **Positive Feedback:** The effector's response **amplifies or reinforces** the original stimulus, driving the variable further from its set point to complete a specific event rapidly. Less common.
 - *Process:* Stimulus initiates process → Response intensifies stimulus → Cycle continues until endpoint/climax.
 - *Examples:*
 - **Blood Clotting:** Platelet aggregation releases chemicals that attract more platelets, enlarging the clot.
 - **Childbirth (Parturition):** Fetal head pressure on cervix → oxytocin release → stronger contractions → more pressure.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

21. Homeostasis



Homeostasis: One Liner

- **Homeostasis** is the maintenance of a stable internal environment within an optimal range despite external changes.
- Homeostasis represents a **dynamic equilibrium**, not a static state.
- The concept of the internal environment (*milieu intérieur*) was pioneered by **Claude Bernard**.
- The term "homeostasis" was coined by **Walter B. Cannon**.
- It is essential for optimal **enzyme function** and cellular metabolism.
- **Allotaxis** refers to achieving stability through physiological or behavioral change, often by anticipating needs.
- A homeostatic control system operates via a **feedback loop**.
- The **receptor (sensor)** is a specialized structure that detects changes (stimuli) in the environment.
- The **control center (integrator)**, often the **hypothalamus** or an endocrine gland, processes information and determines the response.
- The **effector** (muscle or gland) carries out the corrective response directed by the control center.
- The **set point** is the desired or optimal value for a physiological variable.
- **Negative feedback** is the most common homeostatic mechanism where the response counteracts the initial stimulus.
- Examples of negative feedback include body temperature regulation and blood glucose control via insulin/glucagon.
- **Positive feedback** amplifies the original stimulus to drive a process to completion.
- Examples of positive feedback include blood clotting, childbirth (oxytocin release), and nerve impulse generation (depolarization).
- **Feed-forward mechanisms** are anticipatory responses that begin *before* a change occurs (e.g., salivation at the sight of food).
- Homeostasis regulates **body temperature, blood pH, blood glucose, fluid volume, and electrolyte balance**.
- The **liver** contributes to homeostasis through nutrient balance (glucose/glycogen), detoxification, and urea synthesis.
- The **kidneys** are primary organs for osmoregulation, excretion, pH balance, and blood pressure regulation.
- The **lungs** regulate blood O₂ and CO₂ levels, which impacts pH (acid-base balance).
- The **skin** is involved in thermoregulation and minor excretion.
- The **endocrine system** provides hormonal coordination for long-term control (e.g., ADH, aldosterone, insulin).
- The **nervous system** provides rapid, short-term coordination (e.g., reflexes, hypothalamic control).
- **Blood glucose** is primarily regulated by pancreatic hormones **insulin** (lowers glucose) and **glucagon** (raises glucose).
- The normal human blood pH range is **7.35 to 7.45**.
- Blood pH is regulated by **chemical buffers** (fastest), **respiratory compensation** (lungs), and **renal compensation** (kidneys).
- The **bicarbonate buffer system** ($\text{CO}_2 + \text{H}_2\text{O} \rightleftharpoons \text{H}_2\text{CO}_3 \rightleftharpoons \text{H}^+ + \text{HCO}_3^-$) is the major extracellular buffer.
- **Circadian rhythms** cause many homeostatic set points (e.g., core body temperature, cortisol) to fluctuate on a 24-hour cycle.
- The body's master biological clock is located in the **suprachiasmatic nucleus (SCN)** of the hypothalamus.
- **Homeostatic imbalance** leads to dysfunction and disease (e.g., diabetes, hypertension).

Practice MCQs

1. Which part of the nephron is primarily responsible for the reabsorption of water via the countercurrent multiplier mechanism?

- A) Proximal convoluted tubule
- B) Distal convoluted tubule
- C) Collecting duct
- D) Loop of Henle

Answer: Loop of Henle

2. Antidiuretic hormone increases water reabsorption by altering the permeability of which nephron segment?

- A) Proximal convoluted tubule
- B) Distal convoluted tubule
- C) Collecting duct
- D) Loop of Henle

Answer: Collecting duct

3. During peritoneal dialysis, the dialysis fluid is introduced into which part of the body?

- A) Liver
- B) Abdomen
- C) Kidney
- D) Pancreas

Answer: Abdomen

4. Aldosterone primarily promotes the active reabsorption of which ion in the nephron?

- A) Sodium
- B) Calcium
- C) Potassium
- D) Bicarbonate ions

Answer: Sodium

5. In which part of the nephron does the majority of tubular reabsorption occur?

- A) Distal convoluted tubule
- B) Villi
- C) Cortical tissue
- D) Proximal convoluted tubule

Answer: Proximal convoluted tubule

6. The maximum reabsorption of filtrate components takes place in which segment of the nephron?

- A) Distal convoluted tubule
- B) Proximal convoluted tubule
- C) Ascending limb of Henle
- D) Descending limb of Henle

Answer: Proximal convoluted tubule

7. The process of detecting a change and signaling an effector's response is known as what?

- A) Negative feedback
- B) Positive feedback
- C) Inter-coordination
- D) Feedback mechanism

Answer: Feedback mechanism

8. What are the three essential components of a homeostatic regulatory mechanism?

- A) Receptors, control center, and effectors
- B) Sensory, motor, and associative neurons
- C) CNS, PNS, and diffused nervous system
- D) Cerebrum, cerebellum, and pons

Answer: Receptors, control center, and effectors

9. Blood enters the glomerulus through which vessel?

- A) Efferent arteriole
- B) Afferent arteriole
- C) Renal artery
- D) Renal vein

Answer: Afferent arteriole

10. Which portion of the nephron is primarily influenced by antidiuretic hormone?

- A) Bowman's capsule
- B) Ascending limb of Henle
- C) Distal and collecting ducts
- D) Descending limb of Henle

Answer: Distal and collecting ducts

11. The site of ultrafiltration in the nephron is known as what?

- A) Glomerulus and Bowman's capsule
- B) Proximal and distal tubules
- C) Ascending and descending limbs
- D) Loop of Henle

Answer: Glomerulus and Bowman's capsule

12. Antidiuretic hormone specifically increases the reabsorption of which substance?

- A) Amino acids
- B) Salts
- C) Ammonia
- D) Water

Answer: Water

13. Aldosterone promotes the active uptake of which ion in the ascending limb of Henle?

- A) Potassium
- B) Chloride
- C) Calcium

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Chapter 22

Support and Movement

Support and movement are fundamental characteristics of animals, enabled by specialized organ systems. In complex animals, particularly chordates, this involves an integrated system of **bones, cartilage, joints,** and **skeletal muscles**. These structures work together to provide a rigid framework, protect organs, facilitate movement through leverage, produce blood cells, and store minerals. The evolution of robust support systems became critical with increases in body size and the transition from water to land. The scientific study of bones is called **osteology**, and the study of muscles is **myology**.

THE SKELETAL SYSTEM

Functions of the Skeletal System

1. **Support:** Provides a rigid framework that maintains body shape and supports the weight of tissues.
2. **Protection:** Encloses and shields vital organs (e.g., skull protects the brain, rib cage protects the heart and lungs).
3. **Movement:** Acts as levers that are pulled by skeletal muscles to produce movement at joints.
4. **Mineral Storage:** Serves as a reservoir for **calcium** and **phosphorus**, which can be released into the bloodstream as needed. Also stores sodium and potassium.
5. **Blood Cell Production (Hemopoiesis):** **Red bone marrow** within certain bones produces erythrocytes, leukocytes, and platelets.
6. **Energy Storage:** **Yellow bone marrow** primarily consists of adipose (fat) cells, which store energy.

Types of Skeletal Systems in Animals

Type	Composition & Description	Found In	Key Features
Hydrostatic Skeleton	Fluid pressure within body compartments (coelom or gastrovascular cavity).	Soft-bodied invertebrates (e.g., annelids, cnidarians).	Uses antagonistic muscle contractions against incompressible fluid to change shape; flexible but permits only gross movements.
Exoskeleton	Rigid external covering (e.g., chitin in arthropods, calcium carbonate in molluscs).	Arthropods (insects, crustaceans), molluscs.	Provides excellent protection and muscle attachment; must be molted (ecdysis) for growth; limits maximum body size.
Endoskeleton	Internal framework of living, rigid elements (bone and/or cartilage).	Echinoderms and all chordates/vertebrates.	Grows with the organism; provides superior leverage for muscles; allows for larger body size and efficient oxygen delivery.

BONE (Osseous Tissue)

Bone is a dynamic, living, vascular **connective tissue**. It is strong yet lightweight due to its composite structure.

Chemical Composition of Bone Matrix

- **Organic Component (~35%):** Primarily **collagen type I fibers**. Provides **tensile strength** and flexibility.
- **Inorganic Component (~65%):** Primarily **hydroxyapatite crystals** (calcium phosphate, $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$). Provides **compressive strength**, hardness, and rigidity.

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

22. Support & Movement

M
K
P
R
E
P
A
R
A
T
I
O
N
S



- **System Coordination:** Effective locomotion requires the integrated function of the muscular, skeletal, nervous, circulatory, respiratory, and endocrine systems.

Support And Movement: One Liner

- The **skeletal system** provides structural support and enables movement.
- The study of bones is called **osteology**.
- The study of muscles is called **myology**.
- A robust **support system** became critical with evolutionary increases in body size and the transition from water to land.
- The primary functions of the skeletal system are **support, protection, movement, mineral storage, blood cell production, and energy storage**.
- **Support** refers to providing a rigid framework that maintains body shape.
- **Protection** involves enclosing and shielding vital organs like the brain and heart.
- **Movement** is enabled as bones act as levers pulled by skeletal muscles.
- **Mineral storage** involves serving as a reservoir for **calcium and phosphorus**.
- **Blood cell production (Hemopoiesis)** occurs in the **red bone marrow**.
- **Energy storage** is a function of **yellow bone marrow**, which consists of adipose cells.
- Animals utilize three main types of skeletal systems: **hydrostatic, exoskeleton, and endoskeleton**.
- A **hydrostatic skeleton** uses fluid pressure within body compartments.
- Hydrostatic skeletons are found in soft-bodied invertebrates like **annelids and cnidarians**.
- An **exoskeleton** is a rigid external covering.
- In arthropods, the exoskeleton is made of **chitin**.
- A key limitation of an exoskeleton is that it must be **molted (ecdysis)** for growth.
- An **endoskeleton** is an internal framework of living, rigid elements.
- Endoskeletons are found in **echinoderms and all chordates/vertebrates**.
- The endoskeleton grows with the organism and allows for larger body size.
- **Bone** is a dynamic, living, vascular connective tissue.
- The bone matrix has an **organic component (~35%)** and an **inorganic component (~65%)**.
- The organic component is primarily **collagen type I fibers**, providing **tensile strength**.
- The inorganic component is primarily **hydroxyapatite crystals**, providing **compressive strength**.
- A typical long bone has a **diaphysis (shaft)** and **epiphyses (ends)**.
- The **diaphysis** is composed of thick **compact bone** surrounding a **medullary cavity**.
- The **epiphysis** is composed of **spongy (cancellous) bone** covered by a thin layer of compact bone.
- **Articular cartilage** is a layer of hyaline cartilage covering joint surfaces.
- The **periosteum** is a tough fibrous membrane covering the outer bone surface.
- The **endosteum** is a thin cellular membrane lining inner bone surfaces.
- The **epiphyseal plate (growth plate)** is responsible for longitudinal growth in children.
- The functional unit of compact bone is the **osteon (Haversian system)**.
- The **central (Haversian) canal** contains blood vessels and nerves.
- **Concentric lamellae** are rings of calcified matrix surrounding the central canal.
- **Lacunae** are small spaces between lamellae that house **osteocytes**.
- **Canaliculi** are tiny channels allowing nutrient/waste exchange between osteocytes.
- **Perforating (Volkmann's) Canals** connect the periosteum to central canals.
- **Spongy bone** consists of a network of bony spines and plates called **trabeculae**.
- Trabeculae align along lines of mechanical stress to provide strength with minimal weight.
- **Osteoprogenitor cells** are stem cells that differentiate into osteoblasts.
- **Osteoblasts** are **bone-forming cells** that synthesize and secrete the organic matrix (**osteoid**).

M
K

P
R
E
P
A
R
A
T
I
O
N
S

22. Support & Movement

Practice MCQs

1. Bone-forming cells are called:

- A) Osteocytes
- B) Osteoclasts
- C) Osteoblasts
- D) Chondrocytes

Answer: Osteoblasts

2. The functional unit of compact bone is the:

- A) Trabecula
- B) Osteon
- C) Lacuna
- D) Canaliculus

Answer: Osteon

3. Which type of joint is immovable?

- A) Synovial
- B) Cartilaginous
- C) Fibrous
- D) Diarthrosis

Answer: Fibrous

4. The contractile unit of a muscle fiber is the:

- A) Myofibril
- B) Sarcomere
- C) Sarcoplasmic reticulum
- D) Sarcolemma

Answer: Sarcomere

5. Which protein blocks the myosin-binding sites on actin in a relaxed muscle?

- A) Troponin
- B) Titin
- C) Tropomyosin
- D) Myosin

Answer: Tropomyosin

6. Yellow bone marrow is primarily involved in:

- A) Blood cell production
- B) Mineral storage
- C) Energy storage
- D) Protection

Answer: Energy storage

7. The skull bone that is unpaired is the:

- A) Parietal
- B) Temporal
- C) Occipital
- D) Zygomatic

Answer: Occipital

8. The joint between the atlas and axis vertebrae is a:

- A) Hinge joint
- B) Pivot joint

C) Gliding joint

D) Ball-and-socket joint

Answer: Pivot joint

9. Which type of cartilage is found in the intervertebral discs?

- A) Hyaline cartilage
- B) Elastic cartilage
- C) Fibrocartilage
- D) Articular cartilage

Answer: Fibrocartilage

10. During muscle contraction, the band that shortens is the:

- A) A band
- B) I band
- C) H zone
- D) Z line

Answer: I band

11. Osteoporosis results from:

- A) Excessive bone deposition
- B) Decreased bone resorption
- C) Increased bone density
- D) Decreased bone mass

Answer: Decreased bone mass

12. The hormone that lowers blood calcium levels is:

- A) Parathyroid hormone
- B) Calcitonin
- C) Calcitriol
- D) Estrogen

Answer: Calcitonin

13. Which is a freely movable joint?

- A) Suture
- B) Symphysis
- C) Synchondrosis
- D) Synovial

Answer: Synovial

14. The bone cell responsible for bone resorption is the:

- A) Osteoblast
- B) Osteocyte
- C) Osteoclast
- D) Osteoprogenitor cell

Answer: Osteoclast

15. The scientific study of bones is called:

- A) Osteology
- B) Arthrology
- C) Myology

M
K
P
R
E
P
A
R
A
T
I
O
N
S

Chapter 23

Nervous & Sensory System

The **nervous system** is a specialized, rapid communication network that uses **electrochemical signals (nerve impulses)**. It allows animals to **detect, process, and respond** to changes in their internal and external environments. It works in tandem with the **endocrine system** (which uses hormones) to maintain **homeostasis**. The study of the nervous system is called **neurology**.

Nervous coordination involves three fundamental steps:

1. **Reception:** Detection of a stimulus by sensory receptors.
2. **Processing/Integration:** Analysis and interpretation of the sensory information within the Central Nervous System (CNS).
3. **Response:** Execution of a motor command by effectors (muscles or glands).

Sensory Receptors: Biological Transducers

A **transducer** converts energy from one form to another. **Sensory receptors** are specialized cells or neurons that act as **biological transducers**.

- **Function:** They detect specific stimuli (e.g., light, pressure, chemicals) and convert this stimulus energy into an **electrochemical signal** (a receptor or generator potential) within a sensory neuron.
- **Organization:** Receptors can be clustered to form complex **sense organs** (e.g., eye, ear) or scattered individually (e.g., in skin, viscera).


Cells of the Nervous System

Neurons (Nerve Cells)

Neurons are the **excitable**, signaling units of the nervous system, specialized for generating, conducting, and transmitting **nerve impulses (action potentials)**.

Structure of a Typical Multipolar Neuron:

- **Cell Body (Soma):** Contains the nucleus and organelles. **Nissl's granules** (clusters of ribosomes and rough ER) are present for high levels of protein synthesis. It is the metabolic and biosynthetic center.
- **Dendrites:** Short, branched, tapering extensions. They are the primary **receptive sites**, receiving signals from other neurons or receptors and conducting them *toward* the cell body.



- **Axon:** A single, long, cylindrical extension of constant diameter.
 - **Axolemma:** The plasma membrane of the axon.
 - **Axoplasm:** The cytoplasm within the axon.
 - **Axon Hillock:** Cone-shaped region where the axon originates; the site of **action potential initiation** (integrates signals from dendrites/soma).

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091



The Nervous & Sensory System: One Liner

- The **nervous system** is a highly specialized network for rapid communication within an animal's body.
- Its primary functions are **sensory input, integration, and motor output** to effectors.
- It is the foundation for behavior, homeostasis, learning, and memory.
- The study of the nervous system is called **neurology**.
- **Nervous coordination** involves three components: **receptors, neurons, and effectors**.
- The two main cell classes are **neurons** (excitable, signaling) and **neuroglia** (supportive, non-excitable).
- **Neurons** are specialized for the generation, conduction, and transmission of nerve impulses.
- A typical neuron consists of a **cell body (soma), dendrites, and an axon**.
- The **cell body** contains the nucleus and most organelles; it is the biosynthetic center.
- **Dendrites** are short, branched processes that **receive signals** and convey them *toward* the cell body.
- The **axon** is a single, long process that **conducts nerve impulses away from the cell body**.
- The axon arises from a cone-shaped region called the **axon hillock**, the site of action potential initiation.
- The axon terminates in **synaptic terminals (boutons)**, which release neurotransmitters.
- A **synapse** is the functional junction where a neuron communicates with a target cell.
- **Sensory (afferent) neurons** transmit impulses *from* sensory receptors *to* the CNS.
- **Motor (efferent) neurons** transmit impulses *from* the CNS *to* effectors (muscles/glands).
- **Interneurons (association neurons)** are located entirely within the CNS and connect sensory and motor neurons.
- **Multipolar neurons** have one axon and many dendrites; they are the most common type (e.g., motor neurons).
- **Bipolar neurons** have one axon and one dendrite (e.g., in the retina and olfactory epithelium).
- **Unipolar (Pseudounipolar) neurons** have a single process that divides; they are common in sensory neurons (dorsal root ganglia).
- **Neuroglia (glial cells)** are non-excitable, supportive cells essential for normal neuronal function.
- **Astrocytes** (CNS) are the most abundant glia; they maintain the **blood-brain barrier (BBB)** and regulate the extracellular environment.
- **Oligodendrocytes** (CNS) **myelinate** axons; one cell can myelinate segments of several axons.
- **Microglia** (CNS) are the resident **phagocytic cells** (macrophages) of the CNS.
- **Ependymal cells** (CNS) line brain ventricles and the central canal; they produce and circulate **cerebrospinal fluid (CSF)**.
- **Schwann Cells** (PNS) **myelinate** axons; one cell myelinates a single segment of one axon.
- **Satellite Cells** (PNS) surround neuron cell bodies in ganglia, providing support and regulating exchange.
- **Myelination** is the process where glial cells wrap their plasma membranes around an axon to form an insulating **myelin sheath**.
- The myelin sheath **dramatically increases the speed of nerve impulse conduction** via **saltatory conduction**.
- Gaps in the myelin sheath are called **Nodes of Ranvier**.
- In the PNS, myelin is formed by **Schwann cells**.
- In the CNS, myelin is formed by **oligodendrocytes**.
- **White matter** consists of **bundles of myelinated axons** (tracts) in the CNS.
- **Gray matter** contains neuron **cell bodies**, dendrites, unmyelinated axons, and glia; it is the site of synaptic integration.
- The **resting membrane potential (RMP)** is the voltage difference across the membrane of a non-conducting neuron, typically **-70 mV** (inside negative).

M
K

P
R
E
P
A
R
A
T
I
O
N
S

23. Nervous and Sensory System

Practice MCQs

1. Which part of the neuron primarily receives signals from other neurons?

- A) Axon
- B) Myelin sheath
- C) Dendrites
- D) Node of Ranvier

Answer: Dendrites

2. The resting membrane potential of a neuron is approximately:

- A) +70 mV
- B) 0 mV
- C) -70 mV
- D) -90 mV

Answer: -70 mV

3. Rapid, automatic responses to stimuli that do not require brain involvement are called:

- A) Voluntary actions
- B) Reflexes
- C) Instincts
- D) Hormonal responses

Answer: Reflexes

4. Which neuroglial cell forms the myelin sheath in the central nervous system?

- A) Schwann cells
- B) Astrocytes
- C) Oligodendrocytes
- D) Microglia

Answer: Oligodendrocytes

5. The neurotransmitter released at the neuromuscular junction is:

- A) Dopamine
- B) Serotonin
- C) Acetylcholine
- D) GABA

Answer: Acetylcholine

6. The part of the brain responsible for coordinating voluntary movements and balance is the:

- A) Cerebrum
- B) Medulla oblongata
- C) Cerebellum
- D) Hypothalamus

Answer: Cerebellum

7. During an action potential, the rapid influx of which ion causes depolarization?

- A) Potassium (K^+)
- B) Chloride (Cl^-)
- C) Calcium (Ca^{2+})

D) Sodium (Na^+)

Answer: Sodium (Na^+)

8. Which division of the peripheral nervous system is responsible for the "fight-or-flight" response?

- A) Somatic nervous system
- B) Parasympathetic nervous system
- C) Enteric nervous system
- D) Sympathetic nervous system

Answer: Sympathetic nervous system

9. The gaps in the myelin sheath where action potentials are regenerated are called:

- A) Synaptic clefts
- B) Nodes of Ranvier
- C) Axon hillocks
- D) Terminal boutons

Answer: Nodes of Ranvier

10. Which part of the brain acts as a major relay station for all sensory information (except smell)?

- A) Hypothalamus
- B) Thalamus
- C) Hippocampus
- D) Amygdala

Answer: Thalamus

11. The minimum level of depolarization required to generate an action potential is known as:

- A) Resting potential
- B) Threshold potential
- C) Refractory period
- D) Hyperpolarization

Answer: Threshold potential

12. Which type of neuron has one axon and one dendrite, commonly found in special sense organs?

- A) Multipolar
- B) Unipolar
- C) Bipolar
- D) Anaxonic

Answer: Bipolar

13. The vital centers for heart rate, respiration, and blood pressure are located in the:

- A) Cerebellum
- B) Pons
- C) Medulla oblongata

M
K
P
R
E
P
A
R
A
T
I
O
N
S

Chapter 24

Endocrine System

The **endocrine system** is a major **regulatory and communication network** in animals, working in close coordination with the nervous system to maintain **homeostasis**. It consists of **ductless glands** and specialized cells that secrete **hormones** directly into the bloodstream or extracellular fluid. These chemical messengers travel to distant **target cells** possessing specific **receptors**, eliciting slow but prolonged responses. This system is crucial for regulating growth, development, metabolism, reproduction, and adaptation to environmental changes.

Hormones: Nature and Function

A **hormone** is an **organic chemical messenger** secreted in minute quantities by endocrine tissues. It is transported via body fluids to specific target cells, where it regulates the rate of pre-existing biochemical processes without initiating new reactions.

Key Characteristics

- **High Potency:** Effective at extremely low concentrations (e.g., 10^{-12} M).
- **Specificity:** Acts only on target cells with complementary receptors (**Lock-and-Key Model**).
- **Regulatory Role:** Can stimulate or inhibit physiological processes.
- **Integrated Action:** Hormones often work in synergistic or antagonistic pairs (e.g., Insulin and Glucagon) to fine-tune responses.

Chemical Classification of Hormones

Chemical Class	Solubility	Examples	Key Features & Secretion Sites
Proteins/Polypeptides	Water-soluble (Hydrophilic)	Insulin, Glucagon, Growth Hormone (GH), ADH	Most common type. Stored in vesicles. Bind to cell surface receptors .
Amino Acid Derivatives	Variable	Catecholamines: Epinephrine, Norepinephrine (water-soluble). Thyroid Hormones: T ₃ , T ₄ (lipid-soluble).	Derived from tyrosine/tryptophan. Secreted by adrenal medulla (catecholamines) and thyroid.
Steroids	Lipid-soluble (Hydrophobic)	Cortisol, Aldosterone, Estrogen, Testosterone, Progesterone	Derived from cholesterol. Synthesized on demand. Bind to intracellular receptors . Secreted by adrenal cortex and gonads.
Fatty Acid Derivatives	Lipid-soluble	Prostaglandins, Leukotrienes	Act as local hormones (paracrine/autocrine). Derived from arachidonic acid.

Mechanism of Hormone Action

1. Mechanism for Water-Soluble Hormones (Proteins/Peptides, Catecholamines)

- **Receptor Location:** Transmembrane receptors on target cell surface.
- **Signal Transduction:** Involves second messenger systems.
 - **Example (GPCR-cAMP Pathway):** Hormone (1st messenger) binds → activates G-protein → activates Adenylyl Cyclase → converts ATP to cAMP (2nd messenger) →

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

24. Endocrine System

**M
K
P
R
E
P
A
R
A
T
I
O
N
S**



Endocrine System & Hormones: One Liner

- The **endocrine system** consists of **ductless glands** that secrete **hormones** directly into the bloodstream.
- **Hormones** are **chemical messengers** that regulate and coordinate the body's **metabolic activities, growth, development, and homeostasis**.
- Over **50 hormones** have been identified in the human body.
- Hormones act specifically on **target cells** that possess complementary **receptors**.
- Body communication via hormones occurs between **two endocrine glands** or between an **endocrine gland and a target organ**.
- Chemically, hormones are classified as **steroids, proteins/polypeptides, glycoproteins, amines, or amino acid derivatives**.
- **Steroid hormones** (e.g., cortisol, testosterone) are secreted by the **adrenal cortex, gonads, and placenta**.
- **Proteinaceous/Peptide hormones** (e.g., insulin, GH) are secreted by the **anterior pituitary and pancreas**.
- **Amino acid derivatives** include **thyroxine** (from thyroid) and **catecholamines** like adrenaline (from adrenal medulla).
- Unlike enzymes, hormones **do not initiate** new reactions but **regulate** existing biochemical processes.
- The **lock-and-key model** describes the specific interaction between a hormone and its receptor.
- **Endocrinology** is the study of endocrine glands and their hormones.
- Hormones are effective in extremely small amounts and are not altered by the reactions they regulate.
- **Endocrine glands** are ductless and secrete hormones into tissue spaces and body fluids.
- **Exocrine glands** secrete their products (e.g., sweat, saliva) into ducts that lead to body surfaces or cavities.
- The endocrine system is a **slow-acting, long-lasting** integrative system, contrasting with the fast, short-term nervous system.
- Hormones are active at remarkably low concentrations, typically between **10^{-8} to 10^{-10} M** in the blood.
- **Specificity** of hormone action is determined by the presence of specific **receptor molecules** on or in target cells.
- Compared to the nervous system, endocrine responses are **slower** (seconds to days) but **longer-lasting**.
- Endocrine and nervous systems are **interdependent** and function as a single integrated **neuroendocrine system**.
- **Chemical communication** includes endocrine, synaptic, paracrine, and autocrine signaling.
- A **feedback mechanism** is a cycle where the **output of a process regulates its own production**.
- **Negative feedback** is most common; it **reverses a change** to maintain homeostasis (e.g., blood glucose regulation).
- **Positive feedback amplifies a change** to complete a specific process (e.g., oxytocin in childbirth).
- In blood glucose regulation, **insulin** is released for high levels, and **glucagon** for low levels—a classic negative feedback loop.
- The sensation of **thirst** is part of a **negative feedback** loop to restore body fluid balance.
- Hormone secretion is regulated by **feedback control systems** to maintain homeostasis, not continuous.
- **Negative feedback system** counteracts a stimulus to restore stability (e.g., regulation of blood glucose, calcium, metabolic rate).

Practice MCQs

1. Which of the following is NOT a characteristic of hormones?

- A) High potency at low concentrations
- B) Initiation of new metabolic reactions
- C) Specificity for target cells
- D) Regulation of existing processes

Answer: Initiation of new metabolic reactions

2. The "master integrator" linking the nervous and endocrine systems is the:

- A) Pituitary gland
- B) Hypothalamus
- C) Adrenal medulla
- D) Pineal gland

Answer: Hypothalamus

3. Which hormone is synthesized in the hypothalamus but stored and released from the posterior pituitary?

- A) Growth Hormone
- B) Prolactin
- C) Oxytocin
- D) Adrenocorticotropic Hormone

Answer: Oxytocin

4. Insulin and glucagon are secreted by which endocrine structure?

- A) Adrenal cortex
- B) Thyroid gland
- C) Islets of Langerhans
- D) Anterior pituitary

Answer: Islets of Langerhans

5. A lipid-soluble hormone that binds to intracellular receptors is:

- A) Insulin
- B) Epinephrine
- C) Cortisol
- D) Glucagon

Answer: Cortisol

6. Which of the following is a protein hormone?

- A) Testosterone
- B) Thyroxine
- C) Growth Hormone
- D) Aldosterone

Answer: Growth Hormone

7. The hormone primarily responsible for lowering blood calcium levels is:

- A) Parathyroid Hormone
- B) Calcitonin
- C) Calcitriol

D) Aldosterone

Answer: Calcitonin

8. What is the primary effect of Antidiuretic Hormone (ADH)?

- A) Stimulates milk ejection
- B) Increases water reabsorption in kidneys
- C) Raises blood glucose
- D) Promotes sodium retention

Answer: Increases water reabsorption in kidneys

9. Which anterior pituitary hormone is tropic to the adrenal cortex?

- A) TSH
- B) ACTH
- C) FSH
- D) Prolactin

Answer: ACTH

10. A goiter is most commonly associated with a deficiency of:

- A) Iron
- B) Iodine
- C) Calcium
- D) Vitamin D

Answer: Iodine

11. The "fight-or-flight" response is primarily mediated by hormones from the:

- A) Adrenal cortex
- B) Adrenal medulla
- C) Thyroid gland
- D) Pancreas

Answer: Adrenal medulla

12. Which hormone is secreted by the pineal gland and regulates circadian rhythms?

- A) Melatonin
- B) Serotonin
- C) Insulin
- D) Cortisol

Answer: Melatonin

13. In females, luteinizing hormone (LH) directly triggers:

- A) Milk production
- B) Ovulation
- C) Follicle development
- D) Uterine contractions

Answer: Ovulation

14. Which pancreatic cell type secretes glucagon?

- A) Alpha cells

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Chapter 25

Immune System

Immunology is the study of the body's defense mechanisms against **pathogens** (disease-causing agents like bacteria, viruses, fungi, protozoa, helminths) and abnormal cells. The **immune system** is a highly coordinated network of cells, tissues, and molecules that distinguishes **self** from **nonself** and eliminates harmful entities. Its core functions are **surveillance, recognition, response, and regulation**, leading to **immunological memory**.

Immune Response: The coordinated reaction of immune cells and molecules to a foreign substance (**antigen**). A successful response provides **protection**; a dysregulated response causes **hypersensitivity, autoimmunity, or immunodeficiency**.

M Innate Immunity

Innate immunity provides immediate, non-specific defense against pathogens. It is present from birth, found in **all animals**, and does not confer immunological memory.

K Characteristics of Innate Immunity

- **Rapid response:** Activated within minutes to hours.
- **Non-specific:** Recognizes broad patterns common to groups of pathogens.
- **No memory:** The response magnitude is similar upon repeated exposure.
- **First line of defense:** Acts before adaptive immunity.

P Components of Innate Immunity

R A. Barrier Defenses

- **Skin (Integument):**
 - **Stratum Corneum:** Multiple layers of dead, keratinized cells—a formidable physical barrier.
 - **Sebaceous Glands:** Secrete **sebum** containing antimicrobial fatty acids (e.g., lauric acid).
 - **Sweat Glands:** Secretions contain **lysozyme** and **dermcidin** (antimicrobial peptide). Low pH (3-5) inhibits microbial growth.
 - **Normal Microflora:** Commensal bacteria (e.g., *Staphylococcus epidermidis*) compete with pathogens for space/nutrients (**microbial antagonism**).
- **Mucous Membranes (Mucosae):** Line gastrointestinal, respiratory, urogenital tracts.
 - **Mucus:** Viscous glycoprotein secretion that **traps** microbes. Contains **secretory IgA (sIgA)** and lysozyme.
 - **Ciliary Escalator:** In respiratory tract, ciliated epithelial cells propel mucus-trapped pathogens toward the pharynx to be swallowed.
 - **Chemical Defenses:**
 - **Stomach:** HCl and proteases (pepsin) create a highly acidic (pH ~2) proteolytic environment.
 - **Duodenum:** Bile salts disrupt bacterial membranes.
 - **Paneth Cells (Small Intestine):** Secrete **cryptdins** (α -defensins) and lysozyme.
- **Chemical Barriers:**
 - **Lysozyme:** Enzyme in tears, saliva, and mucus; digests bacterial cell walls.
 - **Acidic pH:** Gastric acid (pH ~2) and vaginal secretions create hostile environments.
 - **Antimicrobial peptides (e.g., Defensins):** Disrupt pathogen membranes.

E B. Cellular Innate Defenses

Specialized leukocytes (white blood cells) that detect and destroy pathogens.

Cell Type	Primary Role & Function	Key Feature
Phagocytes	Engulf and destroy pathogens via phagocytosis .	

M
K
P
R
E
P
A
R
A
T
I
O
N
S



Found in

All multicellular organisms

Jawed vertebrates only

Immune System: One-Liner

- **Immunology** is the study of the body's defense mechanisms against disease-causing agents.
- The **immune system** is a coordinated network of cells, tissues, and molecules that distinguishes **self** from **nonself**.
- Core functions of the immune system are **surveillance, recognition, response, and regulation**.
- An **immune response** is the coordinated reaction of immune cells and molecules to a foreign substance (**antigen**).
- A successful immune response provides **protection**; a dysregulated response causes **hypersensitivity, autoimmunity, or immunodeficiency**.
- **Pathogens** are disease-causing agents like bacteria, viruses, fungi, protozoa, and helminths.
- **Innate immunity** provides immediate, non-specific defense and is present from birth in all animals.
- It is characterized by a **rapid response** (minutes to hours), **non-specificity**, and **no immunological memory**.
- It serves as the **first line of defense** before adaptive immunity is activated.
- The **skin (integument)** is a formidable physical barrier composed of dead, keratinized cells in the **stratum corneum**.
- **Sebaceous glands** secrete **sebum** containing antimicrobial fatty acids like lauric acid.
- **Sweat glands** secrete **lysozyme** and **dermcidin**, and maintain a low pH (3–5) that inhibits microbial growth.
- **Normal microflora** (e.g., *Staphylococcus epidermidis*) compete with pathogens via **microbial antagonism**.
- **Mucous membranes** line the gastrointestinal, respiratory, and urogenital tracts.
- **Mucus** traps microbes and contains **secretory IgA (sIgA)** and lysozyme.
- The **ciliary escalator** in the respiratory tract propels mucus-trapped pathogens toward the pharynx.
- **Stomach acid** (HCl, pH ~2) and **proteases** (pepsin) create a highly acidic, proteolytic environment.
- **Bile salts** in the duodenum disrupt bacterial membranes.
- **Paneth cells** in the small intestine secrete **cryptdins (α -defensins)** and lysozyme.
- **Lysozyme** in tears, saliva, and mucus digests bacterial cell walls.
- **Antimicrobial peptides** (e.g., defensins) disrupt pathogen membranes.
- **Phagocytes** engulf and destroy pathogens via **phagocytosis**.
- **Macrophages** are “big eaters” resident in tissues; they derive from monocytes and act as **Antigen-Presenting Cells (APCs)**.
- **Neutrophils** are the most abundant white blood cells, are first responders, short-lived, and a major component of **pus**.
- **Dendritic Cells** are key APCs that phagocytose in tissues and migrate to lymph nodes to activate adaptive immunity.
- **Natural Killer (NK) Cells** destroy virus-infected and cancerous host cells by inducing **apoptosis**; they recognize cells lacking **MHC Class I**.
- **Eosinophils** discharge enzymes against multicellular parasites (e.g., helminths) and are located beneath epithelia.
- **Mast Cells** release **histamine** and other inflammatory chemicals from granules via **degranulation**.
- The **Complement System** is a group of ~30 plasma proteins that, when activated, leads to **opsonization, inflammation, and cell lysis**.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

25. Immune System

Practice MCQs

M
K
P
R
E
P
A
R
A
T
I
O
N
S

1. Which of the following is NOT a component of the body's first line of defense?

- A) Skin
- B) Stomach acid
- C) Macrophages
- D) Mucous membranes

Answer: Macrophages

2. The slightly acidic pH of human skin primarily functions to:

- A) Promote sebum production
- B) Inhibit pathogen growth
- C) Increase sweat secretion
- D) Enhance keratinization

Answer: Inhibit pathogen growth

3. Sebaceous glands contribute to skin defense by secreting sebum containing:

- A) Lysozyme
- B) Hydrochloric acid
- C) Antimicrobial fatty acids
- D) Histamine

Answer: Antimicrobial fatty acids

4. Lysozyme, an enzyme that breaks down bacterial cell walls, is found in:

- A) Sebum
- B) Sweat and tears
- C) Gastric juice
- D) Complement proteins

Answer: Sweat and tears

5. The ciliary escalator mechanism is associated with which part of the body?

- A) Skin
- B) Stomach
- C) Respiratory tract
- D) Kidney

Answer: Respiratory tract

6. Which cell is a phagocyte that is typically the first to arrive at a site of infection?

- A) Macrophage
- B) Neutrophil
- C) Dendritic cell
- D) Natural Killer cell

Answer: Neutrophil

7. Antigen-presenting cells (APCs) include all EXCEPT:

- A) Macrophages
- B) Neutrophils
- C) Dendritic cells

D) B cells

Answer: Neutrophils

8. Natural Killer (NK) cells primarily destroy target cells by releasing:

- A) Histamine and heparin
- B) Perforins and granzymes
- C) Interferons and interleukins
- D) Antibodies and complement

Answer: Perforins and granzymes

9. The complement system proteins are primarily synthesized in the:

- A) Spleen
- B) Bone marrow
- C) Liver
- D) Thymus

Answer: Liver

10. Membrane Attack Complex (MAC) formed by the complement system results in:

- A) Opsonization
- B) Chemotaxis
- C) Lysis of pathogen cell membranes
- D) Antigen presentation

Answer: Lysis of pathogen cell membranes

11. Interferons are cytokines that primarily function to:

- A) Stimulate B cell proliferation
- B) Inhibit viral replication in neighboring cells
- C) Promote histamine release
- D) Activate complement cascade

Answer: Inhibit viral replication in neighboring cells

12. The cardinal signs of inflammation are redness, heat, swelling, pain, and:

- A) Fever
- B) Loss of function
- C) Itching
- D) Pus formation

Answer: Loss of function

13. Which chemical is released by mast cells to initiate the inflammatory response?

- A) Interferon
- B) Histamine
- C) Perforin
- D) Interleukin-2

Answer: Histamine

14. Pyrogens cause fever by acting on the:

- A) Pituitary gland
- B) Hypothalamus



Chapter 26

Reproduction and Development

Developmental biology represents one of the most integrative fields in biological science, seeking to explain how **genetic information** is translated into **three-dimensional form and function**. At its core lies a profound paradox: unlike human-engineered machines that are built first and then function, organisms must **maintain physiological function** while simultaneously **constructing themselves** through embryonic development, growth, and repair. This field transcends traditional **embryology** (development from fertilization to birth) to encompass the entire lifespan, including:

- **Metamorphosis:** Radical post-embryonic transformation (e.g., caterpillar to butterfly)
- **Regeneration:** Replacement of lost body parts (e.g., salamander limbs, zebrafish heart)
- **Tissue Turnover:** Continuous renewal of cells in skin, gut, and blood
- **Ageing:** Progressive changes in structure and function over time

Historical Foundations: The Epigenesis-Preformationism Debate

The philosophical struggle to understand development shaped early biological thought:

Preformationism (17th-18th Centuries)

- **Core Belief:** A miniature, fully formed organism (**homunculus**) existed preformed within the egg or sperm
- **Proponents:** Marcello Malpighi (observed structures in unincubated chicken eggs), Nicolas Hartsoeker (drew sperm containing tiny humans)
- **Supporting Observations:** Visible organization in early embryos; difficulty imagining how complexity could arise from simplicity
- **Fatal Flaws:** Could not explain **hybrid traits** (blending of parental characteristics), **regeneration**, or the existence of **parasites within parasites** (infinite regress problem)

Epigenesis (Gained Ascendancy in 18th-19th Centuries)

- **Core Principle:** The embryo forms **de novo** ("from scratch") through progressive **differentiation** and **organization** of initially homogeneous material
- **Key Figures:**
 - **Aristotle:** First proposed epigenetic concepts, distinguishing between "potential" and "actual" form
 - **William Harvey:** "Ex ovo omnia" ("all from the egg"), emphasized gradual emergence
 - **Kaspar Friedrich Wolff:** Provided experimental evidence with chick embryos, showing heart and intestines develop from undifferentiated tissues
- **Modern Support:** Microscopic observations of cell division, differentiation, and morphogenesis

The Modern Synthesis: Genetic Preformation + Epigenetic Construction

Today's understanding reconciles both historical views:

1. **Genetic Preformation:** The fertilized egg contains **preformed genetic instructions** (the complete genome) that guide development
2. **Epigenetic Construction:** The organism is built epigenetically through:
 - **Progressive differentiation** of cells
 - **Cell-cell interactions** and signaling
 - **Responses to environmental cues** (temperature, nutrition, mechanical forces)
 - **Self-organizing properties** of tissues

This synthesis recognizes that while no miniature organism exists, the **developmental program** is pre-encoded in the DNA, awaiting execution through epigenetic processes.


Stages of Animal Development with Key Events and Significance

M
K
P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development

Stage	Definition	Key Processes	Duration (Human Example)	Significance
Fertilization	Fusion of haploid gametes to form diploid zygote	Sperm-egg recognition, membrane fusion, blocks to polyspermy, egg activation, pronuclear fusion	~24 hours	Restores diploidy; combines parental genomes; activates development
Cleavage	Rapid mitotic divisions without growth	Cell division, cytoplasmic partitioning, blastula formation	Days 1-5 (to blastocyst)	Increases cell number; partitions cytoplasm; establishes initial axes
Gastrulation	Reorganization into three-layered embryo	Cell movements (invagination, involution, etc.); germ layer formation; body plan establishment	Week 3	Most critical period; establishes body axes and germ layers
Organogenesis	Formation of organs from germ layers	Tissue interactions, morphogenesis, cell differentiation	Weeks 4-8	Builds organ systems; highly sensitive to teratogens
Fetal Period	Growth and maturation of organs	Histogenesis (tissue specialization), functional maturation, growth	Week 9 to birth	Organs become functional; massive growth
Metamorphosis	Radical transformation to adult form	Tissue remodeling, apoptosis, new growth	Variable (e.g., ~12 weeks in frogs)	Adapts organism to different ecological niches
Gametogenesis	Production of next generation's gametes	Meiosis, gamete differentiation, maturation	Puberty to reproductive senescence	Completes life cycle; enables reproduction

Early embryonic development



Comparative Embryology and Germ Layer Theory: Deep Principles

The Germ Layer Concept:

The discovery that most animals develop from three primary germ layers represents one of the most profound unifying principles in biology.

Germ Layer Derivatives with Specific Examples

Germ Layer	Major Derivatives	Specific Tissues/Organs
Ectoderm (Outer Layer)	Surface Ectoderm: Epidermis, hair, nails, lens, inner ear, enamel	Keratinocytes, melanocytes, Merkel cells
	Neuroectoderm: CNS (brain, spinal cord), retina, posterior pituitary	Neurons, astrocytes, oligodendrocytes, ependymal cells
	Neural Crest: PNS, facial bones, adrenal medulla, melanocytes	Schwann cells, dorsal root ganglia, odontoblasts
Mesoderm (Middle Layer)	Axial Mesoderm: Notochord	Nucleus pulposus of intervertebral discs
	Paraxial Mesoderm: Somites → vertebrae, skeletal muscle, dermis	Myoblasts, sclerotome, dermatome
	Intermediate Mesoderm: Urogenital system	Nephrotomes, gonads, ducts
	Lateral Plate Mesoderm: Heart, blood vessels, limbs, body wall	Cardiomyocytes, endothelial cells, limb bud mesenchyme
Endoderm (Inner Layer)	Foregut Derivatives: Pharynx, lungs, thyroid, liver, pancreas	Hepatocytes, pancreatic β -cells, pneumocytes
	Midgut Derivatives: Small intestine, proximal colon	Enterocytes, goblet cells, Paneth cells
	Hindgut Derivatives: Distal colon, rectum, bladder epithelium	Colonocytes, urothelial cells

Induction: The Dialogue Between Tissues

- **Definition:** The process by which one group of cells (**inducer**) influences the developmental fate of adjacent cells (**responder**)
- **Historical Example: Spemann-Mangold Organizer (1924)** - Transplantation of dorsal lip of blastopore induced secondary axis
- **Molecular Mechanism:** Inducer cells secrete **morphogens** (signaling molecules) that form concentration gradients
- **Types of Induction:**
 1. **Instructive:** Inducer is necessary AND sufficient to change responder fate
 2. **Permissive:** Responder already determined; inducer provides environment for expression

Von Baer's Laws: The Embryological Basis of Evolution

Karl Ernst von Baer's observations (1828) laid groundwork for evolutionary embryology:

1. **Law of General to Specific:** General characters of large taxonomic groups appear earlier than specialized characters of smaller groups.
 - *Example:* All vertebrate embryos develop a **notochord**, **pharyngeal arches**, and **post-anal tail** before developing mammalian hair or bird feathers.

2. **Law of Developmental Divergence:** Embryos of different species increasingly diverge during development; early stages are more similar.
 - *Example:* Early fish, reptile, bird, and human embryos are remarkably similar; differences emerge progressively.
3. **Law Against Recapitulation:** An embryo does NOT pass through adult stages of its ancestors (**contradicts Haeckel's Biogenetic Law**).
 - *Critical Evidence:* No mammalian embryo has functional gills like adult fish; rather, all have **pharyngeal arches** that develop differently.
4. **Law of Embryonic Similarity:** The early embryo of a "higher" animal resembles the early embryo (not adult) of a "lower" animal.
 - *Modern Concept: Phylotypic stage* - a conserved embryonic stage when embryos within a phylum look most similar (e.g., **pharyngula stage** in vertebrates).


Comprehensive Analysis of Cleavage Types and Mechanisms

Cleavage Type	Yolk Amount	Mechanism	Regulation	Examples	Developmental Significance
Holoblastic: Radial	Isolecithal (sparse, even)	Cleavage furrows pass completely through egg; blastomeres arranged radially around animal-vegetal axis	Equal partitioning of cytoplasm; microtubule asters define cleavage planes	Sea urchins, amphibians, mammals	Regulative development - early blastomeres are totipotent; enables identical twinning
Holoblastic: Spiral	Isolecithal or mesolecithal	Cleavage planes diagonal to animal-vegetal axis; blastomeres spiral around embryo	Asymmetric division influenced by spindle orientation determinants	Annelids, mollusks (except cephalopods), flatworms	Mosaic development - cytoplasmic determinants unequally partitioned; cell fate determined early
Holoblastic: Rotational	Isolecithal (mammals)	First division: meridional; second division: one meridional, one equatorial with rotation	Unique to mammals; involves compaction at 8-cell stage	Eutherian mammals, some nematodes	Compaction forms inner/outer cells; establishes first differentiation (trophoblast vs. ICM)
Meroblastic: Discoidal	Telolecithal (dense vegetal yolk)	Cleavage restricted to small blastodisc of yolk-free cytoplasm at animal pole	Cleavage furrows cannot penetrate yolk mass; regulated by marginal zone	Birds, reptiles, most fish, monotremes	Forms bilaminar disc (epiblast/hypoblast); embryo develops from epiblast only

Meroblastic: Superficial	Centrolecithal (central yolk)	Early nuclear divisions without cytokinesis → syncytium ; nuclei migrate to periphery before cellularization	Controlled by centrosom e cycles ; cellularization occurs simultaneously	Most insects (Drosophila), some crustaceans	Syncytial specification - morphogen gradients in common cytoplasm pattern nuclei before cellularization
-----------------------------	----------------------------------	---	---	--	---

M
K

P
R
E
P
A
R
A
T
I
O
N
S





Key Molecular Regulators of Cleavage:

- **Cyclin-CDK Complexes:** Control cell cycle progression; modified to eliminate G1/G2 phases
- **Aurora Kinases & Polo-like Kinases:** Regulate spindle assembly and cytokinesis
- **PAR Proteins:** Establish polarity in *C. elegans* and *Drosophila*
- **Cortical Rotation:** Microtubule-dependent movement of dorsal determinants in amphibians

Gastrulation: The Most Important Time in Your Life

Lewis Wolpert's famous quote underscores gastrulation's significance: "It is not birth, marriage, or death, but gastrulation which is truly the most important time in your life."

Core Objectives of Gastrulation:

1. **Form three germ layers** (ectoderm, mesoderm, endoderm)
2. **Establish body axes** (anteroposterior, dorsoventral, left-right)
3. **Position germ layers appropriately** for subsequent organogenesis
4. **Create primitive gut** (archenteron) from endoderm

Gastrulation Strategies Across Model Organisms

Organism	Key Structure	Cell Movements	Signaling Centers	Evolutionary Significance
Sea Urchin	Vegetal Plate	Primary mesenchyme ingression → skeletogenic cells; archenteron invagination	Wnt/β-catenin at vegetal pole specifies endoderm; Nodal specifies oral-aboral axis	Simple model for deuterostome gastrulation; shows convergent extension
Amphibian (Xenopus)	Dorsal Lip of Blastopore (Spemann organizer)	Involution of mesoderm; epiboly of ectoderm; convergent extension	Wnt/β-catenin (dorsalization); BMP4 (ventralization); Chordin/Noggin (organizer signals)	Classic model for induction and organizer function
Chick	Primitive Streak (Hensen's node at anterior end)	Epiblast cells ingress through streak → form mesoderm and endoderm; streak regresses	FGF8 maintains streak; SHH from notochord; BMP4 lateral mesoderm	Excellent for fate mapping; accessible for microsurgery
Mouse	Primitive Streak (similar to chick)	As in chick, but embryo rotates; anterior visceral endoderm prevents ectopic primitive streak	Nodal establishes streak; Brachyury (T) expressed in streak cells; Cripto co-receptor	Mammalian model; embryonic lethality of mutations informs human development
<i>Drosophila</i>	Ventral Furrow (mesoderm); Posterior Midgut Invagination	Ventral cells constrict apically → invaginate; germband extension	Twist/Snail specify mesoderm; Decapentaplegic (Dpp/BMP) patterns dorsoventral axis	Genetic model for cell shape changes and invagination


Cellular Mechanisms of Gastrulation Movements:

M
K

P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development

- Apical Constriction:** Contraction of **actin-myosin networks** at apical cell surface creates wedge-shaped cells that drive invagination
 - Regulation:* **Folded gastrulation** and **T48** genes in *Drosophila*; **Shroom** proteins in vertebrates
- Convergent Extension:** Cells intercalate mediolaterally, narrowing tissue in one dimension while lengthening in another
 - Mediation:* **Planar cell polarity (PCP)** pathway (Frizzled, Dishevelled, Van Gogh)
 - Example:* Elongation of archenteron in sea urchin; neural plate in vertebrates
- Epiboly:** Spreading of cell sheets to cover embryo
 - Mechanisms:* Cell division, cell shape change (radial intercalation), directed migration
- Cell Migration:** Individual cells move through extracellular matrix
 - Guidance:* Chemotaxis, haptotaxis, contact inhibition
 - Example:* Primordial germ cells, neural crest cells



Sex Determination

Introduction to Sexual Reproduction

Sexual reproduction generates genetic variation through fusion of haploid gametes. Sex is often determined chromosomally at fertilization, but other mechanisms exist.

Primary Sex Determination

Refers to the development of gonads (testes or ovaries) from a bipotential precursor.

Chromosomal Sex Determination Mechanisms:

Taxon	System	Key Feature
Mammals	XX = female, XY = male	SRY gene on Y chromosome triggers testes.
Birds	ZZ = male, ZW = female	System reversed compared to mammals.
<i>Drosophila</i>	XX = female, XY = male	X:A ratio determines sex; Y chromosome only for spermatogenesis.
Hymenopterans	Haplodiploidy	Fertilized (diploid) eggs → females; unfertilized (haploid) eggs → males.

The Mammalian Pathway:

- **Testis-Determining (XY): SRY** → activates **SOX9** → Sertoli cell differentiation → testes form → produce **Testosterone** and **Anti-Müllerian Hormone (AMH)**.
- **Ovary-Determining (XX):** Absence of SRY → **WNT4/RSPO1** stabilize **β-catenin** → ovary formation → **FOXL2** maintains ovarian identity.

Key Genes in Mammalian Sex Determination

Gene/Protein	Location	Role	Phenotype if Mutated
SRY	Y Chromosome	Triggers testis pathway.	XY sex reversal (female).
SOX9	Autosome	Master regulator of testis development.	XY sex reversal.
WNT4/RSPO1	Autosome	Promote ovary development.	XX sex reversal (masculinization).
FOXL2	Autosome	Maintains ovarian identity.	Postnatal ovary-to-testis transdifferentiation.

Secondary Sex Determination in Mammals: Hormonal Regulation

Development of genitalia and secondary sexual characteristics.

- **Male Phenotype:** AMH causes regression of Müllerian ducts. **Testosterone** stabilizes Wolffian ducts; converted to **DHT** by **5α-reductase** for external genitalia.
- **Female Phenotype:** Default pathway in absence of testes; Müllerian ducts persist and differentiate under **estrogen**; Wolffian ducts degenerate.

Clinical Correlates - Disorders of Sexual Development (DSD):

- **Androgen Insensitivity Syndrome (AIS):** XY, testes, but androgen receptor mutation → female external phenotype.
- **5α-Reductase Deficiency:** XY, cannot make DHT → ambiguous genitalia, virilization at puberty.
- **Congenital Adrenal Hyperplasia (CAH):** XX, excess androgens → masculinized external genitalia.

Sex Determination in Drosophila: A Cell-Autonomous Model

- Each cell decides its sex based on the X:A ratio.
- Regulatory cascade: X:A ratio → **Sex-lethal (Sxl)** activation (in females) → splicing of **transformer (tra)** → female-specific **Doublesex (Dsx^F)** protein.
- The Y chromosome is irrelevant for sex determination.

Environmental Sex Determination (ESD)

In some reptiles, sex is determined by incubation temperature.

- **Patterns:** Ia (Turtles: low = male, high = female); Ib (some lizards: opposite); II (Alligators: extremes = one sex, intermediate = other).
- **Molecular Basis:** Temperature influences expression of genes like **DMRT1**, **SOX9**, aromatase, and **β-catenin**.

Comparative Animal Reproduction and Strategies

Feature	Asexual Reproduction	Sexual Reproduction
Genetic Variation	None (clonal).	High (genetic recombination).
Number of Parents	One.	Two (typically).
Gamete Formation	None.	Involves meiosis.
Speed & Energy	Rapid, inexpensive.	Slower, costly.
Adaptability	Low in changing environments.	High; promotes diversity.



Examples	Hydra, sponges, many invertebrates.	Most vertebrates, many invertebrates.
----------	-------------------------------------	---------------------------------------

Asexual Reproduction in Invertebrates

Asexual reproduction involves the production of offspring from a single parent without the fusion of gametes, relying on mitotic cell division. It is common in stable environments where a successful genotype can be perpetuated efficiently.

Modes of Asexual Reproduction

- Fission:** The division of one cell, body, or body part into two.
 - Binary Fission:** Equal division, common in protists like amoebas and many bacteria.
 - Longitudinal Fission:** Division along the longitudinal axis, e.g., planarian flatworms.
 - Multiple/Transverse Fission:** Formation of transverse constrictions leading to a chain of individuals, e.g., some annelids like *Naididae*.
- Budding:** New individuals develop from outgrowths (buds) on the parent's body.
 - External Budding:** Buds form on the body surface, as seen in **cnidarians** (e.g., *Hydra*) and **poriferans** (sponges). Detached buds form new individuals; attached buds form colonies.
 - Internal Budding (Gemmulation):** In some freshwater sponges, aggregates of stem cells are encased in tough, cyst-like **gemmules**. Upon the parent's death and under favorable conditions, these cells form a new organism.
- Fragmentation:** A new organism regenerates from a lost body part. This requires significant regenerative capacity and is seen in some cnidarians (e.g., sea anemones), platyhelminths, and echinoderms.
- Parthenogenesis:** Development of an embryo from an unfertilized egg.
 - Eggs may be diploid (via suppressed meiosis) or haploid.
 - Offspring can be all female (diploid eggs), all male (haploid eggs, as in hymenopteran drones), or a mix.
 - It allows rapid population growth in stable conditions and is integral to social structure in bees, ants, and wasps. Some parthenogenetic species (e.g., certain rotifers, aphids) can revert to sexual reproduction under environmental stress to generate genetic diversity.
 - Also occurs in some vertebrates: certain lizards (e.g., *Aspidoscelis neomexicana*), snakes (e.g., *Indotyphlops braminus*), and fishes (e.g., *Stegostoma fasciatum*).

Genetic and Evolutionary Implications

- Asexual populations are **clonal**, leading to low genetic diversity.
- Evolution relies solely on mutation as a source of variation, without genetic recombination from independent assortment or crossing over.
- This can lead to rapid fixation of adaptive mutations but also makes populations vulnerable to environmental change or novel pathogens, as deleterious mutations are passed to all offspring.

Advantages and Disadvantages of Asexual Reproduction

Advantages	Disadvantages
Rapid population increase from a single individual.	Lack of genetic diversity reduces adaptability to changing environments.
Energetically efficient; no need to find a mate.	Accumulation and inheritance of deleterious mutations.
Successful genotypes are preserved intact in stable environments.	High risk of population collapse from a single disease or environmental shift.
Ideal for colonizing predictable, favorable habitats.	Evolutionary change is slower and dependent on rare beneficial mutations.



Sexual Reproduction in Invertebrates

Sexual reproduction involves meiosis to produce haploid gametes (eggs and sperm), followed by fertilization (**syngamy**) to form a genetically unique diploid zygote. This process generates significant genetic diversity within populations.

Modes of Fertilization

1. External Fertilization (Broadcast Spawning):

- Gametes are released into the surrounding environment (typically water).
- Common in aquatic invertebrates (e.g., many sponges, corals, polychaete worms).
- Requires synchronization via environmental cues (temperature, light cycles) and pheromones.
- Gonads are often simple and transient. High numbers of gametes are produced to offset low fertilization rates.

2. Internal Fertilization:

- Sperm are directly transferred from male to female via a **copulatory organ** (e.g., penis, cirrus).
- Requires specialized reproductive systems.
- **Spermatophores**: Protective packets containing sperm are used by some groups (e.g., cephalopods, leeches, arthropods) for efficient transfer.

Variations in Sexual Systems

- **Gonochorism (Dioccy)**: The common condition with separate male and female individuals.
- **Hermaphroditism (Monoecy)**: An individual possesses both functional male and female reproductive systems.
 - **Simultaneous Hermaphroditism**: Individuals can produce both gametes simultaneously, often mating reciprocally (e.g., earthworms, *Lumbricus terrestris*; many pulmonate snails). Beneficial for sessile or low-density populations.
 - **Sequential Hermaphroditism**: Individuals change sex during their lifecycle.
 - **Protandry**: Male first, then female (e.g., some oysters, *Ostrea edulis*).
 - **Protogyny**: Female first, then male. Often induced by population sex ratio changes.

Advantages and Disadvantages of Sexual Reproduction

Advantages	Disadvantages
Generates extensive genetic diversity through crossing over, independent assortment, and random fertilization.	Energetically costly: requires production of gametes, finding mates, and often complex behaviors/structures.
Facilitates removal of deleterious alleles from populations.	"Dismantles" successful parental genotypes in offspring.
Provides the raw material (variation) for evolution by natural selection.	Many gametes go unfertilized, representing a metabolic loss.
Enhances population resilience to parasites, diseases, and environmental fluctuations.	Requires coordination between two individuals.

Sexual Reproduction in Vertebrates



Vertebrates exhibit diverse reproductive strategies shaped by the transition from aquatic to terrestrial life. Key adaptations include internal fertilization and specialized eggs/gestation.

Basic Vertebrate Reproductive Strategies

Strategy is defined by where fertilization occurs and how the embryo is nourished.

Strategy	Fertilization	Embryonic Development & Nourishment	Examples
Oviparity	External or Internal	Eggs laid externally; embryo nourished by yolk. No direct maternal nourishment after laying.	Most fishes, amphibians, reptiles (incl. birds), monotremes.
Ovoviviparity	Internal	Eggs retained inside female; embryo nourished by yolk. Young born live.	Some sharks, rays, reptiles (e.g., garter snakes), and invertebrates.
Viviparity	Internal	Young retained in female reproductive tract; embryo nourished directly by mother via a placenta or similar structure.	Most mammals (eutherians & marsupials), some sharks, reptiles.

- **Gestation:** The period of carrying developing young within the female.

Reproduction in Major Vertebrate Classes

1. Fishes:

- **Actinopterygii & Sarcopterygii (Bony Fishes):** Mostly oviparous with **external fertilization**. Produce vast numbers of eggs with small yolk reserves. High mortality offset by rapid development.
- **Chondrichthyes (Cartilaginous Fishes):** **Internal fertilization** via modified pelvic fins (claspers). Many are ovoviviparous or viviparous. Some are oviparous, laying eggs in protective cases ("mermaid's purses").

2. Amphibians:

- Lifecycle tied to water due to permeable skin and (usually) aquatic larvae.
- **External fertilization** is most common (e.g., anurans use **amplexus** for sperm release over eggs).
- Internal fertilization occurs in salamanders and caecilians.
- Exhibit **biphasic development**: aquatic larval stage (tadpole) undergoes **metamorphosis** into (often) terrestrial adult.

3. Reptiles (Nonavian & Avian):

- Key innovations for terrestrial life: **internal fertilization** and the **amniotic egg**.
- **Amniotic Egg:** Contains extraembryonic membranes—**amnion** (fluid-filled protection), **chorion** (gas exchange), **allantois** (waste storage/gas exchange)—and a leathery or calcareous shell.
- **Nonavian Reptiles:** Mostly oviparous; eggs often deposited and left unattended (except crocodylians).
- **Birds (Avian Reptiles):** All oviparous. Thick, calcified shells allow **brooding** (incubation). Exhibit high levels of **parental care** for altricial (helpless) or precocial (mobile) hatchlings. Internal fertilization via cloacal contact.

4. Mammals:

- **Monotremes (Prototheria):** Oviparous (e.g., platypus, echidna). Retain egg-laying from synapsid ancestors.
- **Therians:** Viviparous.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development

- **Marsupials (Metatheria):** Short gestation, followed by prolonged development in an external **marsupial pouch** (e.g., kangaroos, opossums).
- **Placental Mammals (Eutheria):** Long gestation; embryo nourished via a complex **placenta**. After birth, young nourished with milk from **mammary glands**.
 - **Primate Reproduction:** Many are **asynchronous breeders** (year-round mating). Human females have concealed ovulation and can mate throughout the cycle. In promiscuous primates (e.g., chimpanzees), **sperm competition** leads to larger testes and higher sperm counts.

HUMAN MALE REPRODUCTIVE SYSTEM

The human male reproductive system is a complex network of organs designed for three primary functions:

1. **Spermatogenesis:** The production of male gametes (spermatozoa).
2. **Hormone Production:** The secretion of androgens, primarily **testosterone**, which regulate reproductive function and secondary sexual characteristics.
3. **Sperm Delivery:** The transport, maturation, and deposition of sperm into the female reproductive tract via the seminal fluid.

It becomes functional at **puberty** under the influence of the hypothalamic-pituitary-gonadal (HPG) axis and remains active throughout adult life.

Anatomical Organization and Histology

A. Primary Sex Organ: The Testes


- **Location & Temperature Regulation:** Paired, oval organs housed within the **scrotum**, an external skin pouch. The scrotum maintains the testes at approximately **34–35°C**, which is **2–3°C below core body temperature**, a critical condition for viable spermatogenesis. This is achieved via the **cremasteric reflex** (muscle contraction/relaxation) and the **pampiniform plexus** (a countercurrent heat exchange system of veins surrounding the testicular artery).
- **Internal Structure:** Each testis is encapsulated by the fibrous **tunica albuginea**. It is divided into approximately **250-300 lobules**.
 - **Seminiferous Tubules:** Each lobule contains 1-4 highly coiled **seminiferous tubules**, the functional units for sperm production (total length ~250 meters per testis). These tubules are lined by a specialized **germinal epithelium**.
 - **Interstitial Tissue:** The spaces between the tubules contain blood vessels, lymphatic vessels, and **Leydig (interstitial) cells**.

Cellular Composition of the Seminiferous Tubule:

1. **Spermatogenic Cells:** A stratified epithelium in various stages of spermatogenesis.
 - **Spermatogonia:** Diploid (2n) stem cells located at the basement membrane. Type A (self-renewing) and Type B (committed to differentiation).
 - **Primary Spermatocytes:** Large cells undergoing Meiosis I.
 - **Secondary Spermatocytes:** Quickly undergo Meiosis II.
 - **Spermatids:** Haploid (n) cells undergoing spermiogenesis (differentiation).
 - **Spermatozoa:** Mature sperm released into the tubule lumen.
2. **Sertoli Cells (Sustentacular Cells):** Large, columnar somatic cells that span the entire epithelium from basement membrane to lumen. They are crucial for:
 - **Structural Support:** Form the **blood-testis barrier (BTB)** via tight junctions, creating an immunologically privileged adluminal compartment.
 - **Nutrition & Metabolism:** Provide nutrients (lactate, pyruvate) to developing germ cells.
 - **Phagocytosis:** Engulf residual cytoplasm shed during spermiogenesis and apoptotic germ cells.

- **Secretion:** Produce **Androgen-Binding Protein (ABP)** (to maintain high local testosterone), **Anti-Müllerian Hormone (AMH)** during fetal development, **inhibin**, and fluid for sperm transport.
3. **Leydig (Interstitial) Cells:** Located in the connective tissue *between* seminiferous tubules. Their primary function is the **synthesis and secretion of testosterone** in response to Luteinizing Hormone (LH).

Male Reproductive System



The Reproductive (Sperm) Duct System

This series of ducts transports, matures, and stores sperm.

1. **Rete Testis:** A network of channels that collect sperm from the seminiferous tubules.
2. **Efferent Ductules (Vasa Efferentia):** 12-20 small, ciliated ducts that carry sperm from the rete testis **out of the testis** into the epididymis. They also reabsorb testicular fluid.
3. **Epididymis:** A single, highly coiled tube (~6 meters long) located on the posterior surface of each testis. It is divided into:
 - **Head (Caput):** Receives sperm from efferent ductules.
 - **Body (Corpus):** Site of major functional maturation.
 - **Tail (Cauda):** Principal site for sperm **storage** (can remain viable for weeks). The epididymis provides an environment for sperm to gain **motility** and **fertilizing capacity** (capacitation potential).
4. **Vas Deferens (Ductus Deferens):** A thick-walled, muscular tube (~45 cm) that transports sperm from the epididymis tail. It ascends through the **spermatic cord**, passes through the **inguinal canal**, and loops over the ureter to enter the pelvic cavity. Its terminal portion, the **ampulla**, dilates before joining the seminal vesicle duct.
5. **Ejaculatory Duct:** Formed by the union of the **vas deferens** and the **duct of the seminal vesicle**. It is short (~2 cm) and passes through the prostate gland to empty into the prostatic urethra.
6. **Urethra:** The terminal duct serving as a common passage for both semen (**reproductive function**) and urine (**excretory function**). It has three regions:
 - **Prostatic Urethra:** Passes through the prostate.
 - **Membranous Urethra:** A short segment through the urogenital diaphragm.
 - **Spongy (Penile) Urethra:** Runs through the length of the penis.

Accessory Glands

These glands produce seminal plasma, the fluid component of semen, which nourishes sperm, provides a transport medium, and neutralizes hostile environments.

Gland	Contribution to Semen	Key Secretions & Functions
Seminal Vesicles (Paired)	60-70% of volume.	Fructose: Primary energy source for sperm motility. Prostaglandins: Stimulate uterine/oviduct contractions to aid sperm transport. Fibrinogen-like proteins: Aid in coagulation of semen post-ejaculation. Ascorbic acid, Flavins: Antioxidants.
Prostate Gland (Single)	20-30% of volume.	Citrate & Zinc: Nutrient and cofactor. Prostate-Specific Antigen (PSA): A serine protease that liquefies coagulated semen 15-30 minutes after ejaculation, releasing sperm. Phosphatases, Fibrinolysin. Acidic Phosphatase: Enzyme marker for prostatic secretion.
Bulbourethral (Cowper's) Glands (Paired)	~5% (Pre-ejaculate).	Clear, alkaline mucus: Neutralizes traces of acidic urine in the urethra and lubricates the penile urethra during sexual arousal.

Semen Composition: A typical ejaculate is 2-5 ml, containing 20-150 million sperm/ml. It is a mixture of **spermatozoa** (from testes/epididymis) and **seminal plasma** (from accessory glands). It is slightly alkaline (pH 7.2-7.8) to neutralize the acidic vaginal environment.

External Genitalia: The Penis

The penis is the copulatory organ. It consists of:

- **Root:** Attached to the pelvic bone.
- **Body (Shaft):** Composed of three cylindrical masses of **erectile tissue**:
 1. **Two Corpora Cavernosa:** Dorsal columns. Primarily involved in erection.
 2. **One Corpus Spongiosum:** Ventral column, surrounding the urethra. Expands distally to form the **glans penis**, which is covered by the **prepuce (foreskin)** unless removed by circumcision.
- **Erectile Mechanism:** Upon sexual stimulation, parasympathetic nerves release **nitric oxide (NO)**. NO causes vasodilation of the **helicine arteries**, leading to increased blood flow into the erectile tissue sinuses. Engorgement compresses the venous drainage veins, trapping blood and causing **erection**. Detumescence is mediated by sympathetic stimulation.

Hormonal Regulation: The Hypothalamic-Pituitary-Gonadal (HPG) Axis

This neuroendocrine axis provides precise, feedback-controlled regulation of testicular function.

1. **Hypothalamus:** Secretes **Gonadotropin-Releasing Hormone (GnRH)** in a pulsatile manner into the hypophyseal portal system.
2. **Anterior Pituitary:** GnRH stimulates the synthesis and release of two gonadotropins:
 - **Luteinizing Hormone (LH):** Acts on **Leydig cells** to stimulate **testosterone** biosynthesis from cholesterol via a cAMP-mediated pathway.
 - **Follicle-Stimulating Hormone (FSH):** Acts on **Sertoli cells** to stimulate the production of **Androgen-Binding Protein (ABP)** and other factors essential for supporting spermatogenesis (e.g., transferrin, inhibin).
3. **Testicular Hormones & Feedback:**
 - **Testosterone:** The primary androgen. It has both **endocrine** and **paracrine** functions.
 - **Local (Paracrine):** High concentrations within the seminiferous tubule (bound to ABP) are absolutely required for the completion of meiosis and spermiogenesis.



- **Systemic (Endocrine):** Responsible for the development and maintenance of **male secondary sexual characteristics** (facial/body hair, deep voice, muscle mass), libido, and anabolic effects.
- **Feedback:** Exerts **negative feedback** on the hypothalamus (reducing GnRH pulse amplitude) and directly on the pituitary (reducing LH secretion).
- **Inhibin:** A glycoprotein hormone (mainly inhibin B) secreted by **Sertoli cells** in response to FSH. It provides **selective negative feedback** on the anterior pituitary to suppress FSH secretion, allowing fine-tuning of spermatogenesis without affecting LH.

Clinical Correlation: Conditions like **Klinefelter Syndrome (47, XXY)** result in testicular dysgenesis, leading to reduced testosterone, elevated LH/FSH, and impaired spermatogenesis.

HUMAN FEMALE REPRODUCTIVE SYSTEM

Introduction and Functional Overview

The human female reproductive system is designed for:

1. **Oogenesis:** Production of female gametes (oocytes/ova).
2. **Hormone Production:** Secretion of **estrogens** (primarily estradiol) and **progesterone**, which regulate the reproductive cycles, pregnancy, and secondary sexual characteristics.
3. **Fertilization & Gestation:** Provision of a site for fertilization, implantation, and embryonic/fetal development.
4. **Parturition & Lactation:** Facilitation of childbirth and nourishment of the newborn.

It exhibits **cyclical activity** (the menstrual cycle) from menarche to menopause, in contrast to the continuous activity of the male system.

Anatomical Organization and Histology

Primary Sex Organs: The Ovaries

- **Location:** Paired, almond-shaped organs (3-5 cm long) located in the pelvic cavity, attached by the **mesovarium**.
- **Structure:** Each ovary is covered by a single layer of cuboidal **germinal epithelium** (a misnomer; it is not the source of germ cells). Underneath lies the dense connective tissue **tunica albuginea**. The ovary consists of:
 - **Cortex:** Outer region containing **ovarian follicles** at various stages of development, embedded in a stromal matrix.
 - **Medulla:** Central region with loose connective tissue, blood vessels, and nerves.

The Ovarian Follicle: The functional unit, consisting of an **oocyte** surrounded by somatic support cells.

1. **Primordial Follicle:** The resting pool. A **primary oocyte** arrested in **Prophase I**, surrounded by a single layer of flattened **pregranulosa cells**. Millions present at birth; only a few hundred will ovulate.
2. **Primary Follicle:** The oocyte enlarges, and granulosa cells become cuboidal and proliferate to form multiple layers. A glycoprotein layer, the **zona pellucida**, forms between the oocyte and granulosa cells.
3. **Secondary (Antral) Follicle:** Fluid-filled spaces appear among granulosa cells, coalescing into a single **antrum**. The oocyte is surrounded by a mound of granulosa cells (**cumulus oophorus**) and sits off to one side. Stromal cells differentiate into the **theca interna** (androgen-producing) and **theca externa** (fibrous capsule).
4. **Mature (Graafian) Follicle:** The antrum is large and the follicle bulges from the ovarian surface. The primary oocyte completes Meiosis I just before ovulation, producing the **secondary oocyte** and first polar body.

The Reproductive (Genital) Ducts

1. **Uterine Tubes (Fallopian Tubes/Oviducts):** ~10 cm long, divided into regions:
 - **Infundibulum:** Funnel-shaped opening with finger-like **fimbriae** that sweep over the ovary to capture the ovulated oocyte.
 - **Ampulla:** Widest, longest region; the **most common site of fertilization**.

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091




- **Isthmus:** Narrow segment connecting to the uterus.
- **Histology:** Lined by a mucosa with **ciliated cells** (beat toward the uterus) and **secretory peg cells** (provide nutrients). The muscularis has inner circular and outer longitudinal layers for peristaltic movement.
- 2. **Uterus (Womb):** A pear-shaped, thick-walled, muscular organ.
 - **Regions: Fundus** (top), **Body (Corpus)** (main part), **Cervix** (neck projecting into vagina).
 - **Layers:**
 - **Perimetrium:** Outer serosal layer (visceral peritoneum).
 - **Myometrium:** Thick middle layer of smooth muscle arranged in interlacing bundles. Contracts rhythmically during labor.
 - **Endometrium:** Inner mucosal lining. It has two layers:
 - **Stratum Functionalis:** Thick, superficial layer rich in glands and spiral arteries. It undergoes **cyclic growth, secretion, and shedding** during each menstrual cycle.
 - **Stratum Basalis:** Thin, deep layer adjacent to myometrium. It remains intact and regenerates the functionalis each cycle.
- 3. **Vagina:** A fibromuscular tube (~8-10 cm) extending from the cervix to the vestibule.
 - **Function:** Receives semen during intercourse, serves as the **birth canal**, and allows passage of menstrual flow.
 - **Histology:** Lined by stratified squamous non-keratinized epithelium that thickens under estrogen influence. The lamina propria is rich in elastic fibers. No glands; lubrication comes from cervical mucus and transudation from vaginal walls.

External Genitalia (Vulva)

- **Mons Pubis:** Fatty pad over the pubic symphysis.
- **Labia Majora:** Hair-covered skin folds (homologous to male scrotum).
- **Labia Minora:** Hairless, vascular folds medial to the labia majora.
- **Vestibule:** Area enclosed by labia minora, containing openings of the **urethra** and **vagina**.
- **Clitoris:** Small, erectile organ composed of **glans, body, and crura** (homologous to the penis). It is highly innervated and a primary focus of sexual sensation.
- **Vestibular Glands (Bartholin's Glands):** Homologous to male bulbourethral glands; secrete mucus for lubrication.

Mammary Glands (Modified Sweat Glands)

- **Structure:** Located within the breasts. Each gland consists of 15-20 **lobes**, each divided into **lobules** containing **alveoli** (milk-secreting sacs). Alveoli drain into **lactiferous ducts**, which converge at the **nipple**.
- **Hormonal Control:** Development is estrogen/progesterone-dependent. Milk production (**lactogenesis**) is stimulated by **prolactin**. Milk ejection (**let-down**) is stimulated by **oxytocin**.



M
K
P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development

The Ovarian and Menstrual (Uterine) Cycles

These two interconnected, hormonally driven cycles (~28 days) prepare the body for potential pregnancy.

The Ovarian Cycle

Phase (Days)	Key Events	Hormonal Control
Follicular Phase (1-13)	Recruitment of a cohort of follicles from the primordial pool. One becomes dominant (Graafian follicle) . The primary oocyte completes Meiosis I.	FSH stimulates growth. Growing follicles secrete estradiol . Low estradiol inhibits FSH/LH (negative feedback). High estradiol near mid-cycle triggers positive feedback , causing the LH surge .
Ovulation (~Day 14)	The LH surge induces final oocyte maturation, follicular wall weakening, and rupture, releasing the secondary oocyte .	LH surge is the direct trigger.
Luteal Phase (15-28)	The ruptured follicle collapses and transforms into the corpus luteum ("yellow body"), a temporary endocrine gland.	LH maintains the corpus luteum. It secretes progesterone and some estradiol. If no pregnancy, it degenerates into a corpus albicans ("white body") after ~10 days.



The Menstrual (Uterine) Cycle

Phase (Days)	Endometrial Condition	Hormonal Driver
Menstrual Phase (1-5)	Sloughing of the Stratum Functionalis. Spiral arteries constrict, causing ischemia and tissue necrosis. Blood, tissue fluid, and cellular debris are discharged (menses).	Sharp decline in estradiol and progesterone from the regressing corpus luteum.
Proliferative (Follicular) Phase (6-14)	Rapid regeneration and growth of the functionalis. Endometrial glands elongate; spiral arteries grow. The endometrium thickens from ~1 to 3-5 mm.	Estradiol from the developing ovarian follicles.
Secretory (Luteal) Phase (15-28)	Endometrium becomes secretory and highly vascularized (up to 6 mm). Glands become tortuous and secrete glycogen-rich fluid. Spiral arteries become highly coiled. This creates a hospitable environment for implantation .	Progesterone (and estradiol) from the corpus luteum .

M
K

P
R
E
P
A
R
A
T
I
O
N
S

Hormonal Regulation: The Integrated HPG Axis

- Early Follicular Phase:** Low ovarian hormones → **negative feedback** on pituitary is lifted → FSH rises → stimulates follicle growth.
- Mid-Late Follicular Phase:** Dominant follicle secretes high **estradiol**. Sustained high estradiol for ~36 hours switches feedback to **positive** → triggers **GnRH surge** and consequent **LH/FSH surge**.
- Luteal Phase:** **Corpus luteum** secretes high progesterone and estradiol → strong **negative feedback** on GnRH and gonadotropins → prevents new follicle development.
- Late Luteal Phase:** If no pregnancy, corpus luteum regresses → progesterone/estradiol plummet → negative feedback removed → FSH begins to rise, initiating the next cycle.
- Pregnancy:** If implantation occurs, the embryo-derived hormone **Human Chorionic Gonadotropin (hCG)** rescues and maintains the corpus luteum until the placenta takes over hormone production (~week 10).

Clinical and Biological Correlations

- **Endometriosis:** Growth of endometrial tissue outside the uterus, causing pain and infertility.
- **Polycystic Ovary Syndrome (PCOS):** Characterized by anovulation, hyperandrogenism, and multiple cystic follicles.
- **Menopause:** Cessation of ovarian follicular activity and cycles (~45-55 yrs), leading to low estrogen and symptoms like hot flashes and osteoporosis.
- **Fertility Window:** Highest probability of conception is in the 5 days leading up to and including the day of ovulation, reflecting sperm survival and oocyte viability.

26. Reproduction and Development



Key Reproductive Hormones and Functions


Hormone	Source (Female)	Primary Functions (Female)	Source (Male)	Primary Functions (Male)
GnRH	Hypothalamus	Stimulates FSH/LH release.	Hypothalamus	Stimulates FSH/LH release.
FSH	Anterior Pituitary	Stimulates follicular growth; increases estradiol.	Anterior Pituitary	Stimulates Sertoli cells; supports spermatogenesis.
LH	Anterior Pituitary	Triggers ovulation; forms/maintains corpus luteum.	Anterior Pituitary	Stimulates Leydig cells to produce testosterone.
Estradiol	Ovarian Follicles, Corpus Luteum	Endometrial proliferation, secondary sex characteristics, positive feedback for LH surge.	Minor (from aromatization)	Minimal role.
Progesterone	Corpus Luteum, Placenta	Endometrial secretory transformation, maintains pregnancy, inhibits uterine contractions.	-	-
Testosterone	Theca Cells (precursor)	Precursor for estradiol.	Leydig Cells	Spermatogenesis, secondary sex characteristics, anabolism.
Inhibin	Granulosa Cells	Inhibits FSH selectively.	Sertoli Cells	Inhibits FSH selectively.
hCG	Syncytiotrophoblast (Embryo)	Rescues corpus luteum (pregnancy test marker).	-	-

- **Menarche:** First menstruation.
- **Menopause:** Cessation of cycles (~age 50).

M
K
P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development

Menstrual cycle



M
K

P
R
E
P
A
R
A
T
I
O
N
S

Gametogenesis

Gametogenesis is the process by which diploid **primordial germ cells (PGCs)** undergo meiotic division and differentiation to form haploid, functional gametes: **spermatozoa** in males and **oocytes/ova** in females. While both processes share the goal of producing haploid cells through meiosis, they differ profoundly in timing, pattern, and outcome, reflecting the distinct biological roles of sperm (motile, numerous) and eggs (large, nutrient-rich). The fundamental difference is encapsulated in the concept of **anisogamy** – the production of two dissimilar gametes.

Primordial Germ Cell (PGC) Specification and Migration

- **Origin:** PGCs are specified very early in embryonic development, outside the gonads (e.g., in the epiblast of mice, yolk-sac endoderm in other vertebrates). They are transcriptionally repressed to maintain pluripotency.
- **Migration:** Guided by chemotactic signals like **SDF1/CXCR4**, PGCs migrate via the hindgut and dorsal mesentery to colonize the developing **bipotential (indifferent) gonadal ridges**. This migration is supported by a niche of cells secreting **Stem Cell Factor (SCF)**.
- **Sexual Differentiation:** Upon reaching the genital ridge, the developmental pathway diverges based on genetic sex. In an XY gonad, PGCs become enclosed within **testis cords** to become **spermatogonia**. In an XX gonad, PGCs undergo mitosis to form clusters of **oogonia**.

SPERMATOGENESIS

Spermatogenesis is the continuous, high-yield production of motile spermatozoa within the **seminiferous tubules** of the testes, beginning at puberty and continuing throughout life. It occurs in a highly organized, wave-like pattern along the tubule length.

Stages of Spermatogenesis

1. Mitotic Proliferation (Spermatogonial Phase):

- **Location:** Basal compartment of seminiferous tubule.
- **Process:** **Spermatogonial stem cells (Type A_s)** undergo self-renewing mitotic divisions. Some of these committed cells differentiate into **Type A1-A4 spermatogonia**, then into **Type B spermatogonia**.
- **Regulation:** Tightly controlled by niche signals. **GDNF** promotes stem cell maintenance, while **SCF** promotes differentiation.


MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

2. Meiotic Phase (Spermatocyte Phase):

- **Location:** Transition from basal to adluminal compartment (crossing the **blood-testis barrier**).
- **Process:**
 - Type B spermatogonia enter a growth phase to become **primary spermatocytes (2n, 4c DNA)**.
 - **Meiosis I:** A reductional division. Homologous chromosomes pair, undergo crossing over (facilitated by the **synaptonemal complex**), and then separate. This yields two **secondary spermatocytes (n, 2c DNA)**.
 - **Meiosis II:** An equational division, where sister chromatids separate. Each secondary spermatocyte divides to produce two **spermatids (n, 1c DNA)**. Thus, one primary spermatocyte yields four genetically unique spermatids.
- **Regulation:** Initiated by **retinoic acid (RA)**, which induces expression of the **STRA8** gene, essential for meiotic entry.


Spermatogenesis



3. Spermiogenesis (Spermatid Differentiation):

- **Location:** Adluminal compartment, progressing towards the tubule lumen.
- **Process:** A complex morphological transformation of round, non-motile spermatids into streamlined spermatozoa, **without further cell division**. Key events include:
 - **Nuclear Condensation & Repackaging:** Histones are replaced by smaller, positively charged **protamines**, enabling extreme DNA compaction and inactivation of transcription.
 - **Acrosome Formation:** The Golgi apparatus forms a cap-like **acrosome** over the nucleus, filled with hydrolytic enzymes (hyaluronidase, acrosin) for egg penetration.
 - **Flagellum Assembly:** One centriole elongates to form the axoneme (9+2 microtubule arrangement) of the tail. The motor protein **dynein** generates motility.

- **Mitochondrial Sheath Formation:** Mitochondria spiral around the proximal part of the flagellum (midpiece) to provide ATP.
- **Cytoplasmic Shedding:** Excess cytoplasm is extruded as a **residual body**, which is phagocytosed by Sertoli cells.
- **Cytoplasmic Bridges:** Throughout meiosis and spermiogenesis, descendant cells remain connected by intercellular bridges, allowing synchronous development and sharing of gene products from the single X chromosome in XY cells.



4. Spermiation and Release:

Mature **spermatozoa** are released into the tubule lumen. They are still non-motile and undergo further functional maturation in the **epididymis** (gaining motility and fertilizing capacity) and final **capacitation** in the female reproductive tract.

Hormonal Regulation of Spermatogenesis

The **Hypothalamic-Pituitary-Gonadal (HPG) Axis** provides precise control:

- **Hypothalamus:** Secretes **Gonadotropin-Releasing Hormone (GnRH)** in pulses.
- **Anterior Pituitary:** GnRH stimulates release of:
 - **Follicle-Stimulating Hormone (FSH):** Binds to Sertoli cells, stimulating production of **Androgen-Binding Protein (ABP)** and other factors crucial for supporting meiosis and spermiogenesis.
 - **Luteinizing Hormone (LH):** Stimulates **Leydig cells** in the interstitium to produce **testosterone**.
- **Testosterone:** High local concentration within the tubule (maintained by ABP) is absolutely required for the completion of meiosis and spermiogenesis.
- **Feedback:** Testosterone inhibits GnRH and LH secretion. **Inhibin**, secreted by Sertoli cells, provides negative feedback specifically on FSH release.

Summary of Spermatogenesis

Stage	Cell Type (Ploidy)	Key Events	Duration (Human)
Mitosis	Spermatogonia (2n) → Primary Spermatocyte (2n)	Stem cell renewal, differentiation.	~16 days
Meiosis I	Primary Spermatocyte (2n) → Secondary Spermatocyte (n)	Synapsis, crossing over, homologous chromosome separation.	~24 days
Meiosis II	Secondary Spermatocyte (n) → Spermatid (n)	Separation of sister chromatids.	A few hours
Spermiogenesis	Spermatid (n) → Spermatozoon (n)	Nuclear condensation, acrosome & flagellum formation, cytoplasmic loss.	~24 days
Total		~74 days from spermatogonia to released sperm.	

OOGENESIS

Oogenesis is the discontinuous, yield-conserving production of large, nutrient-rich oocytes within the **ovarian follicles**. It begins in the fetal stage, involves prolonged meiotic arrests, and is completed only upon fertilization.

Stages of Oogenesis

1. Fetal Development (Mitotic Proliferation & Meiotic Arrest):

- **Process:** In the fetal ovary, PGCs divide mitotically to form **oogonia (2n)**. These enter meiosis I to become **primary oocytes (2n)** but arrest immediately at the **diplotene stage of Prophase I** (also called **dictyate arrest**).
- **Outcome:** A female is born with her lifetime supply of ~1-2 million primary oocytes, each encapsulated by a single layer of somatic **pregranulosa cells** to form a **primordial follicle**. No oogonia remain postnatally.


2. Follicular Growth (Recruitment & Cytoplasmic Maturation):

- **Process:** From puberty to menopause, cohorts of primordial follicles are periodically "recruited" to begin growth, stimulated by FSH. The primary oocyte undergoes enormous growth (e.g., a 100µm frog oocyte grows to ~1mm), accumulating:
 - **Yolk (Vitellogenin):** Proteins and lipids for early embryonic nutrition.
 - **Maternal mRNAs & tRNAs:** Stored in a translationally repressed state for use after fertilization.
 - **Ribosomes:** In massive numbers.
 - **Morphogenetic Determinants:** Proteins and mRNAs with asymmetric localization that will pattern the future embryo.
 - **Protective Components:** Pigments, antioxidants, defensive chemicals.
- **Folliculogenesis:** The surrounding somatic cells proliferate to form **granulosa cells**. A fluid-filled **antrum** forms, creating a **secondary** and then a **Graafian (tertiary) follicle**.

3. Resumption of Meiosis I and Ovulation:

- **Process:** Just before ovulation, the **LH surge** triggers the primary oocyte to resume and complete **Meiosis I**. This division is **highly asymmetric**:
 - One daughter cell receives almost all the cytoplasm → **secondary oocyte (n, 2c DNA)**.

- The other receives minimal cytoplasm → **first polar body** (which may or may not divide again).
- **Ovulation:** The Graafian follicle ruptures, releasing the **secondary oocyte**, which is arrested at **Metaphase of Meiosis II**. It is surrounded by the **zona pellucida** (glycoprotein coat) and a layer of granulosa cells (**corona radiata**).



4. Completion of Meiosis II (Triggered by Fertilization):

- **Process:** The secondary oocyte completes Meiosis II only upon fertilization by a sperm. Sperm penetration triggers a calcium signal that inactivates **Cytostatic Factor (CSF)**, allowing the cell cycle to proceed.
- **Outcome:** Another asymmetric division produces:
 - The large, mature **ovum** ($n, 1c$ DNA).
 - A tiny **second polar body**.
- The first polar body may also divide, resulting in a total of **three polar bodies** that degenerate.

Hormonal Regulation of Oogenesis

- **Follicular Phase:** FSH stimulates follicle growth and estradiol production. Rising estradiol thickens the endometrium and, at a critical threshold, triggers the **positive feedback LH surge**.
- **Ovulation:** The LH surge induces final oocyte maturation, resumption of Meiosis I, and follicle rupture.
- **Luteal Phase:** The ruptured follicle becomes the **corpus luteum**, secreting progesterone to prepare the endometrium. If no pregnancy, the corpus luteum regresses, hormone levels fall, and menstruation occurs.

Modern Insight: The Ovarian Reserve

The long-held dogma of a fixed number of oocytes is being challenged. Evidence in mice, primates, and humans suggests the presence of **female germline stem cells** in adult ovarian surface epithelium or cortex, capable of generating new oocytes. This has significant implications for reproductive biology and fertility preservation, though its physiological importance remains under investigation.

COMPARATIVE ANALYSIS: SPERMATOGENESIS vs. OOGENESIS

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091




Major Differences Between Spermatogenesis and Oogenesis

M
K
P
R
E
P
A
R
A
T
I
O
N
S


Feature	Spermatogenesis	Oogenesis
Site	Seminiferous tubules of testes .	Ovarian follicles .
Timing & Continuity	Continuous process from puberty to old age.	Cyclic & Discontinuous . Begins in fetus, arrested for decades, resumes cyclically from puberty to menopause.
Release	Millions per ejaculation.	Typically one secondary oocyte per menstrual cycle.
Meiotic Divisions	Equal cytokinesis.	Extremely Unequal cytokinesis, conserving cytoplasm in the functional gamete.
Product from 1 Germ Cell	4 functional, motile spermatozoa.	1 functional ovum + 2-3 polar bodies (which degenerate).
Arrests	No prolonged meiotic arrest.	Two arrests: Prophase I (from fetal stage to pre-ovulation) and Metaphase II (from ovulation to fertilization).
Gamete Size & Motility	Extremely small, motile , streamlined for delivery.	Very large (largest cell in body), non-motile , packed with nutrients and developmental machinery.
Cytoplasmic Content	Most cytoplasm eliminated; few mitochondria (paternally inherited mtDNA is degraded post-fertilization).	Massive cytoplasm retained; many mitochondria, ribosomes, stored mRNAs, yolk.
Stem Cell Population	Spermatogonial stem cells (SSCs) maintained throughout life, allowing continuous production.	Fixed pool of primary oocytes at birth (traditional view), though adult stem cells are a subject of ongoing research.
Hormonal Trigger for Meiosis	Retinoic acid (RA) induces STRA8 .	Luteinizing Hormone (LH) surge triggers resumption of Meiosis I; sperm-induced Ca²⁺ wave triggers completion of Meiosis II.

26. Reproduction and Development

Spermatogenesis



Oogenesis



Clinical and Biological Significance

- **Male Infertility:** Can result from defects at any spermatogenic stage (e.g., **azoospermia**, **oligospermia**), hormonal imbalances, or genetic issues like Y-chromosome microdeletions.
- **Female Fertility & Maternal Age:** The **ovarian reserve** declines with age. More critically, the prolonged dictyate arrest makes oocytes susceptible to age-related damage, particularly to the **cohesin** complexes that hold homologous chromosomes together. This is the primary reason for the dramatic increase in **aneuploidy** (e.g., Trisomy 21/Down syndrome) with advanced maternal age.
- **Conservation of Resources:** Oogenesis invests heavily in one gamete to support early development, while spermatogenesis prioritizes quantity and motility to ensure fertilization success. This anisogamy is a fundamental aspect of sexual reproduction.
- **Epigenetic Reprogramming:** Both processes involve extensive epigenetic remodeling (DNA methylation, histone modification) that is crucial for genomic imprinting and normal embryonic development. Errors here can lead to transgenerational epigenetic disorders.

FERTILIZATION

Fertilization is the fundamental biological process involving the fusion of two highly specialized **haploid gametes**—the **spermatozoon** (male) and the **oocyte/ovum** (female)—to form a single **diploid zygote**.

This event accomplishes two primary, interconnected functions:

1. **Sexual Reproduction:** It combines genetic material from two parents, generating a genetically unique individual and promoting genetic diversity within a population.
2. **Activation of Development:** It triggers the resumption of the arrested metabolism and cell cycle of the mature oocyte, initiating the complex program of embryogenesis.

The process is a tightly coordinated sequence of molecular and cellular events designed to ensure **species specificity** and prevent lethal **polyspermy** (entry of multiple sperm). Mechanisms vary between **external fertilizers** (e.g., sea urchins, many fish) and **internal fertilizers** (e.g., mammals, birds), but core principles are conserved.

MK PREPARATIONS: Let's Make It Happen

+92 333 2605045, +92 342 4470091

Structural Preparation: The Specialized Gametes

A. The Spermatozoon: A Cell Designed for Delivery

The sperm is a minimalist, motile cell optimized for transporting the paternal genome to the egg.

Anatomy and Functional Compartments:

1. Head:

- **Nucleus:** Contains a tightly compacted, transcriptionally inactive haploid genome. **Protamines** replace histones for extreme DNA condensation.
- **Acrosome:** A Golgi-derived, membrane-bound vesicle capping the nucleus. It contains a cocktail of **hydrolytic enzymes** (e.g., **hyaluronidase**, **acrosin**, proteases) essential for penetrating the egg's outer investments.

2. Midpiece (Neck):



- Packed with **spirally arranged mitochondria** that generate ATP via oxidative phosphorylation to power flagellar motility.
- Contains the **centriole(s)** that will nucleate microtubules for pronuclear migration and first cleavage spindle.

3. Tail (Flagellum):

- The propulsive apparatus with a canonical "**9+2**" **axoneme** of microtubules.
- Motility is generated by the motor protein **dynein**, which hydrolyzes ATP to create sliding forces between microtubule doublets.

Maturation Events Post-Testis:

- **Epididymal Maturation:** Sperm gain forward motility and fertilizing capacity.
- **Capacitation (in mammals):** A final maturation step occurring in the female reproductive tract. It involves cholesterol efflux from the sperm plasma membrane, leading to increased membrane fluidity, hyperactivated motility, and preparation for the **acrosome reaction**.



B. The Oocyte/Ovum: A Cell Designed for Reception and Support


The egg is a large, metabolically quiescent cell packed with reserves for early development.

Cytoplasmic Stores (Deutoplasm):

- **Nutrients:** Yolk proteins (vitellogenin), lipids, and glycogen.
- **Protein Synthesis Machinery:** Ribosomes, tRNAs, and **maternally-loaded mRNAs** stored in a translationally repressed state.
- **Morphogenetic Determinants:** Transcription factors and signaling molecules (e.g., bicoid in *Drosophila*) often localized to specific regions to establish future body axes.
- **Protective Substances:** Antioxidants, DNA repair enzymes, antimicrobial compounds.

M
K

P
R
E
P
A
R
A
T
I
O
N
S



Protective Membranes and Envelopes:

1. **Plasma Membrane:** Contains species-specific receptors for sperm binding.
2. **Extracellular Coats:**
 - **Vitelline Envelope:** In many invertebrates (e.g., sea urchins).
 - **Zona Pellucida (ZP):** A thick glycoprotein matrix in mammals (composed of ZP1, ZP2, ZP3, ZP4).
 - **Jelly Coat:** An outer, non-cellular layer in many external fertilizers containing chemoattractants.
3. **Cortical Region:** A gel-like layer just beneath the plasma membrane, rich in globular actin and membrane-bound **cortical granules**. These granules contain enzymes and mucopolysaccharides crucial for the **slow block to polyspermy**.

The Sequential Stages of Fertilization


Stage 1: Sperm Attraction and Activation (Chemotaxis)

- **Purpose:** To guide sperm toward the egg, increasing the probability of encounter.
- **Mechanism:** Eggs release species-specific **chemoattractants**, creating a chemical gradient.
- **Examples:**
 - **Sea Urchin:** The egg jelly secretes small peptides like **resact** (in *Arbacia*). Resact binds to receptor-guanylyl cyclases on the sperm flagellum, increasing intracellular cGMP, opening CatSper calcium channels, and inducing directed, **chemotactic** motility.

- **Mammals:** A multi-step guidance system:
 1. **Rheotaxis:** Sperm swim against the flow of fluid in the oviduct.
 2. **Thermotaxis:** Sperm are guided by a temperature gradient toward the warmer ampullary-isthmic junction.
 3. **Chemotaxis:** Capacitated sperm are attracted by factors like **progesterone** secreted by the **cumulus cells** surrounding the oocyte.

Stage 2: The Acrosome Reaction

- **Purpose:** A triggered exocytosis that releases acrosomal enzymes and exposes proteins necessary for binding to and penetrating the egg's outer layers.
- **Induction:**
 - **Sea Urchin:** Sulfated polysaccharides (fucose sulfate) in the egg jelly bind to sperm receptors, causing membrane depolarization and an influx of Ca^{2+} and Na^+ , which triggers the reaction.
 - **Mammals:** Binding of sperm to the **Zona Pellucida (ZP)**, specifically to glycoproteins like **ZP3** (in the classical model) or **ZP2** (in newer models), induces the reaction.
- **Events:**
 1. Fusion of the outer acrosomal membrane with the sperm plasma membrane.
 2. Formation of hybrid vesicles and release of soluble enzymes to create a path.
 3. In many species (e.g., sea urchin), polymerization of globular actin forms a finger-like **acrosomal process**, which extends and presents **bindin** (or equivalent) for species-specific adhesion.



Stage 3: Species-Specific Binding

- **Purpose:** To ensure only conspecific sperm can fertilize the egg, a key mechanism of **reproductive isolation**.

- **Mechanism:** A molecular "lock-and-key" interaction between sperm surface proteins and receptors on the egg coat.
 - **Sea Urchin: Bindin** on the acrosomal process binds to **bindin receptors** (glycoprotein complexes like EBR1) on the vitelline envelope.
 - **Mammals:** Acrosome-reacted sperm bind to the ZP. Current evidence points to binding of sperm proteins to a cleaved form of **ZP2** as the primary interaction for maintaining sperm binding post-acrosome reaction.

Stage 4: Sperm-Egg Membrane Fusion

- **Purpose:** To allow the sperm nucleus, centriole, and activating factors to enter the egg cytoplasm.
- **Fusogenic Proteins:** Specific proteins mediate the merger of the two plasma membranes.
 - **Mammals:** The sperm protein **IZUMO1** (exposed post-acrosome reaction) interacts with its egg receptor **JUNO (FOLR4)**. This interaction recruits tetraspanin proteins like **CD9** and **CD81** in the egg membrane, which facilitate membrane fusion.

Stage 5: Blocks to Polyspermy


Polyspermy is lethal, leading to aberrant ploidy, multipolar spindles, and developmental arrest. Successful species employ sequential blocks.

1. Fast Block (Electrical):

- **Mechanism:** Sperm-egg fusion causes an immediate influx of Na^+ into the egg, changing the membrane potential from negative (≈ -70 mV) to positive ($\approx +20$ mV). This **depolarization** prevents the fusion of additional sperm.
- **Occurrence:** Well-documented in external fertilizers with fast, synchronous fertilization (sea urchins, frogs).
- **Duration:** Transient (≈ 1 minute), serving as a rapid but temporary barrier until the slow block is established.

2. Slow Block (Cortical Granule Reaction):

- **Mechanism:** A permanent, physical/chemical barrier.
 - Sperm fusion triggers a wave of **calcium ions (Ca^{2+})** released from the egg's endoplasmic reticulum (via **IP_3** receptor channels).
 - The Ca^{2+} wave causes **cortical granules** to fuse with the egg plasma membrane and exocytose their contents into the perivitelline space.
- **Consequences:**
 - **Sea Urchin:** Enzymes clip sperm receptors (e.g., bindin receptors), mucopolysaccharides cause osmotic swelling, and cross-linking enzymes harden the elevated vitelline envelope into an impenetrable **fertilization envelope**.
 - **Mammals:** Cortical granule enzymes (e.g., **ovastacin**, a zinc metalloprotease) cleave **ZP2**, making it unable to bind sperm. The **JUNO** receptor is also shed from the egg membrane. This is the **primary block** in mammals.





M
K
P
R
E
P
A
R
A
T
I
O
N
S

Stage 6: Activation of Egg Metabolism

The sperm-induced Ca^{2+} signal is the master trigger for reactivating the dormant egg.

- **Completion of Meiosis:** In mammals, Ca^{2+} oscillations inactivate **MAP kinase** and **Cyclostatic Factor (CSF)**, allowing the secondary oocyte to complete **Meiosis II**, form the female pronucleus, and extrude the second polar body.
- **Resumption of Cell Cycle:** Ca^{2+} activates **calmodulin-dependent kinase II (CaMKII)**, which leads to the degradation of cyclin B and other inhibitors, allowing the cell cycle to progress from its **Metaphase II arrest**.
- **Cortical Granule Exocytosis:** As described above.
- **Metabolic Activation:** Increased cytoplasmic pH and calcium activate translation, leading to a **burst of protein synthesis** from stored maternal mRNAs (e.g., for cyclins, histones). DNA replication is initiated.

Post-Fusion Events: Formation of the Zygote

1. Sperm Nuclear Decondensation and Pronuclear Formation

- The tightly packaged sperm chromatin is unwound. In mammals, protamines are actively removed and replaced by histones from the egg cytoplasm.
- The remodeled sperm nucleus swells to form the **male pronucleus**.

2. Pronuclear Migration and Syngamy

- The **sperm-derived centriole** nucleates microtubules, forming a **sperm aster** that radiates through the egg cytoplasm.
- The female pronucleus is captured by these microtubules, and the two pronuclei migrate toward each other, meeting near the cell center.
- **Syngamy** (genome union):
 - **Sea Urchin/Frog:** Pronuclear membranes fuse to create a single diploid **zygote nucleus**.
 - **Mammals:** Pronuclear membranes dissolve without full fusion. The parental chromosomes align on a common **mitotic spindle** derived from the sperm centriole. A true diploid nucleus is first seen after the first cleavage division.

3. Cytoskeletal and Organelle Inheritance

- **Centrosome:** Typically paternally inherited (from the sperm centriole) in most animals, establishing the mitotic poles for cleavage.
- **Mitochondria: Exclusively maternal inheritance.** Paternal mitochondria are actively tagged with ubiquitin and degraded by the egg's autophagy machinery shortly after fertilization.

Comparison of Fertilization Events in Sea Urchins vs. Mammals

Event	Sea Urchin (External Fertilizer)	Mammal (Internal Fertilizer)
Sperm Maturation	Mature upon spawning; activated by seawater pH.	Requires capacitation in female tract (membrane remodeling, protein phosphorylation).
Egg at Fertilization	Meiosis complete; haploid female pronucleus present.	Arrested at Metaphase II ; meiosis completed after sperm entry.
Chemoattraction	Strong, using peptides like resact/speract .	Modest; involves thermotaxis, rheotaxis, chemotaxis (e.g., progesterone).
Acrosome Reaction Site	Triggered by egg jelly at a distance from the egg.	Occurs on the surface of the zona pellucida.



MK PREPARATIONS



M
K

P
R
E
P
A
R
A
T
I
O
N
S

Primary Sperm-Binding Protein	Bindin on sperm binds to receptors on vitelline envelope.	Acrosome-reacted sperm bind to ZP2 on zona pellucida.
Fast Block	Yes. Na ⁺ -dependent membrane depolarization.	Likely absent or minor. Slow block is primary.
Slow Block Mechanism	Cortical granule exocytosis forms a fertilization envelope .	Cortical granules modify the zona pellucida (ZP2 clipping).
Source of Activating Ca ²⁺	Released from egg ER via IP ₃ produced by egg PLCγ .	Released from egg ER via IP ₃ produced by sperm-derived PLCζ .
Pronuclear Behavior	Pronuclei migrate and fuse before first mitosis.	Pronuclei remain separate; chromosomes align on common spindle.
Centrosome Inheritance	Derived from sperm centriole .	Derived from sperm centriole (in most).
Mitochondrial Inheritance	Exclusively maternal.	Exclusively maternal.

Clinical and Applied Significance

- **Infertility:** Defects in any step (sperm motility, acrosome reaction, sperm-egg binding, fusion, egg activation) can cause infertility. Understanding these steps underpins **Assisted Reproductive Technologies (ART)** like **ICSI**.
- **Contraception:** Targeting key steps (e.g., sperm capacitation, acrosome reaction, zona pellucida binding) is a strategy for developing non-hormonal contraceptives.
- **Prevention of Polyspermy:** In IVF, ensuring monospermic fertilization is critical. Techniques like **intracytoplasmic sperm injection (ICSI)** bypass all natural barriers but must be performed with precision.
- **Pronuclear Transfer:** For preventing mitochondrial diseases, techniques involve transferring the pronuclei from a fertilized egg with defective mitochondria into an enucleated donor egg with healthy mitochondria.

Pregnancy and Human Development

Fertilization

Fertilization is the process by which a sperm fuses with an oocyte (secondary oocyte) to form a **zygote**. This typically occurs in the **ampulla of the oviduct** (fallopian tube), the widest section near the ovary.

Key Steps in Fertilization:

1. **Sperm Capacitation:** Sperm undergo final maturation within the female reproductive tract, gaining the ability to fertilize the egg.
2. **Acrosomal Reaction:** Upon contact with the **zona pellucida** (glycoprotein layer surrounding the oocyte), the sperm's **acrosome** releases hydrolytic enzymes (hyaluronidase, acrosin) that digest the zona pellucida.
3. **Sperm Penetration:** A single sperm penetrates the zona pellucida and fuses with the oocyte's plasma membrane.



4. **Cortical Reaction:** Fusion triggers the **cortical reaction** – cortical granules in the oocyte cytoplasm release enzymes that modify the zona pellucida, making it impenetrable to other sperm. This **blocks polyspermy** (fertilization by multiple sperm).
5. **Completion of Meiosis II:** The sperm entry activates the oocyte to complete **meiosis II**, forming the mature **ovum** and a second polar body.
6. **Pronuclei Formation:** The sperm nucleus swells to form the **male pronucleus**; the ovum nucleus becomes the **female pronucleus**.
7. **Syngamy:** The pronuclei fuse, combining their chromosomes to form a diploid **zygote**.

Cleavage

Cleavage is a series of rapid **mitotic divisions** of the zygote without overall growth. The cells produced are called **blastomeres**.

Stages:

- **Day 1-3:** Zygote divides into 2, then 4, then 8 cells, forming a **morula** (solid ball of 16+ cells) by day 4.
- **Day 4-5:** The morula develops into a **blastocyst** as it enters the uterus. The blastocyst has:
 - **Trophoblast:** Outer cell layer that will form the placenta and extra-embryonic membranes.
 - **Inner Cell Mass (Embryoblast):** Cluster of cells that will develop into the embryo.
 - **Blastocoel:** Fluid-filled cavity.

Implantation

Implantation is the attachment and embedding of the blastocyst into the **endometrium** (uterine lining), occurring approximately **6–7 days after fertilization**.

Process:

1. **Hatching:** The blastocyst "hatches" from the zona pellucida.
2. **Attachment:** The **trophoblast** adheres to the endometrium.
3. **Invasion:** Trophoblast cells proliferate and invade the endometrium, forming two layers:
 - **Cytotrophoblast** (inner cellular layer).
 - **Syncytiotrophoblast** (outer multinucleated layer that erodes maternal tissues to establish nutrient exchange).
4. **Decidual Reaction:** The endometrium undergoes changes (increased vascularity, edema) to form the **decidua**, which supports implantation.

By day 7–10, the blastocyst is fully embedded. The **inner cell mass** differentiates into the **bilaminar germ disc** (epiblast and hypoblast), the precursor to the embryo.

Placenta Formation and Function

Placental Structure

The **placenta** is a temporary organ formed from both fetal and maternal tissues.

Fetal Components:

- **Chorionic villi:** Finger-like projections of the **chorion** (derived from trophoblast) containing fetal capillaries.
- **Chorionic plate:** Fetal side of the placenta.
- **Umbilical cord:** Connects fetus to placenta; contains **two umbilical arteries** (carry deoxygenated blood from fetus) and **one umbilical vein** (carries oxygenated blood to fetus).


Maternal Components:

- **Decidua basalis:** The portion of the endometrium underlying the implantation site.
- **Maternal blood pools (lacunae):** Spaces where maternal blood bathes the chorionic villi.

Important: Maternal and fetal blood **do not mix**; exchange occurs across the **placental barrier** (syncytiotrophoblast, connective tissue, and fetal capillary endothelium).

M
K

P
R
E
P
A
R
A
T
I
O
N
S



Placental Functions

A. Exchange & Transport:

- **Passive Diffusion:** Oxygen, carbon dioxide, water, electrolytes.
- **Facilitated Diffusion:** Glucose (via GLUT1 transporters).
- **Active Transport:** Amino acids, ions (e.g., Ca^{2+} , Fe^{2+}).
- **Pinocytosis:** Maternal antibodies (IgG) for passive immunity.
- **Waste Removal:** Urea, creatinine, bilirubin transferred to maternal blood.

B. Endocrine Secretion:

- **hCG (Human Chorionic Gonadotropin):** Produced by syncytiotrophoblast; maintains the **corpus luteum** for progesterone secretion until the placenta takes over (~week 8–10). Basis for pregnancy tests.
- **Progesterone:** Maintains endometrium, suppresses uterine contractions, prevents menstruation, prepares mammary glands.
- **Estrogens (primarily estriol):** Stimulate uterine growth, increase blood flow, promote breast duct development.
- **hPL (Human Placental Lactogen):** Modulates maternal metabolism to ensure fetal nutrient supply; has growth hormone-like and lactogenic effects.
- **Relaxin:** Softens cervix and relaxes pelvic ligaments for parturition.

Developmental Timeline (Gestation ~40 weeks)

Gestation is calculated as **40 weeks from the last menstrual period (LMP)** or **38 weeks from fertilization**.

First Trimester (Weeks 1–12)

- **Week 1–2:** Fertilization, cleavage, implantation.
- **Week 3: Gastrulation** – formation of three germ layers: ectoderm, mesoderm, endoderm.
- **Week 4–8: Organogenesis** – major organs form from germ layers.
 - **Ectoderm:** Nervous system, epidermis, sensory organs.
 - **Mesoderm:** Muscles, bones, circulatory system, kidneys.
 - **Endoderm:** Lining of digestive and respiratory tracts, glands.
- **Week 8:** Embryo becomes a **fetus**; all major organs are present in rudimentary form.
- **Vulnerability:** This period is highly sensitive to **teratogens** (alcohol, drugs, infections, radiation) that can cause congenital malformations.



Second Trimester (Weeks 13–26)

- **Growth:** Rapid increase in length; fetal movements felt by mother (**quickening** ~week 18–20).
- **Differentiation:** Sex distinguishable by ultrasound; hair, nails form.
- **Organ Maturation:** Bones ossify; urine production begins; surfactant production in lungs starts.
- **Viability:** By ~24 weeks, survival outside uterus may be possible with intensive care.

Third Trimester (Weeks 27–40)

- **Final Growth:** Fetus gains weight (~50% of birth weight in last 10 weeks).
- **Maturation:** Lungs complete surfactant production; nervous system develops (myelination); fat deposits form for thermoregulation.
- **Positioning:** Fetus typically moves into **cephalic presentation** (head down) by week 36.
- **Immunity:** Maternal antibodies (IgG) are actively transported to fetus, providing passive immunity for the first few months after birth.

Parturition (Birth)

Parturition is the process of childbirth, typically occurring ~40 weeks after LMP. It is initiated by a hormonal cascade.

Hormonal Triggers:

1. **Fetal Cortisol:** Rising levels from the fetal adrenal cortex stimulate the placenta to convert progesterone to **estrogens**.
2. **Estrogen Surge:** Increases expression of **oxytocin receptors** on uterine myometrium and promotes formation of **prostaglandins** (PGE₂, PGF_{2α}) that stimulate contractions.
3. **Progesterone Withdrawal:** The relative decrease in progesterone (compared to estrogen) removes its quieting effect on the uterus.
4. **Oxytocin Release:** From the maternal posterior pituitary; stimulated by cervical stretching (**Ferguson reflex**) and positive feedback.

Stages of Labor:

Stage 1 – Dilation:

- Cervix effaces (thins) and dilates (opens) to ~10 cm.
- Can last 6–12 hours (longer in first pregnancies).
- Regular contractions increase in frequency and intensity.

Stage 2 – Expulsion:

- From full dilation to delivery of the baby.
- Mother bears down with contractions.
- Typically lasts up to a few hours.

Stage 3 – Placental Delivery:

- Uterine contractions detach and expel the placenta and membranes ("afterbirth") within 30 minutes.
- Important to ensure complete expulsion to prevent hemorrhage.


Positive Feedback Mechanism:

Contractions → Cervical stretching → Oxytocin release → Stronger contractions → More stretching → Continued until delivery.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

Onset of labour



Flexion



Internal rotation of head




Extension



External rotation of head



Uterus immediately after birth



Lactation

Mammary Gland Preparation:

- During pregnancy, **estrogen** and **progesterone** stimulate duct and alveolar growth.
- **Prolactin** levels rise but milk production is inhibited by high progesterone.

Postpartum Lactation:

- After delivery, progesterone drops, allowing **prolactin** to initiate **lactogenesis** (milk production).
- **Colostrum** is produced first (days 1–3): yellowish fluid rich in **antibodies (IgA)**, proteins, and vitamins, but lower in fat and lactose. Provides passive immunity and laxative effect to clear meconium.
- **Mature milk** appears by ~day 4: higher in lactose and fat.

Milk Ejection Reflex (Let-down):

- Stimulated by **oxytocin** released in response to suckling (neuroendocrine reflex).
- Oxytocin causes myoepithelial cells around alveoli to contract, ejecting milk into ducts.

Maintenance of Lactation:

- **Suckling** stimulates prolactin secretion (maintains milk production) and oxytocin release (ejection).
- Emotional stress can inhibit oxytocin release, hindering milk let-down.


Benefits of Breastfeeding:

- **Infant:** Optimal nutrition, immune protection, bonding, reduced risk of allergies, infections, and chronic diseases.
- **Mother:** Promotes uterine involution, delays ovulation (natural contraception), reduces risk of breast/ovarian cancer, and enhances bonding.

EXTRAEMBRYONIC MEMBRANES IN TERRESTRIAL VERTEBRATES

These membranes support embryonic development on land by providing protection, nutrition, gas exchange, and waste storage. They are derived from the germ layers but are not part of the embryo proper.

Membrane	Germ Layer Origin	Primary Function in Birds/Reptiles	Function in Mammals
Chorion	Ectoderm & Mesoderm	Outermost membrane; major site of gas exchange with the environment.	Contributes to the placenta ; involved in gas and nutrient exchange with maternal blood.
Amnion	Ectoderm & Mesoderm	Encloses embryo in amniotic fluid ; prevents desiccation, cushions against shock.	Same essential functions. Amniocentesis samples this fluid for prenatal diagnosis.
Allantois	Endoderm & Mesoderm	Stores nitrogenous wastes; its blood vessels become part of the chorionic circulation for gas exchange.	Small; its blood vessels contribute to umbilical circulation. Waste is handled by the placenta.
Yolk Sac	Endoderm & Mesoderm	Encloses and digests yolk, making nutrients available to the embryo.	Vestigial in placental mammals (little yolk), but an important early site of blood cell formation .





M
K

P
R
E
P
A
R
A
T
I
O
N
S

Mammalian Placenta and Development

Placental mammals retain the extraembryonic membranes but modify their functions for internal development.

- **Implantation:** The **blastocyst** implants into the uterine endometrium. The **trophoblast** forms the chorion and **chorionic villi**, which interdigitate with maternal tissue to form the placenta.
- **Placental Function:** Facilitates nutrient/waste exchange and gas exchange. It evades maternal immune rejection via specialized proteins.
- **Developmental Periods:**
 - **Germinal Period (First 2 weeks):** Cleavage, implantation, resistant to teratogens.
 - **Embryonic Period (Weeks 3-8):** Organogenesis; highly sensitive to teratogens.
 - **Fetal Period (Week 9 to birth):** Growth and continued differentiation.

Organogenesis: Derivatives of Germ Layers

All adult structures derive from one of the three embryonic germ layers.

Major Derivatives of the Primary Germ Layers in Vertebrates

Ectoderm	Mesoderm	Endoderm
<ul style="list-style-type: none"> • Epidermis, hair, nails, skin glands • Lining of mouth, enamel of teeth • Lens & cornea of eye, inner ear • Neural Tube: Brain, spinal cord, motor nerves • Neural Crest: Sensory nerves, adrenal medulla, melanocytes, craniofacial cartilage/bone, dentine 	<ul style="list-style-type: none"> • Notochord • Dermis of skin • All muscle (skeletal, cardiac, smooth) • Bone & cartilage (except neurocranium) • Circulatory system (heart, blood vessels, blood cells) • Kidneys, gonads, reproductive ducts • Lining of body cavities (peritoneum) 	<ul style="list-style-type: none"> • Epithelial lining of digestive tract and respiratory tract • Parenchyma of liver and pancreas • Pharyngeal pouches → tonsils, thymus, parathyroid, thyroid glands • Lining of urethra, urinary bladder

Reproductive Health and Technologies

Disorders and Infertility

- **Infertility Causes:**
 - **Male:** Azoospermia, oligospermia, duct blockages.
 - **Female:** Anovulation, blocked oviducts (e.g., from **Pelvic Inflammatory Disease, PID**), endometriosis, uterine fibroids.
- **Endocrine Disrupting Contaminants (EDCs):** Chemicals interfering with hormone action.

Assisted Reproductive Technology (ART)

- **In Vitro Fertilization (IVF):** Eggs fertilized in lab, embryos transferred to uterus.
- **Intracytoplasmic Sperm Injection (ICSI):** Single sperm injected into oocyte.
- **Gamete Intrafallopian Transfer (GIFT):** Gametes transferred into oviduct.

Sexually Transmitted Diseases (STDs)

Infections transmitted through sexual contact.

Disease	Causative Agent	Key Notes
Gonorrhea	<i>Neisseria gonorrhoeae</i>	Can cause PID, infertility. Curable with antibiotics.



MK PREPARATIONS



Syphilis	<i>Treponema pallidum</i>	Progresses in stages; can cause severe long-term damage. Curable with penicillin.
Genital Herpes	Herpes Simplex Virus (HSV-2)	Recurrent outbreaks; no cure ; antivirals manage symptoms.
HIV/AIDS	Human Immunodeficiency Virus (HIV)	Destroys Helper T-cells (CD4+). Managed with Antiretroviral Therapy (ART) ; no cure .
HPV	Human Papillomavirus	High-risk strains cause cervical cancer. Vaccination available .

Contraception (Birth Control)

Method	Category	Mode of Action	Effectiveness
Condoms	Barrier	Block sperm entry.	Moderate-High (also prevents STIs).
Oral Contraceptives	Hormonal	Inhibit ovulation, thicken cervical mucus.	Very High (with perfect use).
IUD	Device/Hormonal	Spermicidal or hormonal effects to prevent implantation.	Very High.
Sterilization	Surgical	Vasectomy or Tubal Ligation blocks gamete transport.	Permanent, Very High.
Emergency Contraception	Hormonal	Delays ovulation or disrupts implantation.	Moderate.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development

PREPARATIONS
LET'S MAKE IT HAPPEN



Reproduction & Development: One Liner

- **Developmental biology** is the study of how organisms grow and develop from a single fertilized egg into complex multicellular structures.
- It encompasses **embryology** (development from fertilization to birth) and post-embryonic processes like **metamorphosis**, regeneration, and aging.
- A central question is how a single **zygote** with one genome can produce over **250 different human cell types**.
- **Morphogenesis** is the process by which cells organize into tissues and organs through coordinated division, migration, shape change, and death.
- Development is precisely regulated to ensure organs and limbs grow to the correct size, shape, and symmetry.
- **Germ cells** (sperm and egg) are set aside early in development to pass genetic information to the next generation.
- Some animals, like salamanders, can **regenerate** limbs, while mammals rely on **stem cells** for more limited repair.
- Environmental factors like **temperature** can influence development, as seen in **Temperature-Dependent Sex Determination (TSD)** in many reptiles.
- Evolutionary changes in developmental processes, such as alterations in gene regulation, lead to new body forms (e.g., evolution of the one-toed horse).
- Historically, **epigenesis** (development from scratch) was opposed to **preformationism** (performed miniature homunculus); the modern synthesis combines genetic instructions with epigenetic construction.
- The typical animal life cycle includes **fertilization, cleavage, gastrulation, organogenesis**, and sometimes **metamorphosis**.
- **Gastrulation** forms the three primary **germ layers: ectoderm, mesoderm, and endoderm**.
- **Von Baer's laws** state that general embryonic features appear before specialized ones, and embryos of different species diverge early and do not pass through adult stages of their ancestors.
- Cleavage patterns are influenced by yolk amount: **holoblastic** (complete) in eggs with little yolk (e.g., mammals, sea urchins) and **meroblastic** (incomplete) in yolk-rich eggs (e.g., birds, reptiles).
- **Isolecithal** eggs have sparse, even yolk and undergo equal holoblastic cleavage.
- **Mesolecithal** eggs have moderate yolk concentrated at the vegetal pole and undergo unequal holoblastic cleavage (e.g., amphibians).
- **Discoidal** cleavage is a meroblastic pattern where cleavage is restricted to a small disc of cytoplasm (e.g., birds, reptiles).
- **Superficial** cleavage is a meroblastic pattern where cleavage occurs in peripheral cytoplasm surrounding a central yolk mass (e.g., most insects).
- **Gastrulation movements** include **invagination, involution, ingression, delamination, and epiboly**.
- **Cell behaviors** during morphogenesis include changes in division, shape, migration, growth, death (**apoptosis**), and adhesion/secretions.
- **Fate mapping** uses dyes, fluorescent markers, or genetic labels (e.g., **GFP**) to trace which embryonic cells become which adult structures.
- Embryonic similarities between species indicate common ancestry, supporting **evolutionary embryology**.
- **Homologous structures** (e.g., human arm, bird wing) share a common evolutionary origin, while **analogous structures** (e.g., bird wing, insect wing) have similar functions but different origins.

M
K

P
R
E
P
A
R
A
T
I
O
N
S

26. Reproduction and Development



- **Evo-Devo** (Evolutionary Developmental Biology) studies how changes in developmental genes and processes drive evolutionary morphological change.
- **Teratogens** (e.g., alcohol, thalidomide, radiation) can cause birth defects during critical periods of organogenesis.
- Understanding normal development is essential for diagnosing **congenital disorders** and advancing **regenerative medicine**.
- **Sexual reproduction** combines genetic material from two parents via **meiosis** and fertilization, increasing genetic diversity.
- **Primary sex determination** refers to the development of gonads (testes or ovaries).
- In mammals, primary sex determination is chromosomal: **XX** = female, **XY** = male.
- The **SRY gene** on the Y chromosome triggers testis development by activating **SOX9**.
- In the absence of SRY, the ovarian pathway is activated through actions of **WNT4**, **RSPO1**, and **FOXL2**.
- **Secondary sex determination** is the development of genitalia and secondary sexual characteristics, driven by gonadal hormones.
- In males, **testosterone** and **Anti-Müllerian Hormone (AMH)** drive secondary sex determination.
- In females, the default pathway occurs without testicular hormones; **estrogen** promotes female reproductive tract development.
- **Disorders of Sexual Development (DSD)** include **Androgen Insensitivity Syndrome (AIS)**, **5 α -Reductase Deficiency**, and **Congenital Adrenal Hyperplasia (CAH)**.
- In **Drosophila**, sex is determined by the **X:A ratio** (number of X chromosomes to sets of autosomes), not by hormones, via a splicing cascade involving **Sex-lethal (Sxl)** and **doublesex (dsx)**.
- Some reptiles and fish use **Environmental Sex Determination (ESD)**, where incubation temperature determines sex.
- Common ESD patterns include: low temperature = males, high = females (Pattern Ia, turtles); the opposite (Pattern Ib, some lizards); and extremes produce one sex, intermediate the other (Pattern II, alligators).
- The molecular basis of ESD involves temperature-sensitive enzymes like **aromatase** (converts androgens to estrogens) and expression of genes like **DMRT1** and **SOX9**.
- **Gametogenesis** is the process by which diploid **Primordial Germ Cells (PGCs)** undergo meiosis and differentiation to form haploid gametes.
- PGCs are specified early in development and migrate to the gonadal ridges guided by chemotactic signals like **SDF1/CXCR4**.
- **Spermatogenesis** occurs in the **seminiferous tubules** of the testes and is continuous from puberty.
- **Oogenesis** begins in the fetal ovary, involves long meiotic arrests, and is cyclic from puberty to menopause.
- **Spermatogenesis** produces four functional spermatozoa from one germ cell through two meiotic divisions.
- **Oogenesis** produces one functional ovum and two or three polar bodies from one germ cell through highly unequal cytokinesis.
- Spermatogenesis has **no prolonged meiotic arrest**, while oogenesis has two arrests: at **Prophase I** (from fetal stage to ovulation) and **Metaphase II** (from ovulation to fertilization).
- In spermatogenesis, stem cells (**spermatogonial stem cells**) are maintained throughout life.
- Traditionally, females are born with a fixed pool of primary oocytes, though the existence of adult female germline stem cells is debated.



- **Spermiogenesis** is the morphological transformation of round spermatids into streamlined spermatozoa, involving nuclear condensation, acrosome formation, flagellum assembly, and cytoplasmic loss.
- In oogenesis, the primary oocyte undergoes enormous cytoplasmic growth, accumulating yolk, ribosomes, maternal mRNAs, and morphogenetic determinants.
- The **LH surge** triggers the resumption and completion of **Meiosis I** in the oocyte, leading to ovulation of a **secondary oocyte**.
- Fertilization triggers the completion of **Meiosis II** in the oocyte via a sperm-induced calcium signal.
- **Meiosis** generates genetic diversity through **independent assortment** of chromosomes and **crossing over** between homologous chromosomes.
- **Maternal age** increases the risk of **aneuploidy** (e.g., Down syndrome/Trisomy 21) due to age-related degradation of **cohesin** proteins holding chromosomes together.
- **Retinoic acid (RA)** and the **STRA8** gene are crucial for initiating meiosis in spermatogenesis.
- **Fertilization** is the fusion of sperm and egg to form a diploid **zygote**, restoring diploidy and initiating development.
- Sperm are small, motile cells with an **acrosome** containing digestive enzymes, a midpiece with mitochondria, and a flagellum for movement.
- Eggs are large, non-motile cells storing nutrients, ribosomes, maternal mRNAs, and morphogenetic factors, surrounded by protective coats like the **zona pellucida** (mammals) or vitelline envelope.
- **Capacitation** is a final maturation step for mammalian sperm in the female reproductive tract, involving membrane remodeling.
- Steps of fertilization include **sperm attraction** (chemotaxis/thermotaxis/rheotaxis), **acrosome reaction**, **species-specific binding**, **sperm-egg membrane fusion**, and **blocks to polyspermy**.
- The **fast block to polyspermy** is a rapid electrical change (membrane depolarization) in some species like sea urchins.
- The **slow block to polyspermy** is the **cortical granule reaction**, a physical/chemical modification of the egg's envelopes (e.g., fertilization envelope, ZP2 clipping).
- Sperm-egg fusion triggers a **calcium wave** (**Ca²⁺ oscillations** in mammals) that activates the egg, resumes meiosis, and initiates embryonic cycles.
- In mammals, the sperm contributes **PLC ζ** (phospholipase C zeta) to trigger the calcium oscillations.
- **Pronuclei** from the sperm and egg form and migrate toward each other; their genomes unite in a process called **syngamy**.
- **Polyspermy** (entry of multiple sperm) is typically lethal and is prevented by fast and slow blocks.
- Gamete recognition proteins like **bindin** (sea urchin) and the **Izumo1-Juno** interaction (mammals) evolve rapidly, contributing to **reproductive isolation**.
- **Genomic imprinting** means some genes are expressed only from the maternal or paternal allele, with marks established during gametogenesis.
- **Mitochondria** are almost always **inherited exclusively from the mother**; paternal mitochondria are degraded after fertilization.
- The **sperm centriole** is typically inherited and nucleates the first mitotic spindle.
- The male reproductive system produces sperm (**spermatogenesis**), secretes hormones (**testosterone**), and delivers sperm.
- **Testes** are housed in the **scrotum**, which maintains a temperature **2–3°C below core body temperature** for optimal spermatogenesis.
- Temperature regulation involves the **cremasteric reflex** and the **pampiniform plexus**, a countercurrent heat exchanger.



- **Seminiferous tubules** are the site of sperm production, containing **spermatogenic cells** and **Sertoli cells**.
- **Sertoli cells** provide structural support, form the **blood-testis barrier**, nourish germ cells, and secrete **Androgen-Binding Protein (ABP)** and **inhibin**.
- **Leydig cells**, located in the interstitial tissue between tubules, produce **testosterone** in response to LH.
- Sperm travel from the testes through the **epididymis** (for maturation and storage), **vas deferens**, **ejaculatory duct**, and **urethra**.
- **Accessory glands** (seminal vesicles, prostate, bulbourethral glands) produce **seminal fluid**.
- **Seminal vesicles** contribute ~60-70% of semen volume, providing fructose, prostaglandins, and coagulating proteins.
- The **prostate gland** contributes ~20-30% of semen volume, secreting a milky fluid containing citrate, zinc, and **Prostate-Specific Antigen (PSA)** for semen liquefaction.
- **Bulbourethral glands** produce a clear, alkaline pre-ejaculate that lubricates and neutralizes urine residue.
- **Semen** is a mixture of sperm and seminal plasma, typically 2-5 ml per ejaculate with 20-150 million sperm/ml.
- Hormonal control involves the **HPG axis**: **GnRH** from hypothalamus → **FSH & LH** from pituitary → FSH acts on Sertoli cells, LH acts on Leydig cells → testosterone and spermatogenesis.
- **Testosterone** promotes spermatogenesis, male secondary sex characteristics, libido, and anabolic effects.
- **Negative feedback** by testosterone (on GnRH/LH) and inhibin (from Sertoli cells on FSH) regulates hormone levels.
- The **penis** is the copulatory organ; erection is mediated by parasympathetic release of **nitric oxide (NO)** causing vasodilation.
- The female reproductive system produces ova (**oogenesis**), receives sperm, supports pregnancy, and gives birth.
- **Ovaries** contain **follicles** (each with an oocyte and surrounding somatic cells) and produce **estrogen** and **progesterone**.
- Follicle stages include **primordial**, **primary**, **secondary (antral)**, and **mature (Graafian) follicle**.
- The **oviducts (Fallopian tubes)** are the site of fertilization; the **ampulla** is the most common site.
- The **uterus** has three layers: **perimetrium**, **myometrium** (smooth muscle), and **endometrium**.
- The **endometrium** has a superficial **stratum functionalis** (shed during menstruation) and a deep **stratum basalis** (regenerative).
- The **menstrual cycle** (~28 days) has coordinated **ovarian** and **uterine** phases.
- **Ovarian cycle phases**: **Follicular phase** (days 1-13, follicle growth), **Ovulation** (~day 14), **Luteal phase** (days 15-28, corpus luteum activity).
- **Uterine cycle phases**: **Menstrual phase** (days 1-5, shedding), **Proliferative phase** (days 6-14, estrogen-driven rebuilding), **Secretory phase** (days 15-28, progesterone-driven secretion).
- **Oogenesis** begins prenatally; primary oocytes arrest in **Prophase I** until puberty.
- Each month, one **secondary oocyte** (arrested at **Metaphase II**) is ovulated after the **LH surge**.
- Fertilization triggers completion of **Meiosis II**, producing one **ovum** and a second polar body.
- The **follicular phase** is driven by **FSH** and rising **estradiol**; high estradiol triggers a **positive feedback LH surge**.
- The **luteal phase** is maintained by **progesterone** (and some estradiol) from the **corpus luteum**.
- If no pregnancy occurs, the corpus luteum degenerates, causing a drop in hormones and **menstruation**.
- **Menopause** is the cessation of ovarian cycles due to follicular depletion (~age 45-55).



- The **vagina** is a fibromuscular tube that receives semen and serves as the birth canal.
- The **mammary glands** are modified sweat glands that produce milk; **prolactin** stimulates milk production, **oxytocin** stimulates milk ejection.
- After fertilization in the oviduct, the zygote undergoes **cleavage** to form a **morula**, then a **blastocyst**.
- The **blastocyst** consists of an **inner cell mass** (becomes embryo) and **trophoblast** (becomes placenta).
- **Implantation** into the endometrium occurs around day 7 after fertilization.
- The **placenta** forms from fetal **chorionic villi** and maternal **decidua basalis**, facilitating nutrient/gas/waste exchange and hormone production.
- The placenta produces **Human Chorionic Gonadotropin (hCG)**, which rescues the corpus luteum early in pregnancy.
- **Human prenatal development** is divided into three **trimesters**.
- The **first trimester** (weeks 1-12) is the period of **organogenesis** and is highly susceptible to **teratogens**.
- The **second trimester** (weeks 13-26) involves rapid growth and fetal movement.
- The **third trimester** (weeks 27-40) involves final maturation, especially of lungs and the nervous system.
- **Parturition** (birth) is initiated by a hormonal cascade: fetal cortisol → placental estrogen → increased oxytocin receptors → **oxytocin** release → uterine contractions in a positive feedback loop.
- **Lactation: Prolactin** stimulates milk production; **oxytocin** triggers the milk let-down reflex.
- **Colostrum** is the first milk, rich in antibodies.
- **Infertility** is the failure to conceive after 12 months of unprotected intercourse.
- **Male infertility** causes include **azoospermia** (no sperm), **oligospermia** (low sperm count), duct blockages, hormonal imbalances, and genetic issues.
- **Female infertility** causes include **anovulation** (e.g., PCOS), blocked oviducts (e.g., from PID), **endometriosis**, and uterine fibroids.
- **Assisted Reproductive Technology (ART)** includes **In Vitro Fertilization (IVF)**, **Intracytoplasmic Sperm Injection (ICSI)**, and Gamete Intrafallopian Transfer (GIFT).
- **Miscarriage** is spontaneous pregnancy loss, while **abortion** is induced termination.
- **Sexually Transmitted Diseases (STDs)** include bacterial (gonorrhea, syphilis, chlamydia), viral (HIV, herpes, HPV), and others.
- **HIV** attacks **CD4+ T-cells**, leading to AIDS; managed with Antiretroviral Therapy (ART).
- **HPV** infection with high-risk strains can cause cervical cancer; vaccination is available.
- **Contraception methods** include barrier (condoms), hormonal (pills, implants), intrauterine devices (IUDs), sterilization, and emergency contraception.
- **Sterilization** methods are **vasectomy** (cutting/blocking vas deferens) in males and **tubal ligation** (cutting/blocking oviducts) in females.
- **Prenatal diagnosis** techniques include ultrasound, amniocentesis, Chorionic Villus Sampling (CVS), and Non-Invasive Prenatal Testing (NIPT).
- Ethical issues surround abortion, genetic testing, embryo research, and ART.
- **Asexual reproduction** produces genetically identical offspring from a single parent without gamete fusion.
- Modes include **fission** (binary/transverse), **budding** (external/internal), **fragmentation**, and **parthenogenesis**.
- **Parthenogenesis** is development from an unfertilized egg, common in insects (e.g., bees for drones), some fish, and reptiles.
- Advantages of asexual reproduction: rapid population growth, no mate needed, preserves successful genotypes.



- Disadvantages: low genetic diversity, vulnerability to environmental change and disease, accumulation of deleterious mutations.
- **Sexual reproduction** involves meiosis, gamete formation, and fertilization, increasing genetic diversity.
- **External fertilization** occurs in water (e.g., many fish, frogs, invertebrates); requires synchronization and produces many gametes.
- **Internal fertilization** is common on land (e.g., reptiles, birds, mammals); requires copulatory organs and produces fewer, protected gametes.
- **Hermaphroditism** (both sexes in one individual) can be **simultaneous** (e.g., earthworms) or **sequential** (**protandry**: male first, then female; **protogyny**: female first, then male).
- Based on embryo nourishment and development, strategies are **oviparity** (egg-laying), **ovoviviparity** (egg retained, yolk-nourished), and **viviparity** (live birth, maternally nourished).
- **Oviparity**: Eggs laid externally; embryo nourished by yolk (e.g., most fish, amphibians, reptiles, birds, monotremes).
- **Ovoviviparity**: Eggs retained inside female; embryo nourished by yolk; young born live (e.g., some sharks, snakes).
- **Viviparity**: Young develop within female; nourished directly by mother via a placenta (e.g., most mammals, some sharks/reptiles).
- Reproductive cycles are often seasonal, controlled by hormones and environmental cues (photoperiod, temperature).
- The "**twofold cost of sex**" (males don't produce offspring, only females do) is outweighed by benefits of genetic diversity.
- **Cleavage**: Rapid mitotic divisions without growth, forming a multicellular **blastula** (with a **blastocoel** cavity).
- **Gastrulation**: Cells rearrange to form a three-layered **gastrula** with ectoderm, mesoderm, and endoderm.
- **Organogenesis**: Germ layers differentiate into specific organs and tissues.
- The **blastopore** fate differs: becomes the mouth in **protostomes** and the anus in **deuterostomes**.
- **Neurulation** forms the neural tube from ectoderm, induced by the underlying **notochord**.
- **Neural crest cells** are a migratory cell population from the dorsal neural tube that form diverse structures (nerves, skull bones, pigment cells).
- **Somites** are blocks of mesoderm that form alongside the neural tube and give rise to vertebrae, skeletal muscle, and dermis.
- **Apoptosis** (programmed cell death) sculpts structures by removing cells (e.g., between digits, in neural development).
- **Induction** is the process by which one group of cells influences the development of adjacent cells (e.g., Spemann organizer).
- **Hox genes** are highly conserved transcription factors that control body plan and segment identity along the anterior-posterior axis.
- Limb development is guided by signaling centers: the **Apical Ectodermal Ridge (AER)** secreting FGFs (controls proximal-distal axis) and the **Zone of Polarizing Activity (ZPA)** secreting Sonic hedgehog (**Shh**) (controls anterior-posterior axis).
- **Cell fate specification** can be **cytoplasmic/autonomous** (via maternal determinants), **conditional** (via cell-cell interactions), or **syncytial** (in a multinucleate cell).
- **Evo-Devo** studies how changes in developmental processes and gene regulation lead to evolutionary morphological change.
- **Conserved developmental genes** like **Hox** (body patterning) and **Pax-6** (eye development) are used across animal phyla for similar functions.



MK PREPARATIONS



- Changes in the **timing (heterochrony), location, or amount** of gene expression can lead to major evolutionary innovations.
- Vertebrate reproductive strategies are defined by fertilization site and embryonic nourishment: **oviparity, ovoviviparity, viviparity.**
- The **amniotic egg** (with shell and extraembryonic membranes) was a key innovation for reproduction on land in reptiles, birds, and monotremes.
- **Extraembryonic membranes** in the amniotic egg are the **amnion, chorion, allantois, and yolk sac.**
- In placental mammals, these membranes are modified: the **chorion** forms the fetal part of the placenta, the **allantois** contributes to umbilical circulation, and the **yolk sac** is an early site of blood cell formation.
- Early vertebrate embryos share a conserved **pharyngula stage**, with a notochord, dorsal neural tube, pharyngeal pouches, and post-anal tail.
- **Germ layer derivatives: Ectoderm** → epidermis, nervous system; **Mesoderm** → muscles, bones, circulatory system, kidneys; **Endoderm** → gut lining, liver, pancreas, lungs.

M
K

P
R
E
P
A
R
A
T
I
O
N
S



26. Reproduction and Development

Practice MCQs

1. What is the primary focus of developmental biology?

- A) Study of genetic mutations in adults
- B) Study of how multicellular organisms grow and develop from a zygote
- C) Analysis of evolutionary relationships only
- D) Examination of animal behavior

Answer: Study of how multicellular organisms grow and develop from a zygote

2. The historical concept that a miniature, fully formed organism pre-exists in the gamete is called?

- A) Epigenesis
- B) Homunculus theory
- C) Modern synthesis
- D) Germ layer theory

Answer: Homunculus theory

3. Which of the following is the correct sequence of early developmental stages in animals?

- A) Gastrulation, Cleavage, Fertilization, Organogenesis
- B) Fertilization, Cleavage, Gastrulation, Organogenesis
- C) Cleavage, Fertilization, Organogenesis, Gastrulation
- D) Organogenesis, Gastrulation, Cleavage, Fertilization

Answer: Fertilization, Cleavage, Gastrulation, Organogenesis

4. During gastrulation, which germ layer gives rise to the nervous system and epidermis?

- A) Mesoderm
- B) Endoderm
- C) Ectoderm
- D) Trophoblast

Answer: Ectoderm

5. Von Baer's laws of embryology state that:

- A) Embryos of higher animals pass through adult stages of lower animals
- B) General features appear before specialized features
- C) Embryonic development is identical across all species
- D) All embryos look the same at birth

Answer: General features appear before specialized features

6. Cleavage pattern in mammals is typically:

- A) Meroblastic and discoidal
- B) Holoblastic and isolecithal
- C) Superficial
- D) Meroblastic and superficial

Answer: Holoblastic and isolecithal

7. Which gastrulation movement involves the infolding of a cell sheet?

- A) Involution
- B) Ingression
- C) Delamination
- D) Invagination

Answer: Invagination

8. Programmed cell death that sculpts structures like digits is known as:

- A) Necrosis
- B) Mitosis
- C) Apoptosis
- D) Metastasis

Answer: Apoptosis

9. A fate map is used to:

- A) Determine the genetic sequence of an embryo
- B) Trace which embryonic cells give rise to which adult structures
- C) Map the migration of birds
- D) Identify teratogenic agents

Answer: Trace which embryonic cells give rise to which adult structures

10. The study of how changes in developmental genes drive evolution is called:

- A) Teratology
- B) Evo-Devo
- C) Phylogenetics
- D) Ontogeny

Answer: Evo-Devo

11. An external agent that causes birth defects during critical periods is a:

- A) Mutagen
- B) Carcinogen
- C) Teratogen
- D) Pathogen

Answer: Teratogen

12. In mammals, primary sex determination is triggered by which gene on the Y chromosome?

- A) SOX9
- B) WNT4
- C) SRY

M
K
P
R
E
P
A
R
A
T
I
O
N
S

D) FOXL2

Answer: SRY

13. Which hormone causes the regression of the Müllerian ducts in male fetal development?

- A) Testosterone
- B) Estrogen
- C) Anti-Müllerian Hormone
- D) Follicle-Stimulating Hormone

Answer: Anti-Müllerian Hormone

14. Androgen Insensitivity Syndrome (AIS) results from a mutation in the:

- A) SRY gene
- B) Androgen receptor gene
- C) 5 α -reductase enzyme
- D) Aromatase enzyme

Answer: Androgen receptor gene

15. In Drosophila, sex determination is primarily based on:

- A) Presence of a Y chromosome
- B) Temperature
- C) The X:A ratio
- D) Hormonal signals

Answer: The X:A ratio

16. Environmental Sex Determination (ESD) where low temperature produces males and high produces females is seen in:

- A) Alligators
- B) Some lizards
- C) Turtles (Pattern Ia)
- D) Birds

Answer: Turtles (Pattern Ia)

17. Primordial Germ Cells (PGCs) migrate to the gonadal ridges guided by which chemotactic signal?

- A) SCF
- B) SDF1/CXCR4
- C) GDNF
- D) Retinoic acid

Answer: SDF1/CXCR4

18. Spermatogenesis occurs in the:

- A) Epididymis
- B) Seminiferous tubules
- C) Vas deferens
- D) Prostate gland

Answer: Seminiferous tubules

19. The phase of spermatogenesis where round spermatids transform into spermatozoa is called:

- A) Spermiogenesis

B) Meiosis I

C) Mitotic proliferation

D) Capacitation

Answer: Spermiogenesis

20. What triggers the initiation of meiosis in spermatogenesis?

- A) Luteinizing Hormone
- B) Follicle-Stimulating Hormone
- C) Retinoic acid
- D) Testosterone

Answer: Retinoic acid

21. In oogenesis, the primary oocyte is arrested at which stage until puberty?

- A) Metaphase II
- B) Anaphase I
- C) Prophase I (Dictyate arrest)
- D) Telophase II

Answer: Prophase I (Dictyate arrest)

22. The LH surge directly triggers which event in the ovarian cycle?

- A) Follicle recruitment
- B) Completion of Meiosis II
- C) Ovulation
- D) Formation of the corpus luteum

Answer: Ovulation

23. How many functional ova are produced from one primary oocyte after meiosis?

- A) One
- B) Two
- C) Three
- D) Four

Answer: One

24. The risk of aneuploidy (e.g., Down syndrome) increases with maternal age primarily due to degradation of:

- A) Spindle fibers
- B) Centromeres
- C) Cohesin proteins
- D) Telomeres

Answer: Cohesin proteins

25. The cap-like structure on the sperm head containing digestive enzymes is the:

- A) Flagellum
- B) Acrosome
- C) Midpiece
- D) Nucleus

Answer: Acrosome

26. In mammals, the final maturation step for sperm in the female tract is called:

- A) Spermiogenesis

- B) Capacitation
- C) Acrosome reaction
- D) Spermiation

Answer: Capacitation

27. The fast block to polyspermy involves:

- A) Cortical granule reaction
- B) Zona pellucida hardening
- C) Membrane depolarization
- D) Sperm receptor clipping

Answer: Membrane depolarization

28. Which sperm-specific protein triggers calcium oscillations in the mammalian egg?

- A) Izumo1
- B) Juno
- C) PLC ζ
- D) Bindin

Answer: PLC ζ

29. The fusion of sperm and egg pronuclei is known as:

- A) Cleavage
- B) Gastrulation
- C) Syngamy
- D) Implantation

Answer: Syngamy

30. Mitochondrial DNA inheritance in humans is:

- A) Paternal only
- B) Maternal only
- C) Both paternal and maternal
- D) Random

Answer: Maternal only

31. The scrotum's function is to maintain testicular temperature:

- A) Equal to core body temperature
- B) 2-3°C above core body temperature
- C) 2-3°C below core body temperature
- D) At a variable temperature

Answer: 2-3°C below core body temperature

32. Which cells form the blood-testis barrier and nourish developing germ cells?

- A) Leydig cells
- B) Spermatogonia
- C) Sertoli cells
- D) Theca cells

Answer: Sertoli cells

33. Testosterone is produced by which testicular cells?

- A) Sertoli cells
- B) Spermatocytes
- C) Leydig cells

- D) Germ cells

Answer: Leydig cells

34. The primary energy source for sperm motility in seminal fluid is:

- A) Glucose
- B) Fructose
- C) Citrate
- D) Lactate

Answer: Fructose

35. Which gland produces Prostate-Specific Antigen (PSA) to liquefy semen?

- A) Seminal vesicles
- B) Bulbourethral glands
- C) Prostate gland
- D) Cowper's glands

Answer: Prostate gland

36. Erection of the penis is primarily mediated by the release of:

- A) Adrenaline
- B) Nitric oxide
- C) Acetylcholine
- D) Dopamine

Answer: Nitric oxide

37. In the HPG axis, GnRH is secreted by the:

- A) Anterior pituitary
- B) Hypothalamus
- C) Testes
- D) Adrenal gland

Answer: Hypothalamus

38. Inhibin, produced by Sertoli cells, provides negative feedback specifically on:

- A) LH
- B) FSH
- C) Testosterone
- D) GnRH

Answer: FSH

39. The functional unit of the ovary containing an oocyte and surrounding somatic cells is the:

- A) Corpus luteum
- B) Follicle
- C) Germinal epithelium
- D) Antrum

Answer: Follicle

40. The glycoprotein layer between the oocyte and granulosa cells is the:

- A) Zona pellucida
- B) Corona radiata
- C) Theca interna



D) Cumulus oophorus

Answer: Zona pellucida

41. Fertilization in humans typically occurs in the:

- A) Uterus
- B) Vagina
- C) Ampulla of the oviduct
- D) Cervix

Answer: Ampulla of the oviduct

42. The layer of the uterus that is shed during menstruation is the:

- A) Myometrium
- B) Perimetrium
- C) Stratum basalis
- D) Stratum functionalis

Answer: Stratum functionalis

43. Which hormone surge triggers ovulation?

- A) FSH surge
- B) Estradiol surge
- C) LH surge
- D) Progesterone surge

Answer: LH surge

44. During the menstrual cycle, the secretory phase is driven by which hormone?

- A) Estrogen
- B) Progesterone
- C) FSH
- D) LH

Answer: Progesterone

45. The cessation of ovarian cycles around age 45-55 is called:

- A) Menarche
- B) Menopause
- C) Ovulation
- D) Parturition

Answer: Menopause

46. Milk production is stimulated by the hormone:

- A) Oxytocin
- B) Prolactin
- C) Progesterone
- D) Estrogen

Answer: Prolactin

47. The hollow ball of cells formed after cleavage is the:

- A) Morula
- B) Blastula
- C) Gastrula
- D) Zygote

Answer: Blastula

48. The embryonic structure that implants into the uterine wall is the:

- A) Zygote
- B) Morula
- C) Blastocyst
- D) Trophoblast only

Answer: Blastocyst

49. The hormone produced by the embryo that maintains the corpus luteum is:

- A) LH
- B) FSH
- C) hCG
- D) Progesterone

Answer: hCG

50. Organogenesis occurs primarily during which trimester?

- A) First trimester
- B) Second trimester
- C) Third trimester
- D) Throughout pregnancy

Answer: First trimester

51. Parturition is initiated by a positive feedback loop involving:

- A) Estrogen and FSH
- B) Oxytocin and uterine contractions
- C) Progesterone and relaxin
- D) Prolactin and oxytocin

Answer: Oxytocin and uterine contractions

52. Asexual reproduction that involves an outgrowth from the parent is:

- A) Fission
- B) Budding
- C) Fragmentation
- D) Parthenogenesis

Answer: Budding

53. Development of an embryo from an unfertilized egg is called:

- A) Fission
- B) Budding
- C) Fragmentation
- D) Parthenogenesis

Answer: Parthenogenesis

54. A disadvantage of asexual reproduction is:

- A) Rapid population growth
- B) Low genetic diversity
- C) No need for a mate
- D) Energy efficiency

Answer: Low genetic diversity

55. Internal fertilization is an adaptation primarily for:

- A) Aquatic environments
- B) Terrestrial environments
- C) Asexual reproduction
- D) Producing many gametes

Answer: Terrestrial environments

56. An individual that possesses both male and female reproductive systems is a:

- A) Gonochorist
- B) Hermaphrodite
- C) Parthenote
- D) Zygote

Answer: Hermaphrodite

57. The reproductive strategy where eggs are laid and embryos are nourished by yolk is:

- A) Viviparity
- B) Ovoviviparity
- C) Oviparity
- D) Placental

Answer: Oviparity

58. Which vertebrate group typically exhibits external fertilization?

- A) Birds
- B) Mammals
- C) Most amphibians
- D) Reptiles

Answer: Most amphibians

59. The amniotic egg was a key evolutionary adaptation for:

- A) Aquatic life
- B) Terrestrial reproduction
- C) Internal fertilization only
- D) Asexual reproduction

Answer: Terrestrial reproduction

60. Which extraembryonic membrane is involved in waste storage in birds and reptiles?

- A) Amnion
- B) Chorion
- C) Allantois
- D) Yolk sac

Answer: Allantois

61. In placental mammals, the fetal part of the placenta is derived from the:

- A) Amnion
- B) Chorion
- C) Allantois
- D) Yolk sac

Answer: Chorion

62. During neurulation, the neural tube is formed from which germ layer?

- A) Ectoderm
- B) Mesoderm
- C) Endoderm
- D) Trophoblast

Answer: Ectoderm

63. Somites are derived from which germ layer and give rise to:

- A) Ectoderm; nervous system
- B) Mesoderm; vertebrae and skeletal muscle
- C) Endoderm; gut lining
- D) Mesoderm; dermis only

Answer: Mesoderm; vertebrae and skeletal muscle

64. The process where one group of cells influences the development of adjacent cells is:

- A) Apoptosis
- B) Induction
- C) Migration
- D) Determination

Answer: Induction

65. Hox genes are important for regulating:

- A) Cell division rate
- B) Body plan and segment identity
- C) Gamete formation
- D) Hormone secretion

Answer: Body plan and segment identity

66. In limb development, the Apical Ectodermal Ridge (AER) secretes which signaling molecule?

- A) Sonic hedgehog (Shh)
- B) Wnt7a
- C) Fibroblast Growth Factors (FGFs)
- D) Bone Morphogenetic Proteins (BMPs)

Answer: Fibroblast Growth Factors (FGFs)

67. The zone in the limb bud that controls anterior-posterior patterning is the:

- A) Apical Ectodermal Ridge
- B) Zone of Polarizing Activity
- C) Progress Zone
- D) Notochord

Answer: Zone of Polarizing Activity

68. Protostomes are characterized by:

- A) Blastopore becomes the anus
- B) Radial and indeterminate cleavage
- C) Spiral and determinate cleavage
- D) Coelom formed by enterocoely

Answer: Spiral and determinate cleavage

69. Male infertility characterized by the absence of sperm in semen is:

- A) Oligospermia
- B) Azoospermia
- C) Asthenospermia
- D) Teratospermia

Answer: Azoospermia

70. A common cause of female infertility due to the absence of ovulation is:

- A) Endometriosis
- B) Pelvic Inflammatory Disease
- C) Polycystic Ovary Syndrome
- D) Uterine fibroids

Answer: Polycystic Ovary Syndrome

71. In Vitro Fertilization (IVF) involves:

- A) Fertilization inside the fallopian tube
- B) Fertilization in a laboratory dish
- C) Injection of sperm directly into the uterus
- D) Transfer of gametes into the uterus

Answer: Fertilization in a laboratory dish

72. The sexually transmitted infection caused by Treponema pallidum is:

- A) Gonorrhoea
- B) Syphilis
- C) Genital herpes
- D) Chlamydia

Answer: Syphilis

73. Which viral STD is linked to cervical cancer and has an effective vaccine?

- A) HIV
- B) Herpes Simplex Virus
- C) Human Papillomavirus
- D) Hepatitis B

Answer: Human Papillomavirus

74. A barrier method of contraception that also helps prevent STIs is the:

- A) Oral contraceptive pill
- B) Intrauterine device
- C) Condom
- D) Diaphragm

Answer: Condom

75. Surgical sterilization in females is called:

- A) Vasectomy
- B) Tubal ligation
- C) Hysterectomy
- D) Oophorectomy

Answer: Tubal ligation

76. The process by which a zygote undergoes rapid mitotic divisions is called:

- A) Gastrulation
- B) Cleavage
- C) Organogenesis

- D) Neurulation

Answer: Cleavage

77. The primary germ layer that gives rise to the lining of the digestive tract is:

- A) Ectoderm
- B) Mesoderm
- C) Endoderm
- D) Trophoblast

Answer: Endoderm

78. The notochord is derived from which germ layer?

- A) Ectoderm
- B) Mesoderm
- C) Endoderm
- D) Mesenchyme

Answer: Mesoderm

79. Neural crest cells give rise to all EXCEPT:

- A) Sensory nerves
- B) Adrenal medulla
- C) Epidermis
- D) Craniofacial cartilage

Answer: Epidermis

80. The pharyngeal arches in terrestrial vertebrate embryos are repurposed to form the:

- A) Lungs
- B) Liver
- C) Jaw and ear bones
- D) Kidneys

Answer: Jaw and ear bones

81. Cell fate specification by unevenly distributed maternal determinants is called:

- A) Conditional specification
- B) Syncytial specification
- C) Autonomous specification
- D) Inductive specification

Answer: Autonomous specification

82. Maternal effect genes like bicoid in Drosophila establish the:

- A) Dorsal-ventral axis
- B) Anterior-posterior axis
- C) Left-right axis
- D) Segment boundaries

Answer: Anterior-posterior axis

83. The concept that embryos of different species share a similar early stage is the:

- A) Von Baer's law
- B) Biogenetic law
- C) Phylotypic stage

D) Homunculus theory

Answer: Phylotypic stage

84. Holoblastic cleavage that is unequal due to moderate yolk is characteristic of:

- A) Mammals
- B) Birds
- C) Amphibians
- D) Insects

Answer: Amphibians

85. Epiboly is a gastrulation movement involving:

- A) Cell sheet infolding
- B) Inward rolling of cells
- C) Spreading of cells to enclose deeper layers
- D) Splitting of a sheet into two

Answer: Spreading of cells to enclose deeper layers

86. The synaptonemal complex facilitates:

- A) Cytokinesis
- B) Chromosome condensation
- C) Crossing over during meiosis
- D) Sperm-egg fusion

Answer: Crossing over during meiosis

87. In spermatogenesis, the cells that undergo meiosis I are called:

- A) Spermatogonia
- B) Primary spermatocytes
- C) Secondary spermatocytes
- D) Spermatids

Answer: Primary spermatocytes

88. Spermatids are:

- A) Diploid
- B) Haploid
- C) Tetraploid
- D) Aneuploid

Answer: Haploid

89. The blood-testis barrier is formed by tight junctions between:

- A) Leydig cells
- B) Spermatogonia
- C) Sertoli cells
- D) Endothelial cells

Answer: Sertoli cells

90. Androgen-Binding Protein (ABP) is produced by Sertoli cells to:

- A) Initiate meiosis
- B) Maintain high local testosterone concentration
- C) Stimulate Leydig cells
- D) Break down the blood-testis barrier

Answer: Maintain high local testosterone concentration

91. The first polar body is formed during:

- A) Oogenesis at the end of Meiosis I
- B) Spermatogenesis during Meiosis II
- C) Oogenesis at fertilization
- D) Cleavage of the zygote

Answer: Oogenesis at the end of Meiosis I

92. The corpus luteum secretes:

- A) Estrogen only
- B) Progesterone only
- C) Both estrogen and progesterone
- D) FSH and LH

Answer: Both estrogen and progesterone

93. If fertilization does not occur, the corpus luteum degenerates into a:

- A) Corpus albicans
- B) New follicle
- C) Graafian follicle
- D) Theca interna

Answer: Corpus albicans

94. The proliferative phase of the uterine cycle is driven by:

- A) Progesterone
- B) Estrogen
- C) LH
- D) FSH

Answer: Estrogen

95. Human Chorionic Gonadotropin (hCG) is produced by the:

- A) Corpus luteum
- B) Anterior pituitary
- C) Syncytiotrophoblast of the embryo
- D) Maternal ovary

Answer: Syncytiotrophoblast of the embryo

96. The umbilical cord contains:

- A) One artery and one vein
- B) Two arteries and one vein
- C) Two veins and one artery
- D) Two arteries and two veins

Answer: Two arteries and one vein

97. Colostrum is characterized by being rich in:

- A) Fats
- B) Carbohydrates
- C) Antibodies
- D) Red blood cells

Answer: Antibodies

98. Sequential hermaphroditism where an individual is male first, then female is:

- A) Protogyny
- B) Protandry
- C) Simultaneous hermaphroditism
- D) Gonochorism

Answer: Protandry

99. In birds, sex determination is:

- A) XX female, XY male
- B) ZZ male, ZW female
- C) Temperature-dependent
- D) Based on the X:A ratio

Answer: ZZ male, ZW female

100. The enzyme 5 α -reductase converts testosterone to:

- A) Estradiol
- B) Dihydrotestosterone (DHT)
- C) Androstenedione
- D) Progesterone

Answer: Dihydrotestosterone (DHT)

101. Congenital Adrenal Hyperplasia (CAH) in XX individuals can lead to:

- A) Female internal and external anatomy
- B) Masculinized external genitalia
- C) XY sex reversal
- D) Absence of gonads

Answer: Masculinized external genitalia

102. The Drosophila sex determination gene that acts as the master switch is:

- A) Doublesex
- B) Transformer
- C) Sex-lethal
- D) Sox9

Answer: Sex-lethal

103. In ESD Pattern II (e.g., alligators), what sex is produced at intermediate temperatures?

- A) Male
- B) Female
- C) Both sexes
- D) Hermaphrodite

Answer: Female

104. The primary function of the acrosome reaction is to:

- A) Provide energy for motility
- B) Release enzymes to penetrate egg coats
- C) Initiate DNA replication
- D) Trigger meiosis in the egg

Answer: Release enzymes to penetrate egg coats

105. The Izumo1 protein on sperm interacts with which receptor on the egg?

- A) ZP3
- B) Juno
- C) Bindin receptor
- D) CXCR4

Answer: Juno

106. The slow block to polyspermy in mammals involves modification of the:

- A) Sperm plasma membrane
- B) Zona pellucida
- C) Egg cytoplasm
- D) Cortical actin network

Answer: Zona pellucida

107. Pronuclear migration in the zygote is guided by microtubules nucleated by the:

- A) Maternal centrosome
- B) Sperm centriole
- C) Egg nucleus
- D) Mitochondria

Answer: Sperm centriole

108. Genomic imprinting refers to:

- A) Random mutation of genes
- B) Parent-of-origin specific gene expression
- C) Gene duplication during meiosis
- D) Inactivation of all maternal genes

Answer: Parent-of-origin specific gene expression

109. The pampiniform plexus helps regulate testicular temperature by:

- A) Producing testosterone
- B) Acting as a countercurrent heat exchanger
- C) Storing sperm
- D) Secreting fluid

Answer: Acting as a countercurrent heat exchanger

110. The cremaster muscle is involved in:

- A) Sperm production
- B) Moving the testes closer to or farther from the body
- C) Ejaculation
- D) Secreting seminal fluid

Answer: Moving the testes closer to or farther from the body

111. The vas deferens is cut or blocked during a:

- A) Tubal ligation
- B) Vasectomy
- C) Prostatectomy
- D) Circumcision

Answer: Vasectomy

M
K
P
R
E
P
A
R
A
T
I
O
N
S

112. The bulbourethral glands secrete:

- A) A fructose-rich fluid
- B) An alkaline mucus for lubrication
- C) PSA for liquefaction
- D) Testosterone

Answer: An alkaline mucus for lubrication

113. In the ovarian cycle, the dominant follicle is selected during the:

- A) Luteal phase
- B) Follicular phase
- C) Menstrual phase
- D) Secretory phase

Answer: Follicular phase

114. The fimbriae of the oviduct function to:

- A) Secrete hormones
- B) Capture the ovulated oocyte
- C) Implant the embryo
- D) Produce cervical mucus

Answer: Capture the ovulated oocyte

115. The myometrium is the:

- A) Outer serosal layer of the uterus
- B) Muscular layer of the uterus
- C) Inner lining shed during menstruation
- D) Site of implantation

Answer: Muscular layer of the uterus

116. The labia majora are homologous to the male:

- A) Penis
- B) Scrotum
- C) Prostate gland
- D) Vas deferens

Answer: Scrotum

117. The let-down reflex for milk ejection is stimulated by:

- A) Prolactin
- B) Oxytocin
- C) Estrogen
- D) Progesterone

Answer: Oxytocin

118. The morula stage precedes the formation of the:

- A) Zygote
- B) Blastocyst
- C) Gastrula
- D) Neural tube

Answer: Blastocyst

119. The trophoblast layer of the blastocyst gives rise to the:

- A) Embryo proper
- B) Placenta

C) Yolk sac

D) Amnion

Answer: Placenta

120. The critical period for teratogen susceptibility is highest during:

- A) Fertilization
- B) Cleavage
- C) Organogenesis
- D) Fetal growth

Answer: Organogenesis

121. Fetal cortisol plays a key role in initiating:

- A) Implantation
- B) Organogenesis
- C) Parturition
- D) Lactation

Answer: Parturition

122. Asexual reproduction by splitting of the body into two equal parts is:

- A) Budding
- B) Fission
- C) Fragmentation
- D) Parthenogenesis

Answer: Fission

123. Gemmules are involved in asexual reproduction of some:

- A) Insects
- B) Sponges
- C) Flatworms
- D) Starfish

Answer: Sponges

124. An advantage of sexual reproduction is:

- A) Rapid population growth
- B) Preservation of genotypes
- C) Generation of genetic diversity
- D) Energy efficiency

Answer: Generation of genetic diversity

125. Spermatophores are used for sperm transfer in some:

- A) Mammals
- B) Birds
- C) Arthropods and cephalopods
- D) Amphibians

Answer: Arthropods and cephalopods

126. Ovoviviparity is characterized by:

- A) Eggs laid externally
- B) Live birth with placental nourishment
- C) Eggs retained, yolk-nourished, live birth
- D) No yolk provision

Answer: Eggs retained, yolk-nourished, live birth

127. Amplexus is a mating behavior seen in:

- A) Birds
- B) Frogs
- C) Mammals
- D) Fish

Answer: Frogs

128. The chorioallantoic membrane in bird eggs functions in:

- A) Protection
- B) Nutrition
- C) Gas exchange
- D) Waste storage

Answer: Gas exchange

129. In placental mammals, the yolk sac is an early site for:

- A) Gas exchange
- B) Waste storage
- C) Blood cell formation
- D) Nutrient absorption from mother

Answer: Blood cell formation

130. The heart is derived from which germ layer?

- A) Ectoderm
- B) Mesoderm
- C) Endoderm
- D) Neural crest

Answer: Mesoderm

131. The process of cells changing shape to bend an epithelial sheet is crucial in:

- A) Cleavage
- B) Gastrulation
- C) Neurulation
- D) Apoptosis

Answer: Neurulation

132. The Spemann organizer is a classic example of:

- A) Autonomous specification
- B) Induction
- C) Apoptosis
- D) Morphogen gradient

Answer: Induction

133. Homeotic genes code for:

- A) Growth hormones
- B) Transcription factors
- C) Structural proteins
- D) Receptors

Answer: Transcription factors

134. The T-box gene Tbx5 is important for development of the:

- A) Hindlimb
- B) Forelimb
- C) Heart
- D) Brain

Answer: Forelimb

135. Deuterostomes are characterized by:

- A) Spiral cleavage
- B) Blastopore becomes the mouth
- C) Radial cleavage
- D) Schizocoely

Answer: Radial cleavage

136. Pelvic Inflammatory Disease (PID) is a common cause of female infertility due to:

- A) Anovulation
- B) Blocked fallopian tubes
- C) Uterine fibroids
- D) Endometriosis

Answer: Blocked fallopian tubes

137. Intracytoplasmic Sperm Injection (ICSI) involves:

- A) Mixing sperm and egg in a dish
- B) Placing sperm in the uterus
- C) Injecting a single sperm into an oocyte
- D) Transferring an embryo to the uterus

Answer: Injecting a single sperm into an oocyte

138. The bacterial STD caused by Neisseria gonorrhoeae is:

- A) Syphilis
- B) Gonorrhea
- C) Chlamydia
- D) Herpes

Answer: Gonorrhea

139. Antiretroviral Therapy (ART) is used to manage:

- A) HPV
- B) HIV/AIDS
- C) Herpes
- D) Syphilis

Answer: HIV/AIDS

140. Emergency contraception primarily works by:

- A) Causing abortion of an implanted embryo
- B) Preventing fertilization or ovulation
- C) Killing sperm
- D) Blocking implantation permanently

Answer: Preventing fertilization or ovulation

PPSC, FPSC, SPSC, KPPSC, AJKPSC & BPS

LECTURER BIOLOGY & ZOOLOGY MOST REPEATED & PAST PAPERS MCQs

1. The amphibians and the _____ represent the two major branches of the tetrapod lineage.

- A) reptiles
- B) birds
- C) tennospondyls
- D) amniotes

Answer: amniotes

2. Which of the following statements about hair is true?

- A) it is homologous to feathers
- B) it is a dermal structure
- C) it is found in all mammals
- D) it is possessed by all endotherms

Answer: it is found in all mammals

3. The endoskeleton of the echinoderms is composed of.

- A) keratin ossicles
- B) calcareous ossicles
- C) keratin spicules
- D) calcareous spicules

Answer: calcareous ossicles

4. The first hormone that was isolated.

- A) thyroxine
- B) vasopressin
- C) secretin
- D) adrenaline

Answer: secretin

5. The ultimate source of organic variations is.

- A) sexual reproduction
- B) hormonal action
- C) natural selection
- D) mutation

Answer: mutation

6. The structure of molluscs that forms the shell and houses the gills is the.

- A) mantle
- B) epidermis
- C) gastrovascular cavity
- D) odontophore

Answer: mantle

7. In an ecosystem, which one shows one way passage?

- A) nitrogen
- B) carbon
- C) potassium
- D) free energy

Answer: free energy

8. The central fluid filled cavity of the blastula is known as.

- A) archenteron
- B) blastocoel

C) blastocyst

D) morula

Answer: blastocoel

9. The mammalian skull may be characterized as

_____.

- A) anapsid
- B) diapsid
- C) biaspid
- D) synapsid

Answer: synapsid

10. The first terrestrial tetrapods were probably the.

- A) rhipidistians
- B) amniotes
- C) coelacanth
- D) leptocephalans

Answer: rhipidistians

11. Which of the following phyla of animals is exclusively marine?

- A) protozoa
- B) porifera
- C) echinodermata
- D) mollusca

Answer: echinodermata

12. A normal green male Maize is crossed with albino female. The progeny is albino because.

- A) trait for albinism is dominant
- B) the albinos have biochemical to destroy plastids derived from green male
- C) plastids are inherited from female parent
- D) green plastids of male must have mutated

Answer: plastids are inherited from female parent

13. Sertoli cells are found in.

- A) liver
- B) seminiferous tubules
- C) pancreas
- D) gut

Answer: seminiferous tubules

14. The sum total of the populations of the same kind of organisms constitute.

- A) colony
- B) genus
- C) species
- D) community

Answer: species

15. A molluscan shell is made of three layers arranged from the outside to the inside.

- A) periostracum, prismatic layer, nacreous layer
- B) mantle layer, prismatic layer, periostracum
- C) nacreous layer, periostracum, mantle layer
- D) none of these



M

K

P

R

E

P

A

R

A

T

I

O

N

S



Answer: periostracum, prismatic layer, nacreous layer

16. Which one of the following can achieve protection and conservation of biodiversity?

- A) eco-development
- B) biosphere reserve
- C) national park
- D) game sanctuary

Answer: biosphere reserve

17. Lamarck's "Theory of Organic Evolution" is based upon.

- A) effect of environment
- B) use and disuse of body parts
- C) inheritance of acquired characters
- D) all of these

Answer: all of these

18. Myxoedema in adults is caused due to.

- A) hyperthyroidism
- B) deficiency of thyroid hormone
- C) overproduction of PTH
- D) deficiency of PTH

Answer: deficiency of thyroid hormone

19. "The animals of colder countries have small ears, short tail and limbs to avoid more loss of heat" is the theme of.

- A) Allen's law
- B) Dollo's law
- C) Gause's law
- D) Cope's Law

Answer: Allen's law

20. Testosterone is secreted by.

- A) sertoli cells
- B) leydig cells
- C) spermatocyte
- D) histiocyte

Answer: leydig cells

21. Changing from a bilaterally symmetrical larval form to a radially symmetrical adult involves relocation of various parts. In this process, the left side becomes the _____ and the right side becomes the _____.

- A) aboral surface, oral surface
- B) anterior surface, posterior surface
- C) dorsal surface, ventral surface
- D) oral surface, aboral surface

Answer: oral surface, aboral surface

22. The class name Myxini, or the hagfishes, refers to their.

- A) lack of eyes
- B) unique circulatory system
- C) production of slime
- D) parasitic lifestyle

Answer: production of slime

23. The earliest therians (mammals) evolved in the _____.

- A) Permian
- B) Triassic
- C) Cretaceous
- D) Cenozoic

Answer: Triassic

24. The development of limbs probably aided the first amphibians in.

- A) finding mates
- B) swimming
- C) moving between bodies of water
- D) none of these

Answer: moving between bodies of water

25. Which of the following is odd one?

- A) cockroach, spider, silver-fish
- B) whale, bat, lizard
- C) star-fish, sea-cucumber, sea-urchin
- D) cray-fish, cuttle-fish hag-fish

Answer: cray-fish, cuttle-fish hag-fish

26. ABO blood group system is due to.

- A) multifactor inheritance
- B) incomplete dominance
- C) multiple allelism
- D) epistasis

Answer: multiple allelism

27. In human beings, the eggs are.

- A) microlecithal
- B) macrolecithal
- C) mesolecithal
- D) electihal

Answer: mesolecithal

28. Pyramid of numbers deal with number of.

- A) species in area
- B) sub-species in a community
- C) individuals in a community
- D) individuals in a trophic level

Answer: individuals in a trophic level

29. The morphogenetic movement change the hollow spherical blastula into a/an.

- A) embryonic disc
- B) gastrula
- C) morula
- D) neurula

Answer: gastrula

30. The dominant second trophic level, in a lake ecosystem, is.

- A) benthos
- B) plankton
- C) zooplankton
- D) phytoplankton

Answer: zooplankton



M

K

P

R

E

P

A

R

A

T

I

O

N

S



D) Uracil

Answer: Thymine

172. Genetic dominance is associated with.

- A) Personality dominance
- B) Population distribution
- C) Adaptive value
- D) None of these

Answer: Population distribution

173. Amino acids are organic compounds behaving like.

- A) Acid
- B) Base
- C) Both a & b
- D) None of these

Answer: Both a & b

174. XXY individual in Drosophila (fruit fly) is.

- A) Male
- B) Female
- C) Sterile
- D) All of these

Answer: Female

175. Linneous proposed the method of naming organism.

- A) Uninominal
- B) Trinomial
- C) Binomial
- D) None of these

Answer: Binomial

176. Ontogeny represents the history of an organisms.

- A) Embryonic
- B) Both a & c
- C) Revolutionary
- D) None of these

Answer: Embryonic

177. The white blood corpuscles of the cockroach are called as.

- A) Hemocyte
- B) Haemoblast
- C) Haematoclast
- D) Leucocyte

Answer: Hemocyte

178. Nuclear membranes have a structure similar to.

- A) Cell membrane
- B) Endoplasmic reticulum
- C) Mitochondria
- D) All of these

Answer: All of these

179. Trypsin protease secreted in.

- A) Bile
- B) Saliva
- C) Gastric juice

D) Pancreatic juice

Answer: Pancreatic juice

180. Insects mouth Parts are serially homologous.

- A) Claspers
- B) Sting apparatus
- C) Legs
- D) None of these

Answer: Legs

181. At cytokinesis in plants a structure phragmoplast is formed from vesicles which originate from?

- A) Lysosomes
- B) Golgi complex
- C) Centriole
- D) Glyoxisomes

Answer: Golgi complex

182. Sickle cell anemia is the condition where RBCs deflate.

- A) Under oxygen
- B) Both a & c
- C) Absence of Oxygen
- D) None of these

Answer: Absence of Oxygen

183. Inherited and environmental factors responsible for certain trait are termed as.

- A) Epigenetics
- B) Multifactorial
- C) Monogenetic mutation
- D) None of these

Answer: Epigenetics

184. DNA duplicate during.

- A) Prophase
- B) Metaphase
- C) Telophase
- D) Interphase

Answer: Interphase

185. The first living and respiring organisms on earth were.

- A) Aerobes
- B) Anaerobes
- C) Both a & b
- D) None of these

Answer: Anaerobes

186. Zygote containing XY are female in.

- A) Drosophila
- B) In grasshopper
- C) In moths
- D) None of these

Answer: None of these

187. Significant flight muscles in birds is.

- A) Appendicular
- B) Tensor
- C) Pectoral



M

K

P

R

E

P

A

R

A

T

I

O

N

S



- C) Lecithin
- D) Thrombin

Answer: Thrombin

392. During the final stage of cell division, the mitotic apparatus disappears, the chromosomes become attenuated, the centrioles duplicate and split, the nuclear membrane becomes reconstituted and the nucleolus reappears. This phase of cell division is known as.

- A) Prophase
- B) Metaphase
- C) Anaphase
- D) Telophase

Answer: Telophase

393. Sickle cell anemia and Huntington's chorea are both.

- A) Virus-related diseases
- B) Bacteria-related diseases
- C) Congenital disorders
- D) None of the above

Answer: Congenital disorders

394. Which of the following is NOT part of a neuron?

- A) Synapse
- B) Axon
- C) Nissl bodies
- D) Dendrite

Answer: Synapse

395. Melatonin is produced by the.

- A) Skin
- B) Pineal gland
- C) Liver
- D) Pituitary gland

Answer: Pineal gland

396. Which of the following statements is TRUE of insulin? Is it.

- A) Secreted by the pancreas
- B) A protein
- C) Involved in the metabolism of glucose
- D) All of the above

Answer: All of the above

397. Bacteriophage are.

- A) Bacteria
- B) Bacteria precursors
- C) Viruses
- D) Agents which cause the production of bacteria

Answer: Viruses

398. The AIDS viruses specifically attack which kind of cell?

- A) B cells
- B) Helper T cells
- C) Killer T cells

- D) Macrophages

Answer: Helper T cells

399. Which of the following human digestive enzymes is incorrectly matched to its substrate?

- A) Pepsin - protein
- B) Trypsin - starch
- C) Pancreatic amylase - starch
- D) Lipase -- fat

Answer: Trypsin - starch

400. Humans cannot digest cellulose because.

- A) It does not contain sugars
- B) It is made up of disaccharides
- C) It is made up of monosaccharides
- D) Humans lack the proper enzymes

Answer: Humans lack the proper enzymes

401. Oxidation of fats and carbohydrates within a cell would be an example of.

- A) Anabolism
- B) Catabolism
- C) Biosynthesis
- D) None of the above

Answer: Catabolism

402. The ideal vector system would have following characteristics.

- A) Non detectable by immune system
- B) Adequate carrying capacity
- C) Non inflammatory
- D) All

Answer: All

403. Hollow fat molecule in solution is called.

- A) Polysome
- B) Polyester
- C) Lipids
- D) Liposomes

Answer: Liposomes

404. Gene therapy can be used for.

- A) Alzheimer's disease
- B) Huntington's disease
- C) Parkinson's disease
- D) All

Answer: All

405. Nonhuman Primates similarities with human.

- A) Genetic
- B) Physiologic
- C) Anatomic
- D) All of the above

Answer: All of the above

406. Vaccinia virus is isolated from.

- A) Herpes Virus
- B) Pox virus
- C) Adenovirus
- D) None

Answer: Pox virus



M
K

P

R

E

P

A

R

A

T

I

O

N

S

